system. Overt disease might then follow from impaired suppressor T cell function and the deposition of immune complexes in the liver and lung.

FRANCIS J DUNNE

Senior Registrar in Psychiatry,
St Bartholomew's Hospital,
London EC1A 7BE

- 1 Conn HO, Fessel JM. Spontaneous bacterial peritonitis in cirrhosis. Variations on a theme. *Medicine* 1971;50:161-97.
- 2 Watson RR, McMurray DN. The effects of malnutrition on the secretory and cellular immune processes. *CRC Crit Rev Food Sci Nutr* 1979;12:113-59.
- 3 MacGregor RR. Alcohol and immune defense. *JAMA* 1986;256:1474-9.
- 4 Jerrells TR, Marietta CA, Bone G, Weight FF, Eckardt MJ. Ethanol-associated immunosuppression. *Adv Biochem Psychopharmacol* 1988;44:173-85.
- 5 Watson RR, Jackson JC, Hartmann B, Sampliner R, Mobley D, Eskelson C. Cellular immune functions, endorphins, and alcohol consumption in males. *Alcoholism: Clinical and Experimental Research* 1985;9:248-54.
- 6 Liu YK. Effects of alcohol on granulocytes and lymphocytes. *Semin Hematol* 1980;17:130-6.

- 7 Glassman AB, Bennett CE, Randall CL. Effects of ethyl alcohol on human peripheral lymphocytes. *Arch Pathol Lab Med* 1985;109:540-2.
- 8 Lundy J, Raaf JH, Deakens S, et al. The acute and chronic effects of alcohol on the human immune system. *Surg Gynecol Obstet* 1975;141:212-8.
- 9 Drew PA, Clifton PM, LaBrooy JT, Sherman DJC. Polyclonal B cell activation in alcoholic patients with no evidence of liver dysfunction. *Clin Exp Immunol* 1984;57:479-86.
- 10 Roselle GA, Mendenhall CL. Alteration of in vitro human lymphocyte function by ethanol, acetaldehyde, and acetate. *J Clin Lab Immunol* 1982;9:33-7.
- 11 Ristow SS, Starkey JR, Hass GM. Inhibition of natural killer cell activity in vitro by alcohols. *Biochem Biophys Res Commun* 1982;105:1315-21.
- 12 Kay N, Allen J, Morley JE. Endorphins stimulate normal human peripheral blood lymphocyte natural killer activity. *Life Sci* 1984;35:53-9.
- 13 Tewari S, Noble EP. Ethanol and brain protein synthesis. *Brain Res* 1971;26:469-79.
- 14 MacGregor RR, Gluckman SJ, Senior JR. Granulocyte function and levels of immunoglobulins and complement in patients admitted for withdrawal from alcohol. *J Infect Dis* 1978;138:747-53.
- 15 Rimland D. Mechanisms of ethanol-induced defects of alveolar macrophage function. *Alcoholism* 1983;8:73-6.
- 16 Smith FE, Palmer DL. Alcoholism, infection, and altered host defenses. A review of clinical and experimental observations. *J Chronic Dis* 1976;29:35-49.
- 17 Stacey NH. Inhibition of antibody-dependent cell-mediated cytotoxicity by ethanol. *Immunopharmacology* 1984;8:155-61.
- 18 Draper LR, Gyure LA, Hall JG, Robertson D. Effect of alcohol on the integrity of the intestinal epithelium. *Gut* 1983;24:399-404.

Being bullish about medical research

The need to accept concentration

Deep gloom has been the prevailing mood among medical researchers for a long time. In some at least it is lifting. The government's policies of selection, concentration, and exploitation have created some winners as well as some losers. The winners are those who have taken up the tremendous opportunities offered by the new biology, reorganised themselves into larger—often interdisciplinary—units, done well in the selection exercises, accepted life on short term contracts confident that they will be good enough to win more, and recognised the importance of seeking funds not only from governmental sources but also from charities and industry. The losers are those left in shrinking, old fashioned departments, overwhelmed by teaching and service commitments, and finding it impossible to do innovative research: the new elitism has left them behind. Today's report from the Academic Medicine Group (p 573) contains the voices of both winners and losers, making it somewhat bland and contradictory. We believe the painful consequences of concentration must be accepted.

The good news about medical research has not received as much attention as it should—perhaps because the losers have understandably been shouting louder than the winners. But there is good news. The arguments over the new National Centre for Clinical Research have finally been settled, and the government has in the past month agreed to pay £2m for site development at Hammersmith. More money is almost certain to follow, and it is crucial to the future of clinical research in Britain that the centre be made to work. But the centre is accepting that it cannot go it alone and must cooperate fully with other organisations, including the British Postgraduate Medical Federation and the undergraduate medical schools in London. The federation is now working hard to link together London's unique collection of research institutions and specialist postgraduate hospitals, and the University of London is looking at further linking its constituent parts. Meanwhile, great intellectual excitement is being generated by the Medical Research Council's directed research programme on AIDS, and basic and clinical scientists from different disciplines are being brought together in a way that may serve as a model for attacking other medical problems.

Also in the past month extra government money has been given to fund interdisciplinary research centres (p 550): the

Medical Research Council is now funding four—in cell biology, toxicology, protein engineering, and molecular science—and three more are planned—in neurodegeneration, the brain and behaviour, and molecular genetics. There is strong theoretical and empirical evidence to support the thinking behind these interdisciplinary research centres,^{1,2} and they will probably proliferate.

There is also good news on the horizon for training. Although the funding of intercalated BSc degrees is being squeezed, the first British MB-PhD programme is likely to begin next October—at Cambridge University, and others are likely to follow. In these programmes undergraduates train in basic science and medicine simultaneously, thus preparing them to be the people that link the two disciplines. Such programmes started in 1964 in the United States and have expanded rapidly till there are now around 1000 students taking the courses. It may be that money is better invested in these programmes than in the shorter intercalated BSc degrees, although fewer medical schools will be able to offer the more intensive degrees. Certainly, MB-PhD degrees will probably bring richer returns than MD degrees, which are often undertaken unwillingly by doctors with little interest in research. Many MD degrees result in either no publications or none that are worthwhile; often they are not completed.³ There has been too much "Mickey Mouse" research undertaken in British medical schools.

The bad news for British medical research is less about money⁴ and more about manpower. Although the data are sketchy, most senior researchers believe that fewer bright young graduates are entering careers in research than in the past, being deterred by the inadequate facilities, the salaries, and the poor career prospects. Medical graduates in particular recognise that an NHS career combined with private practice offers much richer rewards and more stability than a career in academic medicine. The Academic Medicine Group is thus right to support the call for 250 new career posts in academic medicine, but the posts need to be concentrated. The manpower proposals of *Achieving a Balance* also threaten to make it more difficult for young doctors to combine research and service work, which may not be bad if it separates those who are serious about research from those who are doing it simply to decorate their curriculum vitae.⁵ The government

seems to have paid little attention to research when framing the manpower proposals, and similarly the NHS review devotes only a non-committal paragraph to research.⁶ Yet the opting out of teaching hospitals, the high costs of providing treatment in research oriented hospitals, and the possibility of paying doctors at local rates may all undermine clinical research. It is also unfortunate that the government has not replied to the important report from the House of Lords on medical research despite its publication a year ago.⁷

But the issue that neither the House of Lords nor the Academic Medicine Group has considered adequately is the problem of concentration of medical research in Britain. The group has accepted the principles of concentration, selectivity, priority setting, and better management of the resources needed for research but has not perhaps accepted the consequences. Thus the group says: "Internationally competitive academic clinical departments will need a broad range of skills including top class clinical investigation combined with advanced basic science. There is little sign that many departments can meet these demands." It says too: "The size of most British academic departments now is often inadequate to meet the demands of modern clinical education and research." Yet the group rejects the three tier system of universities put forward by the Advisory Board for the Research Councils (type R universities doing internationally competitive research across the board, type X doing it in selected subjects, and type T doing only a small amount of research needed to support teaching); and the group's first recommendation is that "teaching must take place in schools where high quality research is in progress."

The nettle has not been grasped. It is impossible that all of Britain's 28 undergraduate medical schools will be able to conduct internationally competitive research; indeed, the skimpy evidence available is that it is a fiction that all schools are producing such research at the moment.⁸ Preliminary data suggest, for instance, that only four medical schools in Britain produce more professors and readers than they consume (J Anderson, personal communication). We may be close to an R, T, X system already, and it may be dogma that medical students have to be taught in an environment where high quality research is in progress. Many probably are not, certainly in most subjects. The advisory board argued—from anecdotal evidence in the United States—that students may be better taught in institutions in which teaching rather than research is the priority.² The British medical research and education community has not yet faced this difficult question head on—but it needs to.

RICHARD SMITH

Assistant editor, *BMJ*

- 1 Irvine J, Martin B. *Foresight in science: picking the winners*. London: Pinter, 1984.
- 2 Advisory Board for the Research Councils. *A strategy for the science base*. London: ABRC, 1987.
- 3 Wood GM, Catford JC. Research post-MRCP. *J R Coll Phys Lond* 1986;20:89-94.
- 4 Smith R. The funding of medical research: going up or coming down? *Br Med J* 1988;296:267-70.
- 5 Department of Health and Social Security, Joint Consultants' Committee, Chairmen of Regional Health Authorities. *Hospital medical staffing: achieving a balance*. London: DHSS, 1987.
- 6 Secretaries of State for Health, Wales, Northern Ireland, and Scotland. *Working for patients*. London: HMSO, 1989. (Cmnd 555.)
- 7 House of Lords Select Committee on Science and Technology. *Priorities in medical research*. London: HMSO, 1988.
- 8 Wakeford R, Adams R. Medical research output 1973-81: a romp around the United Kingdom research centres. *Br Med J* 1984;288:297-9.

Screening: the inadequacy of population registers

Screening will fail if population registers are not improved

The current enthusiasm for screening may be stymied by the inadequacies of population registers. The registers being used by district health authorities to screen for cervical cancer are the name and address lists of general practitioners' patients held by family practitioner committees. These will also be used for the national breast screening programme and are sometimes used by health authorities for screening the elderly. The information available from these lists may not, however, be of sufficient quality to deliver these services.

A recent survey of the health of all people aged 85 and over living in City and Hackney Health Authority found that the family practitioner committee's list was less accurate than expected.¹ According to estimates from the Office of Population Censuses and Surveys and the London Research Centre the number of people aged 85 and over living at home in City and Hackney in 1986 was between 1400 and 1700. Yet the family practitioner committee's list contained the names and addresses of 3018 people within this category. Validity checks—with the electoral roll, postal screening questionnaires, and household checks by interviewers—found that this sample was inflated with the names of people who had died or moved. Two thirds of the 3018 addresses from the family practitioner committee's list were inaccurate. Preliminary checks made in 1988 on the accuracy of the addresses of younger people held by the family practitioner committee showed that half of those for people aged 65-74 and two fifths for people aged 75-84 were inaccurate.

The inaccuracy rate of family practitioner committees' lists increases with the age of the population, partly due to increased mortality. These findings are particularly relevant for screening programmes for the elderly and for attempts to screen older women for breast and cervical cancer. Inaccuracy rates for younger people are also unacceptably high. Invitations for cervical cytology screening are generally sent to women aged 35-65 (in some districts the minimum age is lower). Research has shown that between an eighth and a third of non-responders to postal invitations had moved from the address listed on family practitioner committees' registers (or in a few cases had died) and therefore did not receive their invitations.^{2,6}

Women in the lower socioeconomic classes, who are at the highest risk of cervical cancer, are likely to be overrepresented among those who do not receive an invitation for recall because of moving house.³ One study in a deprived inner city area showed that over half of non-attendances for cervical screening were caused by administrative errors in the family practitioner committee's call-recall system; prominent among the reasons for these were incorrect addresses.⁷ This problem is worse in inner city areas because of their high population turnovers.⁸

About three million people each year change their general practitioners in England and Wales. Although three quarters will have registered with a new general practitioner within three months,⁷ problems arise with patients who change their