specific recommendations for a uniform policy of infection control (*BTS Newsletter*, winter 1988).

We thank Keymed, Surgikos, and the Sanatorium Society for their support.

1 Department of Health and Social Security. Disinfection of endoscopes potentially contaminated by mycobacterium species. *Safety Information Bulletin* 1986;28.

2 Working Party of British Society of Gastroenterologists. Cleaning and disinfection of equipment for gastrointestinal flexible endoscopy. Interim recommendations. *Gut* 1988;29:1134-51.

3 Centers for Disease Control. Recommendations for prevention of HIV transmission in health care settings. *MMWR* 1987;36:25.

4 Hanson PJV, Jeffries DJ, Batten JC, Collins JV. Infection control revisited: dilemma facing today's bronchoscopists. *Br Med J* 1988;297:185-7.

(Accepted 7 February 1989)

Carpal tunnel syndrome and type of dialysis membrane

J Chanard, P Bindi, S Lavaud, O Toupance, H Maheut, F Lacour

Service de Néphrologie,
Centre Hospitalier et
Universitaire, Hôpital
Maison Blanche, 51092
Reims Cédex, France
J Chanard, MD, professor of
nephrology
P Bindi, MD, resident
S Lavaud, MD, hospital
practitioner
O Toupance, MD, hospital
practitioner

Clinique Chirurgicale
Mutualiste, 51100 Reims
H Maheut, MD, consultant
nephrologist
F Lacour, MD, consultant
nephrologist

Correspondence to:
Professor Chanard.

Br Med J 1989;298:867-8

The carpal tunnel syndrome is commonly seen in patients who have received haemodialysis for a long time.^{1,2} Compression of the median nerve, flexor tenosynovitis, and destructive arthropathies are associated with hypertrophic material, which generally contains amyloid derived from β_2 microglobulin. The pathogenesis of the syndrome is unknown, although possibly the dialysis membrane has an effect. The syndrome has been reported in patients treated with cellulose membranes non-permeable to β_2 microglobulin. The prevalence of the syndrome increases with the duration of dialysis, and nearly all patients treated for more than 18 years require a decompression operation.

Few data have been reported on the occurrence of amyloidosis due to dialysis in patients treated with synthetic membranes permeable to β_2 microglobulin.¹ Since the early 1970s we have used the highly permeable polyacrylonitrile AN 69 membrane for dialysis. We report the prevalence of the syndrome in our patients.

Patients, methods, and results

Eighty five patients who had received dialysis for more than seven years were entered into the study. Two groups were selected: group 1 comprised 31 patients who had been treated exclusively with AN 69, and group 2 comprised 54 patients who had been treated with cuprophane for an average of 5.7 (SD 3.9) years and subsequently with AN 69. The mean duration of dialysis was 124.2 (33.0) months in group 1 and 126.9 (28.6) months in group 2. The mean time for which patients had been exposed to the artificial membrane was 5533 (1200) hours and 7580 (1984) hours respectively ($p < 0.05$). The mean age of the groups was 54 (3) and 49 (14), and the mean serum β_2 microglobulin concentration was 49 (15) mg/l and 45 (13) mg/l.

The carpal tunnel syndrome was considered to be present only when a decompression operation was needed, thus avoiding clinical bias by using motor nerve conduction velocity as reduced velocity may be due to uraemic toxicity without entrapment of the median nerve. After 10 years of dialysis only one patient in group 1 had had the operation, compared with 16 in group 2 ($p < 0.001$). Amyloidosis derived from β_2 microglobulin was proved in the only patient in group 1 with the syndrome and in eight out of 12 patients in group 2 from whom synovial specimens were available.

Cumulative survival analysis with the Mantel-Cox model applied to the syndrome at the time of operation showed a significant difference ($p = 0.0118$) between the two techniques of dialysis. The proportion of

patients without the syndrome after 10 years was predicted to be 94% ($n = 18$) in those treated with AN 69 and 77% ($n = 21$) in those treated with several membranes; that at 12 years was predicted to be 94% ($n = 8$) and 61% ($n = 11$) respectively.

Comment

The carpal tunnel syndrome was less common in patients treated exclusively with the highly permeable membrane AN 69 than in patients who had been exposed to cuprophane for at least five years. Preliminary studies have indicated a similar trend for juxta-articular cystic bone defects, which have been assumed to be related to amyloidosis derived from β_2 microglobulin.³ Except for the type of membrane used and the time for which patients had been exposed to the membrane there were no clinical or biological differences between the two groups.

Cumulative percentage of patients without carpal tunnel syndrome, calculated from life tables and survival analysis, in patients given dialysis with AN 69 membrane (—) and with several membranes including cellulose membrane (---). Figures are numbers of patients. Date of onset of carpal tunnel syndrome was considered to be date of surgery.

There are several ways in which AN 69 might prevent or postpone the development of the carpal tunnel syndrome. The concentration of β_2 microglobulin substrate may be insufficient to allow amyloid to form after 10 years of dialysis. With a sieving coefficient of 0.40 for β_2 microglobulin up to about 22.5 g of the protein is extracted each year,⁴ but despite this serum β_2 microglobulin concentrations remain raised and it would be naive to assume that the serum β_2 microglobulin concentration alone causes amyloidosis in the absence of other promoting factors such as changes in the structure of the precursor protein in the serum⁵ and transformation of the precursor protein in situ by some local process. Alternatively, because AN 69 is more biocompatible than cellulose mem-

branes it may induce less acute phase reaction. This could play a critical part in preventing or diminishing reactions that favour polymerisation of β_2 microglobulin and generation of amyloid.

- 1 Bardin T, Zingraff J, Kuntz D, Drücke T. Dialysis-related amyloidosis. *Nephrol Dial Transplant* 1986;1:151-4.
- 2 Brown EA, Arnold IR, Gower PE. Dialysis arthropathy: a complication of long-term treatment with haemodialysis. *Br Med J* 1986;292:163-6.

- 3 Vandembroucke JM, Jadoul M, Maldague B, Huaux JP, Noel H, Van Ypersele de Strihou C. Possible role of dialysis membranes characteristics in amyloid osteoarthropathy. *Lancet* 1986;i:1210-1.
- 4 Chanard J, Lavaud S, Toupance O, Melin JP, Gillery P, Revillard JP. β_2 -Microglobulin associated amyloidosis in chronic haemodialysis patients. *Lancet* 1986;i:1212.
- 5 Ogawa H, Saito A, Oda O, Nakajima M, Chung TG. Detection of novel β_2 -microglobulin in the serum of hemodialysis patients and its amyloidogenic predisposition. *Clin Nephrol* 1988;30:158-63.

(Accepted 16 December 1988)

Rat bites and diabetic foot in the West Indies

P Cooles, H Paul

Princess Margaret
Hospital, Roseau,
Dominica, West Indies
P Cooles, MRCP, consultant
physician
H Paul, FRCS, consultant
surgeon

Correspondence to: Dr
Cooles.

Br Med J 1989;298:868

Diabetes mellitus, a disease associated with affluence, is an increasing problem in the West Indies.¹ Diabetic foot is an important cause of morbidity and is often complicated by rat bites. We present four cases of rat bites on a diabetic foot.

Case reports

The table summarises details of the cases.

Details of four cases of rat bites on a diabetic foot

Case No	Age (years)	Sex	Peripheral neuropathy	Foot pulses	Length of hospital stay (days)	Outcome for foot
1	36	M	Yes	Yes	23	Saved
2	70	F	Yes	No	30	Amputated
3	73	F	Yes	No	7	Saved
4	50	M	Yes	Weak	27	Saved

Case 1—An insulin dependent diabetic woke one morning to see a rat eating his left hallux and second toe. He could not feel the bites. Diabetes had been diagnosed four years earlier and peripheral neuropathy one year earlier on the basis of impaired sensation below the knees, absent ankle and knee jerks, and impotence. After two weeks of care by the village nurse he was referred to hospital. Culture of material from a wound swab grew a β haemolytic streptococcus. He was treated with oral cloxacillin and metronidazole, and the wound was cleaned daily. The wound healed satisfactorily, and he was discharged taking 80 units isophane insulin daily in two doses.

Case 2—A woman presented one week after waking with an extensive rat bite on her left hallux. She was not known to be diabetic but had noticed polyuria for several months. Her blood glucose concentration on admission was 19 mmol/l. Her toe was gangrenous, and there were no palpable foot pulses. Sensation in both feet was severely impaired, and both ankle and knee jerks were absent. Culture of material from a wound swab grew a proteus. Amputation below the knee was performed 10 days after admission, and she was discharged taking chlorpropamide 250 mg daily.

Case 3—A woman awoke with a large rat bite on her left hallux. She had had diabetes for 13 years, which

had been controlled with oral chlorpropamide. Amputation below the right knee had been performed two years earlier. After two weeks of care by the village nurse she was sent to hospital. Foot pulses and ankle and knee jerks were absent. Sensation was severely impaired below the knee. Material from a wound swab was sterile. When she was discharged the wound was healing satisfactorily with daily cleaning and chlorpropamide 250 mg daily.

Case 4—A man was referred two weeks after waking to find his feet bleeding from multiple painless rat bites. He had been diabetic for 13 years but had not attended a clinic for over a year. He had absent knee and ankle jerks, impaired sensation in his feet, and weak foot pulses. Material from wound swabs was sterile. His wounds healed with conservative management, and he was discharged taking 30 units insulin zinc suspension daily.

Comment

Rat bites can have serious consequences and even cause death.² Clinton estimated that there were at least 14 000 rat bites a year in the United States,² and Deoras reported 20 000 admissions for rat bites to hospitals in Bombay each year.³ Most bites occur during sleep, and victims usually wake and frighten the rat away, except in the case of small infants or the very infirm. People with peripheral neuropathy, however, do not feel the bites, and the rat can continue to feed undisturbed.

Five per cent of adult Dominicans are diabetic (R Fortune, unpublished data), and the ratio of rats to humans in Dominica may be as high as five to one (B Xavier, personal communication). Housing conditions are poor, largely owing to hurricane David, which in 1979 left three quarters of the people homeless. Diabetic foot is a well recognised problem.⁴ The combination of a high population of rats, poor housing, and a high prevalence of diabetes means that rat bites commonly cause lesions on diabetic feet. The incidence of such lesions is underestimated as patients may not volunteer or even realise that they have been caused by rats. We advise all of our diabetic patients to cover their feet while asleep and seek immediate medical help for any foot lesion.

- 1 Walrond EW, Jordan OW. Suggestions for a new approach to diabetes. *Cajanus* 1985;18:127-30 (Caribbean Food and Nutrition Institute, Jamaica).
- 2 Clinton JM. Rats in urban America. *Public Health Rep* 1969;84:1-7.
- 3 Deoras PJ. Some observations on the probable damage caused by rats in Bombay. *Indian Journal of Entomology* 1966;28:543-7.
- 4 Alberti KGGM, Hockaday TDR. Diabetic foot disease. In: Weatherall DJ, Ledingham JGG, Warrell DA, eds. *Oxford textbook of medicine*. Oxford: Oxford University Press, 1984:9.37-8.

(Accepted 15 September 1988)