- 23 Baran DT, Teitelbaum SL, Berfield MA, *et al*. Effects of alcohol ingestion on bone and mineral metabolism in rats. *Am J Physiol* 1980;238:E507-10.
- 24 Peng TC, Garner SC, Frye GD, *et al*. Evidence of a toxic effect of ethanol on bone in rats. *Alcoholism* 1982;6:96-9.
- 25 Daniell HW. Osteoporosis of the slender smoker: vertebral compression fractures and loss of metacarpal cortex in relation to postmenopausal smoking and lack of obesity. *Arch Intern Med* 1976;136:298-304.
- 26 Kanders B, Dempster DW, Lindsay R. Interaction of calcium nutrition and physical activity on bone mass in young women. *Journal of Bone and Mineral Research* 1988;3:145-9.
- 27 Riggs BL, Wahner HW, Melton LJ, Richelson LS, Judd HL, O'Fallon WM.

Dietary calcium intake and rate of bone loss in women. *J Clin Invest* 1987;80:979-82.

- 28 Stevenson JC, Whitehead MI, Padwick M, *et al*. Dietary intake of calcium and postmenopausal bone loss. *Br Med J* 1988;297:15-7.
- 29 Goldsmith NF, Johnston JO. Bone mineral: effects of oral contraceptives, pregnancy, and lactation. *J Bone Joint Surg* 1975;57A:657-68.
- 30 Christiansen C, Riis BJ, Rodbro P. Prediction of rapid bone loss in postmenopausal women. *Lancet* 1987;i:1105-8.

(Accepted 6 February 1989)

Pulmonary oedema after acute asphyxia in a child

M Boykett

I report a case of severe pulmonary oedema complicating acute upper airway obstruction.

Case report

A previously well 8 year old boy was found having choked on a vitamin tablet (Sanatogen Junior, Fisons). He was apnoeic, unresponsive, and cyanosed. During attempts at basic airway resuscitation he coughed up bloodstained mucus before recovering consciousness.

During the next one to two hours, after admission to hospital, he developed tachypnoea, cyanosis, and bilateral lung crepitations. He had radiological signs of florid bilateral pulmonary oedema without cardiomegaly. Oxygen, ampicillin, and hydrocortisone were given, and a rigid bronchoscopy was performed to ensure that there was no residual foreign body. Anaesthesia and bronchoscopy were rendered difficult by a profuse amount of bloodstained froth pouring from the larynx. No foreign body or vomit was found, and he was given intensive care.

With intermittent positive pressure ventilation with 5 cm H₂O positive end expiratory pressure and an inspired fractional oxygen concentration of 0.6 his oxygen saturation was initially 87.5% (arterial oxygen tension 9.1 kPa). The arterial pH of 7.18 reflected mixed metabolic and respiratory acidosis. His general condition, breathing, and arterial blood gas measurements all improved rapidly over the next six to 12 hours. Subsequently, he maintained a normal arterial oxygen tension during spontaneous breathing at 5-10 cm continuous positive airway pressure and an inspired oxygen fractional concentration of 0.25-0.30. Extubation was delayed for two days because of a persistent pink tracheal aspirate and until the radio-

logical abnormalities had resolved. Further recovery was complete and unremarkable.

Comment

An accurate diagnosis proved elusive. Fluid overload and primary left ventricular failure were dismissed as possible causes, and pulmonary aspiration and chemical pneumonitis were considered to be the main differential diagnoses. The history and clinical findings, however, were not characteristic of aspiration. The manufacturer of the tablet and Guy's Hospital's Poisons Unit were consulted, but chemical irritation was excluded as a likely cause.

Similar cases of postobstructive pulmonary oedema have occurred in, for example, laryngospasm,¹ croup, and epiglottitis.² Most cases were in children but were less severe than the one reported here, and none was associated with an impacted foreign body. Brown reviewed previous reports and proposed mechanisms by which airway obstruction may cause pulmonary oedema.³ These include negative intrathoracic pressure causing transiently low pulmonary interstitial pressure or impaired left ventricular function, or both, and hypoxic postcapillary and venous constriction. Treatment with oxygen alone may be sufficient, but a brief period of tracheal intubation and assisted ventilation may be necessary. Upper airway obstruction occurs frequently, and doctors should be aware of this unfamiliar complication.

I thank Dr M Bexton and Dr A Syed for their help in managing this patient; Dr A Vaizey for permission to report on the patient; Mrs M Chambers and Miss J Norton for typing.

- 1 Lee KWT, Downes JJ. Pulmonary edema secondary to laryngospasm in children. *Anesthesiology* 1983;59:347-9.
- 2 Galvis AG, Stool SE, Bluestone CD. Pulmonary edema following relief of upper airway obstruction. *Ann Otol Rhinol Laryngol* 1980;89:124-8.
- 3 Brown RE. Negative pressure pulmonary edema. In: Berry FA, ed. *Anesthetic management of difficult and routine pediatric patients*. New York: Churchill Livingstone, 1986:168-79.

(Accepted 9 December 1988)

Hallucinations after epidural buprenorphine

M MacEvilly, C O'Carroll

Buprenorphine, a mixed central nervous system opiate agonist-antagonist, is virtually free of hallucinatory effects in parenteral formulation, the incidence being 0.1%.¹ Epidural administration of buprenorphine, however, may not be quite as free of these side effects.

Case reports

Case 1—A 40 year old woman underwent fusion of the posterior lumbar bodies at L4-5 under general anaesthetic. Postoperatively she was given 150 µg

buprenorphine every eight to 12 hours as needed through an epidural cannula. Forty eight hours after surgery and after a total dose of buprenorphine of 900 µg she complained of nightmares (table). The symptoms occurred only at night and within 10 minutes after the drug was topped up. She was also distressed by the shadows in the room and became anxious when left alone. The symptoms disappeared once buprenorphine was stopped.

Case 2—A 38 year old woman underwent posterior spinal fusion between the transverse processes (L1-3). Postoperatively analgesia was maintained through an epidural cannula with 150 µg buprenorphine every eight to 12 hours as necessary. After 48 hours she complained of a recurring dream in which she recalled a frightening incident from her past. The dream occurred so vividly at night that she became extremely distressed. She had received a total dose of 750 µg

Pilgrim Hospital, Boston,
Lincolnshire PE21 9QS
M Boykett, FFARCS,
anaesthetic registrar

Correspondence to:
Leicester Royal Infirmary,
Leicester LE1 5WW.

Br Med J 1989;298:928

Blackrock Clinic,
Blackrock, County Dublin,
Republic of Ireland
M MacEvilly, FFARCSI,
consultant anaesthetist
C O'Carroll, SRN, sister,
Nightingale unit

Correspondence to: Dr
MacEvilly.

Br Med J 1989;298:928-9