dispense with inpatient beds altogether. Even when intensive care is given by enthusiastic and committed staff with residential care facilities outside hospital psychiatric admissions will always be needed.¹⁴ Probably a greater proportion of those who need admission will be more disturbed as others can be dealt with outside hospital. This has been our experience and has economic implications.¹⁵

Although generalising from services in one part of the country to another is difficult, Nottingham is representative of areas with middling demand. Its use of psychiatric beds for the 15-65 age group in 1980 was close to the mean of 0.45 bed per 1000 population estimated by Hirsch to reflect the requirements of England and Wales as a whole.¹⁶ If the fall to the 1985 rate in Nottingham was reflected nationally the requirements for beds would be reduced by over a third. Our findings of reduced use of beds should not be interpreted as showing the newer service to be superior; that is the subject of a separate investigation. Nevertheless, we conclude with some confidence that when community and hospital psychiatric services are combined in an integrated model the benefits are quickly shown and economically feasible.

We thank Professor John Cooper and other members of the Nottingham case register, particularly Dr Sarah White, Pat Mounser, Norah Davis, and Richard Gancarczyk, for their help in preparing and checking data from the case register for this survey in statistical bulletins. We also thank Karen

Robinson of the Department of Health and Social Security for help in obtaining national figures and for estimating the average number of beds used daily in NHS mental illness hospitals.

- 1 Jones K, Poletti A. The Italian experience in mental health care. *Hosp Community Psychiatry* 1986;37:795-801.
- 2 Hafner H. Do we still need beds for psychiatric patients? An analysis of changing patterns of mental health care. *Acta Psychiatr Scand* 1987;75:113-26.
- 3 MacMillan D. Community mental health services and the mental hospital. In: Freeman H, Farndale J, eds. *Trends in the mental health services*. London: Pergamon Press, 1963:303-17.
- 4 Tyrer P. Psychiatric clinics in general practice: an extension of community care. *Br J Psychiatry* 1984;145:9-14.
- 5 Tyrer P, Seivewright N, Wollerton S. General practice psychiatric clinics: impact on psychiatric services. *Br J Psychiatry* 1984;145:15-9.
- 6 Jones SJ, Turner RJ, Grant JE. Assessing patients in their homes. *Bulletin of the Royal College of Psychiatrists* 1987;11:117-9.
- 7 Department of Health and Social Security. *Better services for the mentally ill*. London: HMSO, 1975.
- 8 Tyrer P. The hive system: a model for a psychiatric service. *Br J Psychiatry* 1985;146:571-5.
- 9 Griffiths R. *Community care: agenda for action*. London: HMSO, 1988.
- 10 Murphy E. Griffiths on community care: potentially disastrous. *Br Med J* 1988;296:876-7.
- 11 Donovan CN. Problems of psychiatric practice and community mental health centers. *Am J Psychiatry* 1982;139:456-69.
- 12 Williams P, Balestrieri M. Psychiatric clinics in general practice: do they reduce admissions? *Br J Psychiatry* 1989;154:67-71.
- 13 Burti L, Garzotto N, Siciliano O, Zimmerman-Tansella C, Tansella M. South Verona's psychiatric service—an integrated system of community care. *Hosp Community Psychiatry* 1986;37:809-13.
- 14 Hoult J. Community care of the acutely mentally ill. *Br J Psychiatry* 1986;149:137-44.
- 15 Tyrer P, Gill D, Farr P. Cost of community psychiatry. *Br J Psychiatry* 1986;149:795-6.
- 16 Hirsch SR. Planning for bed needs and resource requirements in acute psychiatry. *Bulletin of the Royal College of Psychiatrists* 1987;11:398-407.

(Accepted 30 May 1989)

Effect of hysterectomy on bowel function

T Taylor, A N Smith, P M Fulton

Hysterectomy is a common, comparatively safe operation. Many complications are known, but its possible relation with bowel function is less well reported. We performed a case-control study to see whether hysterectomy had any long term effect on bowel function and whether this was associated with altered urinary function.

Patients, methods, and results

We studied women in two general practices in north Edinburgh who had had a hysterectomy for benign disease two to eight years previously. Each woman was randomly matched by age and sex with another woman from the same practice. By reading practice records we excluded from the control group women who had had extensive abdominal operations and those with the irritable bowel syndrome. We devised a simple questionnaire to assess bowel function. It was adapted from those used in other studies^{1,2} and validated in a sample of the women. One hundred pairs of women were sent questionnaires; 96 women who had had a hysterectomy and 95 controls responded, giving 91 matched pairs. Their age range was 32 to 63 (mean 48.8) years. We compared each pair by looking at

bowel function (bowel frequency, use of laxatives, and consistency of stools).

The table gives the results. Women who had had a hysterectomy more commonly reported infrequent bowel motions. Fourteen of these women had a bowel motion every four days or less, compared with six controls (Wilcoxon signed rank test on paired data, $p < 0.05$). The women who had had a hysterectomy were more likely to use laxatives and have hard stools, although the differences between the groups were not significant. Twenty one women who had had a hysterectomy but only 10 controls said that they were constipated when the other member of the pair did not. This difference, although not significant, was unlikely to have occurred by chance ($p = 0.07$, McNemar's test). Fifteen women who had had a hysterectomy but only four controls had consulted a doctor about constipation ($p < 0.05$, McNemar's test). Ten of the women who had had hysterectomy had persistently decreased bowel frequency and persistently increased urinary frequency. Of these, eight had noticed both these changes after hysterectomy. This was a highly significant association ($p < 0.01$, χ^2 test).

Comment

Our results show that after a hysterectomy women tend to have less frequent bowel actions than controls. Women who had had a hysterectomy consulted a doctor because of constipation more commonly than controls, which may indicate that their constipation was more severe and required professional help. There are several possible mechanisms for this difference: psychological, hormonal, pharmacological, and surgical. The mechanism is unlikely to be psychological as women are unlikely to become depressed after hysterectomy.³ Women with constipation due to slow transit tend to have increased serum prolactin concentrations and decreased plasma oestradiol concentrations.⁴ Altered hormone concentration may affect bowel habit after hysterectomy, especially in those women who also have oophorectomies. Some women

Western General Hospital,
Edinburgh EH4 2XU

T Taylor, MRCP, senior house
officer, university
gastrointestinal unit

A N Smith, FRCSed,
professor, university
department of
surgery

Department of Community
Medicine, University of
Edinburgh Medical School,
Edinburgh EH8 9AG

P M Fulton, FRCPed, senior
lecturer

Correspondence and
requests for reprints to:
Professor Smith.

Br Med J 1989;299:300-1

Bowel function in women who had had hysterectomy and controls

	Bowel frequency		Use of laxatives		Stool consistency	
	At least every 3rd day	Every 4th day or less often	Fortnightly or more often	Less than monthly or never	Hard	Not hard
Women who had had hysterectomy	75	14	16	71	15	75
Controls	84	6	7	80	10	79

may take large amounts of laxatives for postoperative constipation, leading to the "cathartic colon." If this had been so in our study we would have expected more use of laxatives by the women who had had hysterectomy than we found, although such use might have been underreported deliberately. Another possible mechanism is pelvic autonomic denervation due to an intraoperative manoeuvre,⁵ traction during the operation, or realignment of the pelvic organs after removal of the uterus.

Urinary dysfunction is a well recognised complication of hysterectomy and is thought to be due to disruption of the nerve supply to the bladder. We found a highly significant association between persistently reduced bowel frequency and persistently increased urinary frequency after hysterectomy. This supports the hypothesis that they may have a common aetiology—namely, autonomic denervation of both

viscera, which have a closely related nerve supply in the pelvis.

We thank Dr A A Robertson, Dr A Donald and partners, Dr M A Eastwood, and Mr J Knowles (medical records officer) for their help with this study. We also thank Dr J W Yarnell for giving us a copy of his questionnaire.

- 1 Connell AM, Hilton C, Irvine G, Lennard-Jones JE, Misiewicz JJ. Variation of bowel habit in two population samples. *Br Med J* 1969;iii:1095-9.
- 2 Yarnell JW, Voyle GJ, Richards CJ, Stephenson TP. The prevalence and severity of urinary incontinence of women. *J Epidemiol Community Health* 1981;35:71-4.
- 3 Coppen A, Bishop M, Beard RJ, Barnard GJR, Collins WP. Hysterectomy, hormones and behaviour: a prospective study. *Lancet* 1981;i:126-8.
- 4 Preston DM, Rees LH, Lennard-Jones JE. Gynaecological disorders and hyperprolactinaemia in chronic constipation. *Gut* 1983;24:A480.
- 5 Catchpole BN. Motor pattern of left colon before and after surgery for rectal cancer: possible implications in other disorders. *Gut* 1988;29:624-30.

(Accepted 29 April 1989)

Radioiodine in human thyroid glands and incidence of thyroid cancer in Cumbria

Colin Bowl, Peter Tiplady

Department of Radiation Biology, St Bartholomew's Hospital Medical College, London EC1M 6BQ
Colin Bowl, PhD, senior lecturer

Department of Community Medicine, East Cumbria Health Authority, Cumberland Infirmary, Carlisle CA2 7HY
Peter Tiplady, FFCM, district medical officer

Correspondence to: Dr Bowl.

Br Med J 1989;299:301-2

Concern is increasing over the association between ionising radiation and cancers in populations living close to nuclear installations such as Sellafield in Cumbria. Because the external radiation in Cumbria is only marginally above the national average¹ any appreciable additional dose would have to come from manmade sources of radioactivity.

Both iodine-129 and iodine-131 are discharged from Sellafield into the sea and the atmosphere and since the early 1960s have probably contributed equally to the small additional dose^{2,3} to the thyroid above that of natural background radiation. We measured ¹²⁹I activity in thyroid glands in the population of Cumbria and related them to the incidence of thyroid cancer in the area. Inevitable delays in carrying out these measurements excluded an assessment of ¹³¹I, which has a half life of eight days (that of ¹²⁹I is 15 million years).

November 1984 and September 1987. The thyroid samples were homogenised and dried, and ¹²⁹I activity was measured in a thin walled scintillation detector over 100 minutes. Background levels (which varied during the examination) were subtracted from the count rate for the sample, and occasional negative values resulted. The figure (left) shows ¹²⁹I activity in individual thyroid glands plotted against the distance of the patients' homes from Sellafield. Levels of radioactivity decreased with distance (Spearman's rho = -0.392, p < 0.001).

We also examined the incidence of thyroid cancer, using data from the Northern regional cancer registry for 1969-86. Age standardised registration rates and ratios were calculated by applying the national values for the midpoint of the period. Surprisingly, there was a positive correlation between the distance from Sellafield and the incidence of cancer, with higher incidences being found with greater distance (Spearman's rho = 0.541, p = 0.0007) (figure (right)). In the 40 (out of 288) parishes where thyroid cancers occurred the incidence varied from 0.39 to 40.46/100 000. Calculations using the Poisson distribution,⁴ however, showed only two parishes with significant increases in thyroid cancer: one was 16 km from Sellafield (two cases registered v 0.1 expected, p < 0.01) and the other 24 km away (four cases registered v 0.5 expected, p < 0.01). In Cumbria

Methods and results

Measurements were made on 130 thyroid glands taken opportunistically at necropsy from adults (mean age 67) dying from various causes in Cumbria between

Left: ¹²⁹I activity in thyroid tissue taken at necropsy related to distance of patients' homes from Sellafield. Right: registrations of thyroid cancer in Cumbria, 1969-86

SRR = Standardised registration ratio