Novel role for specialist nurses in managing diabetes in the community

Mary MacKinnon, R Malcolm Wilson, C A Hardisty, J D Ward

Abstract

To develop care of diabetes further a specialist nurse established contact with general practices in Sheffield Health District and identified difficulties in providing a service for diabetics. One hundred and thirty practices were visited, and full data were collected from 104. Each practice agreed to establish a register of diabetics, and information and support were subsequently provided to help in developing services. In collecting information from each practice the nurse covered specific points on staff, facilities, and organisation.

Over two years the service offered in 60 practices considerably improved, allowing a minimum standard of diabetic care to be achieved. This allowed coordinated and effective referral of certain patients from hospital diabetic clinics and improved services to those not attending any clinics.

Introduction

The tradition of the large diabetic clinic serving a district creates deficiencies, not the least being that a large proportion of diabetic patients do not attend. Over the past 15 years several methods of sharing care or collaboration between hospital and general practice have been reported. Thorn established the principle of "miniclinics" in Wolverhampton, initially covering 14 practices.¹ The community care service for diabetics in Poole took two years to become operational and required intensive preparation and dedication from the interested physician and local general practitioners. To facilitate the scheme computer programs were developed to store clinical data and organise clinic visits and screening. Continued efforts have established this as an effective scheme.²

In Sheffield over 1000 patients were discharged from a diabetic clinic over three years. A nurse was available to visit practices, but the scheme was not coordinated effectively, and no attention was paid to the problems many practices faced. As a result many patients were either not being seen regularly by anybody or under the impression that their diabetes was cured.³ In Cardiff, a randomised trial compared hospital care and general practitioners' services, but the doctors were given no special support, and this attempt again highlighted the need for planning, collaboration, preparation, and the provision of suitable support.⁴ Considerable personal collaborative contact is essential in any scheme aimed at improving services for diabetics in a community. Sending a questionnaire to a large number of general practitioners does not develop useful links.⁵ In Ipswich a training programme was available, cooperation booklets were produced, and considerable back up was given regarding retinopathy screening. An efficient register was kept,⁶ but again a further report showed that a high

proportion of patients in this scheme did not receive adequate care.⁷

In many of these studies criticism of the general practitioner is implied by the fact that so many patients were not being seen or were not receiving appropriate care and screening for the complications of diabetes. Such criticism, however, denies the fact that many difficulties arise in developing a service for diabetics in which regular supervision and education of patients are at the centre of management. Moreover, many practices are being encouraged to provide other medical services. We set up this study to identify the practical difficulties that hinder the development of a minimum standard of care of diabetics in general practice.

Subjects and methods

Sheffield has a stable population of about 550 000. Four specialists in diabetes provide care for adults in two main hospitals in the district, there being separate services for children in two paediatric departments.

We compiled a list of practices from the family practitioner list and informed the family practitioner committee of the project. All practices were located, mapped, and coded. The city was divided into sectors for easy recognition and to enable practices to be visited in groups. A study group of practice nurses had been formed by the specialist nurse two years previously, and practices employing nurses were identified. Discussions were held with the specialists in diabetes in the city, and a letter from the specialist nurse and doctors was sent to all general practitioners introducing the concept of the survey and offering help to develop services for diabetics. At the same time other important groups were informed of the project: the local medical committee, senior nursing administration, chiropodists, dietitians, and the local branch of the British Diabetic Association.

The nurse visited each practice for discussions with one or more general practitioners. Other members of the staff were included whenever possible. Discussion in each practice followed a defined course. It included discussion of numbers of patients in the practice, geography, staff, facilities, organisation, and other services provided. Existing services for diabetics and problems encountered in their provision were examined. Where no service existed a whole package was given. This included updates on diabetes, ideas towards a protocol, answers to questions, and the offer of help.

Most practices requested follow up visits, this in itself indicating the value that they attached to the discussions and to the help that was being offered. Particular help requested was, for example, with an initial audit and how best to identify the diabetic population. Follow up visits also included the nurse

**Royal Hallamshire
Hospital, Sheffield S10 2JF**
Mary MacKinnon, RGN,
diabetes nurse specialist
R Malcolm Wilson, DM,
senior lecturer in medicine
J D Ward, MD, consultant
physician

**Northern General
Hospital, Sheffield S5 7AU**
C A Hardisty, MD, consultant
physician

Correspondence to:
Mrs MacKinnon.

Br Med J 1989;299:552-4

being present and giving support at sessions until the practice staff gained confidence.

Results

One hundred and thirty practices were visited, and complete information is presented from 104 practices. Of these, 42 practices consisted of groups of three or more doctors; 31 consisted of two doctors; and 31 were singlehanded practices, in which potentially there were additional problems in developing a service. Eighteen practices had branch surgeries, and in many instances patients living near the branch surgery were reluctant to attend sessions for diabetics at the main surgery. In smaller practices poor buildings and lack of staff and facilities did not allow practitioners to improve their services.

All practices used laboratory facilities for estimates of blood glucose concentrations when required. Fourteen practices that already provided a service for diabetics used visually read blood glucose strips or meters, but there was clearly a need for updating and further education on the use and maintenance of such equipment.

All practices had most of the equipment required to run a service for diabetics. The main difficulties were inaccurate weighing scales, absence of a height gauge, expired equipment for testing blood and urine, and no chart to assess visual acuity. Servicing of sphygmomanometers and ophthalmoscopes was also required in some practices. Advice was given on the type of equipment and where to obtain supplies or servicing at a low cost. Computers or video facilities, or both, were available in six practices. These items of equipment have recently become more widely available.

At the beginning of the survey only 14 practices had attempted to review the feet of diabetics and identify those at risk. Two years later, however, 60 practices aimed to provide such a service. All practices indicated great concern about the provision of chiropody services for their patients who were not attending hospital clinics.

As no community dietitian was available practice nurses were taught simple dietary guidelines for patients with non-insulin dependent diabetes. They were encouraged to "personalise" their advice after receiving daily meal diaries. Simple information was provided to help them. Practice staff also required help and information on food, fluid, and general advice during illness for both non-insulin dependent and insulin dependent diabetics. Referral information about the district dietetic services was given with the name of the chief dietitian in each hospital. Fourteen practices could supply dietary advice at the beginning of the survey; after help from the nurse specialist 60 were able to supply this advice.

There were major concerns about screening of eyes. Fifteen practices carried out routine funduscopy through dilated pupils at the start of the study. The major worries were the small numbers of patients and consequent lack of skill; insufficient time especially in larger practices; the safety of mydriatics and the precipitation of acute glaucoma; and organisation of screening. Most general practitioners in discussion expressed the need for a training scheme; a few thought that a clinical assistantship would be useful. Eighty five practices either periodically suggested examination by the optician or gave no advice. There was no system of checking that eye screening or advice had been given in the 85 practices. A system was subsequently introduced, however, in the 51 practices providing a service. The whole question of visual problems and screening for diabetic retinopathy was discussed in all practices by the specialist nurse, who supplied the

following: correctly set up Snellen charts; information about ophthalmoscopes; organisation and recall systems; and advice about mydriatics.

Only seven practices were concerned with the financial implications of providing or improving services for diabetics, so this did not seem to be a major consideration.

Care, surveillance, and education provided for diabetics in 104 practices at beginning and end of study

	1985	1987
Practice nurse employed	20	59
Register, recall, follow up	16	82
Service for diabetics	16	60
Education for diabetics	13	57
Screening for retinopathy	15	51

After two years 60 practices provided services at differing standards for diabetic patients, and the table summarises these improvements. There were obvious improvements in the practices' awareness and care of diabetic patients where good communication and working relationships were well developed.

Discussion

Each general practice is an individual entity providing flexible treatment and preventive services to its population. Any new system must be organised according to the availability of facilities and the needs of patients; this is particularly so for diabetics. Unlike in the hospital service administrative, organisational, and laboratory support is not near at hand. In attempting to improve services to diabetic patients whether in hospital or in general practice considerable liaison between doctors and nurses is needed. This can be provided by a nurse specialising in care for diabetics. During our study, with the encouragement of the local family practitioner committee, practice premises improved, staff numbers increased, and motivation increased to provide adequate clinical care of and services for diabetics.

The fundamental need in a service for diabetics in the community is the provision of a register of diabetics. Our major achievement has been to establish such a register, which, coupled with the known patients attending hospital clinics, has begun to approach total ascertainment of the diabetic population. In any large city overlapping boundaries (geographical, district, city, family practitioner committee, and practice) may cause problems of follow up and continuity of care. In Sheffield 4000 patients' details are now available on practice registers. Simply constructed, these registers have proved invaluable in finding out where patients are seen, if at all. Education on diabetes and clinical review can be missed when patients have multiple medical problems or have been referred to a doctor with no special interest in diabetes. A named receptionist in each practice (including the branch surgery) in collaboration with the practice nurse can keep the register of diabetics and recall system up to date.

Copies of the registers from individual practices collected by the specialist nurse will be entered into a district computer system. In this way the patients of general practitioners who are not providing a service can be identified and offered clinical and screening services through the hospital services. Eventually, establishing a full time centre purely for diabetics will allow the coordination of such aspects of care.

Obviously the role of the coordinating nurse is important, but many practice nurses are also intimately concerned in organising and running clinical services

for diabetics in general practice.⁸ As with the general practitioners themselves the danger of isolation from discussion and information about clinical management is ever present, and general practitioners and practice nurses should have regular update meetings. As we have shown, the range of help required by general practitioners and nurses is considerable. Most seemed unaware of facilities and services provided by the British Diabetic Association. The folder on diabetes produced by the Royal College of General Practitioners was not commonly used, most practices preferring the opportunity to discuss problems with the specialist nurse. Certainly the study highlighted the need for continuing and adequate communications between hospital and general practice with a sharing of clinical information, particularly when patients are attending hospital whether in specialist diabetes clinics or not. Information and help should be provided for reception and office staff in practices. Their role in contacting patients, organising services, and developing care for diabetics in general practice is important.

People with diabetes require information, support, and surveillance. Provision of services to the diabetic population in a large city can be considerably improved

with a positive approach and the help of a specialist nurse.

We thank Servier Laboratories UK for funding this project; the general practitioners, practice nurses, and staff in Sheffield, who continue to participate in expanding their services; Dr M B Collins and partners and Drs P G Newrick and A M Jennings for their support and advice; and Heather Gibson for clerical help.

- 1 Thorn PA, Russell RG. Diabetic clinics today and tomorrow: mini-clinics in general practice. *Br Med J* 1973;ii:534-6.
- 2 Hill RD. Community care for diabetics in the Poole area. *Br Med J* 1976;ii: 1137-9.
- 3 Wilks E, Lawton EE. The diabetic, the hospital and primary care. *J R Coll Gen Pract* 1980;30:199-206.
- 4 Hayes TM, Harris J. Randomised controlled trial of routine hospital clinic care versus routine general practice care for type II diabetics. *Br Med J* 1984;289:728-30.
- 5 Mellor JG, Samanta A, Blanford RL, Burdon AC. Questionnaire survey of diabetic care in general practice in Leicestershire. *Health Trends* 1985;17: 61-3.
- 6 Day JL. Part 1. The organisation of diabetic care. *Hospital Update* 1980;6: 1177-83.
- 7 Day JL, Humphreys H, Alban-Davis H. Problems of comprehensive shared diabetes care. *Br Med J* 1987;294:1590-2.
- 8 MacKinnon M. The role of nurses in general practice. *Practical Diabetes* 1986;3:232-4.

(Accepted 19 June 1989)

BOOKS RECEIVED

Acquired immune deficiency syndrome

AIDS: Principles, Practices, and Politics. Ed I B Corless, M Pittman-Lindeman. (Pp xvii+252; £58.) Washington: Hemisphere Publishing, 1988. ISBN 0-89116-795-1.

AIDS: the HIV Myth. J Adams. (Pp xii+244; £12.95.) London: Macmillan, 1989. ISBN 0-333-48930-6.

HIV Detection by Genetic Engineering Methods. Ed P A Luciw, K S Steimer. (Pp xiii+301; figs; \$119.50.) New York: Dekker, 1989. ISBN 0-8247-7900-2.

Addiction

EURO Reports and Studies 108. "Alcohol and the Mass Media." J Partanen, M Montonen. (Pp 73; Sw frs 8 paperback.) Copenhagen: World Health Organisation, 1988. ISBN 92-890-1274-9.

Psst... a Really Useful Guide to Alcohol. A Maryon Davis. (Pp viii+152; figs; £3.99 paperback.) London: Pan Books, 1989. ISBN 0-330-31109-3.

Allergy

Allergy: the Facts. R Davies, S Ollier. (Pp viii+200; figs; £4.95 paperback.) Oxford: Oxford University Press, 1989. ISBN 0-19-261858-X.

Alternative medicine

Medicine in Rural China: a Personal Account. C C Chen, in collaboration with F M Bunge. (Pp xix+218; figs; \$35.) Berkeley: University of California Press, 1989. ISBN 0-520-06298-1.

Standard Acupuncture Nomenclature. Part 2. World Health Organisation. (Pp 17; paperback, price not stated.) Manila: World Health Organisation Regional Office for the Western Pacific, 1988.

Anaesthesia

Anaesthesia and Malignant Disease. J Filshie, D S Robbie. (Pp xv+303; figs; £27.50.) London: Arnold, 1989. ISBN 0-7131-4553-6.

Anatomy

Anatomy and Human Movement: Structure and Function. N Palastanga, D Field, R Soames. (Pp iv+891; figs;

£60.) Oxford: Heinemann Medical, 1989. ISBN 0-433-00032-5.

Blood pressure

The BMA Family Doctor Guides. "High Blood Pressure." P Semple. Series editor T Smith. (Pp 95; figs; £2.99 paperback.) London: Equation in association with the British Medical Association, 1989. ISBN 1-85336-053-8.

Calcium in Essential Hypertension. Ed K Aoki, E D Frohlich. (Pp xviii+675; \$150.) Tokyo: Academic Press, 1989. ISBN 0-12-058845-5.

Cardiovascular diseases

Cardiovascular Clinics. "Heart Disease in Women." Ed P S Douglas. Series editor A N Brest. (Pp xx+316; figs; colour plate; £50.) Philadelphia: Davis, 1989. ISBN 0-8036-2751-3.

Developments in Cardiovascular Medicine. Vol 92. "Electrocardiography and Cardiac Drug Therapy." Ed V Hombach, H H Hilger, H L Kennedy. (Pp xiii+369; figs; £68.) Dordrecht: Kluwer, 1989. Distributed by MTP Press. ISBN 0-89838-395-1.

Cardiovascular system

The Circulation in the Female from the Cradle to the Grave. Ed J Ginsburg. (Pp xv+174; figs; £38.) Carnforth: Parthenon, 1989. ISBN 0-940813-68-8.

Computers in medicine

Practical Guides for General Practice. 7. "Computers: a Guide to Choosing and Using." A Willis, T Stewart. (Pp iii+138; figs; £6.95 paperback.) Oxford: Oxford University Press, 1989. ISBN 0-19-261754-0.

Disabled/handicapped people

Dictionary of Mental Handicap. M P Lindsey. (Pp vi+345; £35.) London: Routledge, 1989. ISBN 0-415-02810-8.

Endocrinology

Calcium Homeostasis: Hypercalcaemia and Hypocalcaemia G R Mundy. (Pp vii+248; figs; £49.95.) London: Dunitz, 1989. ISBN 0-948269-67-7.

Pineal Research Reviews. Vol 7. Ed R J Reiter. (Pp xiii+234; £120.) New

York: Liss, 1989. Distributed by John Wiley and Sons. ISBN 0-8451-3606-2.

Environmental and public health

Architectural Design and Indoor Microbial Pollution. Ed R B Kundsinn. (Pp x+317; figs; £37.50.) New York: Oxford University Press, 1989. ISBN 0-19-504436-3.

Engineered Organisms in the Environment. [A 'lay summary' based on the First International Conference on the Release of Genetically Engineered Microorganisms (REGEM 1), Cardiff, 5-8 April 1988.] B Dixon. (Pp 47; figs; £5 paperback, including UK postage, overseas add 50 pence.) Obtainable from Professor A M Paton, University of Aberdeen, School of Agriculture, 581 King Street, Aberdeen AB9 1UD.

EURO Reports and Studies. No 78. "Indoor Air Pollutants: Exposure and Health Effects." Report on a World Health Organisation Meeting. (Pp 42; Sw frs 4 paperback.) Copenhagen: World Health Organisation Regional Office for Europe, 1989. ISBN 92-890-1244-7.

Gastroenterology

Frontiers of Gastrointestinal Research. Vol 15. "Newer Tests and Procedures in Pediatric Gastroenterology. 1. Diagnostic and Therapeutic Procedures." Ed G Dinari, P Rozen, Y Bujanover, E Leblenthal. Series editor P Rozen. (Pp viii+330; figs; £135.) Basel: Karger, 1989. Distributed by John Wiley and Sons. ISBN 3-8055-4646-7.

Frontiers in Gastrointestinal Research. Vol 16. "Newer Tests and Procedures in Pediatric Gastroenterology. 2. Function and Laboratory Tests." Ed G Dinari, P Rozen, Y Bujanover, E Leblenthal. Series editor P Rozen. (Pp viii+296; figs; £121.90.) Basel: Karger, 1989. Distributed by John Wiley and Sons. ISBN 3-8055-4900-8.

Genetics

The Foundations of Human Genetics. K R Dronamraju. (Pp xiii+211; figs; \$40.75.) Illinois: Thomas, 1989. ISBN 0-398-05537-8.

Haematology

Hematology: a Combined Theoretical and Technical Approach. A Simmons. (Pp viii+423; colour plates; figs; £25

paperback.) Philadelphia: Saunders, 1989. Distributed by Harcourt Brace Jovanovich. ISBN 0-7216-1716-6.

Progress in Hemostasis and Thrombosis. Vol 9. Ed B S Coller. (Pp xiv+269; figs; £41.) Philadelphia: Saunders, 1989. Distributed by Harcourt Brace Jovanovich. ISBN 0-7216-2933-4.

Quality Assurance in Haematology. Ed S M Lewis, R L Verwilghen. (Pp xii+297; figs; £25.) London: Baillière Tindall, 1988. Distributed by Harcourt Brace Jovanovich. ISBN 0-7020-1322-6.

Health care issues

EDI Seminar Paper. No 37. "User Charges for Health Care in Principle and Practice." C C Griffin. (Pp vii+80; \$6.50 paperback.) Washington DC: World Bank, 1988. Distributed in the United Kingdom by Microinfo Ltd., PO Box 3, Alton, Hants GU34 2PG. ISBN 0-8213-1139-5.

Facilitating Change in Mental Health. B Lemmer, M Smits. (Pp xviii+222; figs; £10.95 paperback.) London: Chapman and Hall, 1989. ISBN 0-412-33010-5.

From Alma-Ata to the Year 2000: Reflections at the Midpoint. World Health Organisation. (Pp vi+158; Sw frs 30 paperback.) Geneva: World Health Organisation, 1988. ISBN 92-4-156124-6.

Mental Health Care in Crisis. Ed A Brackx, C Grimshaw. (Pp viii+216; £14.95.) London: Pluto Press, 1989. ISBN 0-7453-0221-1.

Talking and Listening to Patients: a Modern Approach. C M Fletcher, P Freeling. (Pp viii+68; £2.50 paperback.) London: Nuffield Provincial Hospitals Trust, 1989. ISBN 0-900574-70-4.

Therapy in Practice. "Counselling Skills for Health Professionals." P Burnard. Series editor J Campling. (Pp xii+201; figs; £10.95 paperback.) London: Chapman and Hall, 1989. ISBN 0-412-32000-2.

History of Medicine

The British Paediatric Association 1928-1988. J O Forfar, A D M Jackson, B M Laurance. (Pp xix+228; figs; £5 including postage.) 1989. Available from the British Paediatric Association, 5 St Andrew's Place, London NW1 4LB.

Resources in Medical History. "An

Essay on Fevers." J Huxham. (Pp xxxi+191; \$19.95.) Canton: Science History Publications, 1989. ISBN 0-88135-083-4.

The Wellcome Institute Series in the History of Medicine. "The Hospital in History." Ed L Granshaw, R Porter. Series editors W F Bynum, R Porter. (Pp ix+273; £35.) London: Routledge, 1989. ISBN 0-415-00375-X.

Human reproduction

Contemporary Issues in Biomedicine, Ethics, and Society. "Reproductive Laws for the 1990s." Ed S Cohen, N Taub. (Pp x+472; \$39.50.) New Jersey: Humana Press, 1989. ISBN 0-89603-157-8.

Reproductive Endocrine Therapeutics. Ed R L Barbieri, I Schiff. (Pp xii+430; figs; \$59.50.) New York: Liss, 1988. Distributed by John Wiley and Sons. ISBN 0-8451-4260-7.

Laboratory medicine

Advances in Biotechnological Processes. Vol 10. "Synthetic Peptides in Biotechnology." Volume and series editor A Mizrahi. (Pp xi+239; figs; \$69.50.) New York: Liss, 1989. Distributed by John Wiley and Sons. ISBN 0-8451-3209-1.

Miscellaneous

Research Publications of the Wellcome Unit for the History of Medicine, Oxford. No IX. "Medicine and Film: a Checklist, Survey and Research Resource." M Shortland. (Pp vi+56; figs; £6.95 paperback.) 1989. Available from The Wellcome Unit for the History of Medicine, 45-47 Banbury Road, Oxford OX2 6PE.

Take It and Leave It: Aspects of Being Ill. R Rubinstein. (Pp 124; £8.95.) London: Marion Boyars, 1989. ISBN 0-7145-2883-8.

WHO Offset Publication No 104. "Agriculture-Health Linkages." M Lipton, E de Kadt. (Pp 111; Sw frs 17 paperback.) Geneva: World Health Organisation, 1988. ISBN 92-4-170104-8.

World Health Organisation Technical Report Series. No 771. "WHO Expert Committee on Biological Standardisation." Thirty-eighth Report. (Pp 221; Sw frs 26 paperback.) Geneva: World Health Organisation, 1988. ISBN 92-4-120771-X.