exercise must—probably for ever—remain controversial.¹⁹ But this is a sterile debate: who would deny that stopping smoking, taking more exercise, controlling high blood pressure, and having nutritional counselling to reduce excess body weight and needlessly high cholesterol concentrations are reasonable strategies in pursuing a healthier lifestyle?

What of those who already have evidence of coronary heart disease? Aspirin, thrombolysis, β blockers, angiotensin converting enzyme inhibitors, risk stratification by cardiological investigation, revascularisation by surgery or angioplasty, and transplantation are some of the factors that have generated justifiable optimism in cardiology during the past decade.²⁰ Yet with resources scarce and under pressure there can be no complacency about the monumental problem of managing coronary heart disease not just in Scotland but world wide. The cost of these techniques and treatments is the greatest stimulus to invest more modest resources in a healthier future. Without doubt, preventive strategies must be strengthened for those already known to have coronary heart disease, with the aim of preventing the progression of atheroma or even inducing regression,^{21 22} especially since the results of trials of secondary prevention have been encouraging.²³⁻²⁵

And primary prevention? We have to acknowledge that the major risk factors and socioeconomic influences explain only part of the overall prevalence of coronary heart disease; other possible mechanisms include climate, latitude, stress, water softness, and viruses.^{26 27} But even if half of all coronary heart disease is not explicable on the basis of current knowledge the other half leaves plenty of scope for action. Thanks to projects such as the Scottish heart health study and the Scottish MONICA study²⁸ we have more information on risk factors and the prevalence of coronary heart disease than anywhere else in the United Kingdom. Yet Scotland is the only country within the United Kingdom that has no national programme aimed specifically at coronary heart disease. The priority that coronary heart disease deserves in Scotland has already been recognised in several local initiatives such as "Good Hearted Glasgow" in its programme of public education and individual counselling. In a population with high cholesterol values nutritional change is the key,²⁹ while the contribution of lipid lowering drugs will become clearer as the results of prevention studies become available.

A holistic approach, as envisaged by the Scottish Health Education Group, has advantages over one that is disease specific.³⁰ Firstly, it addresses other common preventable diseases whose risk factors or markers overlap with those for coronary heart disease. Secondly, it does address the risk factors, albeit indirectly; and, thirdly, it should be supported by those who remain sceptical about the results of trials of risk factor modification in coronary heart disease but who nevertheless accept that their control is in line with the promotion of a healthy lifestyle.

What is required now is a much more vigorous and committed effort in health promotion that recognises the priority of coronary heart disease in Scotland. This population approach should be complemented by a programme to detect and manage people at high risk—justified by the dramatic reduction in risk that can be achieved by simultaneous attention to nutritional counselling and cigarette smoking on an individual basis.^{17 31} The complexities of the direct association between ill health and poor social circumstances should be seen as a challenge, not a deterrent, to everyone concerned in health promotion.³²

Finally, any serious commitment to the problem of coronary heart disease in Scotland requires adequate funding. The results of the Scottish heart health study have shown where extra resources might be needed most. Scotland needs

the diversion of a fraction of that fervour so evident in the sporting arenas of the world towards creating a healthier national lifestyle. Winning the international league championships for coronary heart disease year after year is becoming boring; this is one title the Scots can afford to lose.

HENRY J DARGIE

Consultant Cardiologist,
Western Infirmary,
Glasgow G11 6NT

- 1 Uemura K, Pisa Z. Trends in cardiovascular mortality in industrialised countries since 1950. *World Health Stat Q* 1988;41:155.
- 2 Office of Population Consensus and Surveys. *Mortality statistics*. London: HMSO, 1987. (Series DHI.)
- 3 Registrar General for Scotland. *Annual report 1988*. Edinburgh: HMSO, 1988.
- 4 Fulton M, Adams W, Lutz W, Oliver MF. Regional variations in mortality from ischaemic and cardiovascular disease in Britain. *Br Heart J* 1978;40:563.
- 5 Logan RJ, Thomson M, Riemersma R, et al. Risk factors for ischaemic heart disease in normal men aged 40. Edinburgh-Stockholm study. *Lancet* 1978;i:949.
- 6 Wood DQA, Riemersma R, Oliver MF, et al. Adipose tissue and platelet fatty acids and coronary heart disease in Scottish men. *Lancet* 1984;iii:117.
- 7 Smith WCS, Crombie IK, Tavendale R, Irving JM, Kenicer MB, Tunstall-Pedoe H. The Scottish heart study: objectives and development of methods. *Health Bull (Edinb)* 1987;45:211-7.
- 8 World Health Organisation European Collaborative Group. Multifactorial trial in the prevention of coronary heart disease: recruitment and initial findings. *Eur Heart J* 1980;1:73-80.
- 9 Puska P, Salonen JT, Nissinen A, et al. Change in risk factors for coronary heart disease during 10 years of a community intervention programme. *Br Med J* 1983;287:1840-4.
- 10 Smith WCS, Tunstall-Pedoe H, Crombie IK, Tavendale R. Concomitants of excess coronary deaths—major risk factor and lifestyle findings from 10,359 men and women in the Scottish heart health study. *Scott Med J* 1989;34:550-5.
- 11 Tunstall-Pedoe H, Smith WCS, Crombie IK, Tavendale R. Coronary risk factor and lifestyle variation across Scotland: results from the Scottish heart health study. *Scott Med J* 1989;34:556-60.
- 12 Shaper AG, Pocock SJ, Walker M, Cohen NM, Wale CJ, Thomson AG. British regional heart study: cardiovascular risk factors in middle-aged men in 24 towns. *Br Med J* 1981;283:179-86.
- 13 WHO-ERICA Project. The CHD risk map of Europe. *Eur Heart J* 1988;9(suppl 1).
- 14 Tunstall-Pedoe H. Autres pays, autres moeurs: theories on why the French have less heart disease than the British. *Br Med J* 1988;297:1559.
- 15 Gey KF, Stahelin HB, Puska P, Evans A. Relationship of plasma level of vitamin C to mortality from ischaemic heart disease. *Ann NY Acad Sci* 1987;498:110.
- 16 Neaton JD, Kuller LH, Wentworth D, Borhani NO. Total and cardiovascular mortality in relation to cigarette smoking, serum cholesterol concentrations and diastolic blood pressure among black and white males followed up for five years. *Am Heart J* 1984;108:729.
- 17 Hjermann I, Holme I, Velve Byre K, Leren P. Effect of diet and smoking intervention on the incidence of coronary heart disease. *Lancet* 1981;ii:1303.
- 18 Stallones RE. Mortality due to ischaemic heart disease: observations and explanations. *Atherosclerosis Reviews* 1982;9:43.
- 19 Stamler J. Doing the right things. *N Engl J Med* 1986;312:1053.
- 20 Julian DG, ed. *Diseases of the heart*. London: Baillière Tindall, 1989.
- 21 Arntzenius AC, Kromhout D, Barth JD, et al. Diet lipoproteins and the progression of coronary atherosclerosis. *N Engl J Med* 1985;312:805.
- 22 Blankenhorn DH, Nessim SA, Johnson RL, et al. Beneficial effects of combined colestipol niacin therapy on coronary atherosclerosis and coronary venous bypass grafts. *JAMA* 1987;257:3233.
- 23 Kornitzer M, de Backer G, Dramaix M, et al. Belgian heart disease prevention project: incidence and mortality results. *Lancet* 1983;ii:1066.
- 24 MRFIT Research Group. Exercise electrocardiogram and coronary heart disease mortality in the multiple risk factor intervention trial. *Am J Cardiol* 1985;55:16.
- 25 Daly L, Mulcahy R, Graham IM, Hickey N. Long term effect on mortality of stopping smoking after unstable angina and myocardial infarction. *Br Med J* 1983;287:324.
- 26 Gotto AM. Risk factors for coronary disease. In: Braunwald E ed, *Heart disease*. Philadelphia: W B Saunders, 1988:1153.
- 27 Benditt EP, Barrett T, McDougall JK. Viruses in the etiology of atherosclerosis. *Proc Nat Acad Sci USA* 1983;80:6386.
- 28 World Health Organisation Principal Investigators (prepared by Tunstall-Pedoe H) The World Health Organisation MONICA project (monitoring trends and determinants in cardiovascular disease): a major international collaboration. *J Clin Epidemiol* 1988;41:105-14.
- 29 Tunstall-Pedoe H, Smith WCS, Tavendale R. How-often-that-high graphs of serum cholesterol: findings from the Scottish heart health and Scottish MONICA studies. *Lancet* 1989;ii:540.
- 30 Scottish Health Education Group. *Be all you can be*. Edinburgh: Scottish Health Education Group, 1986.
- 31 Shaper AG, Pocock SJ, Phillips AN, Walker M. Identifying men at high risk of heart attacks: strategy for use in general practice. *Br Med J* 1986;293:474.
- 32 Black D. Inequalities in health: report of a working group. London: HMSO, 1980.

Osteoarthritis

Diagnosis is not as simple as it seems

Osteoarthritis costs Britain millions of pounds annually. It is the most common of the rheumatic diseases, results in many days lost from work, puts more on the national drug bill than does any other arthritic disease, and is the most frequent reason for arthroplasty. It is second only to cardiovascular

disease in producing severe disability, and it affects about 10% of the population over the age of 60.¹

For all these reasons it should have provoked a wealth of research, and at long last it is doing so.^{2,3} As a topic it is fraught with difficulties. Even the simple question, "What do we mean by osteoarthritis?" remains unanswered. Recently the American Rheumatism Association published a set of criteria for reporting osteoarthritis of the knee⁴ and in a preliminary way of the hip and the hand.⁵ Features said to have the most discriminatory power included age over 50, crepitus, bony enlargement, morning stiffness of less than 30 minutes, and osteophytosis—the only radiographic predictor.

The shortcomings of the study have been reviewed by McAlindon and Dieppe.⁶ The controls were not matched for age or sex, were younger than the patients with osteoarthritis, and included many patients with rheumatoid arthritis. This may be the reason why age and osteophytosis appeared to be important discriminators and narrowing of the joint space was not.

There are good reasons for believing that osteophytes are an independent, age related variable.⁷ In some groups, such as professional footballers, traction osteophytes may occur as a physiological response rather than a degenerative process.⁸ The prevalence of features such as crepitus in the population is unknown. Features that seemed to be important in the recognition of osteoarthritis were largely subjective and have not been validated.

In epidemiological studies based on radiology the correlation between radiographic changes and symptoms is poor—apart from in the hip.⁹ Moreover, the distinction between primary osteoarthritis and conditions predisposing to secondary osteoarthritis is by no means firm. There are many clinical variants, such as primary generalised osteoarthritis, erosive inflammatory osteoarthritis, diffuse idiopathic skeletal hyperostosis, and chondromalacia patellae.^{10,11} Nor is it always clear which changes are primary and which are secondary. The fat lady who implores the orthopaedic surgeon for a hip operation assures him that inactivity due to arthritis explains her obesity. Some evidence suggests that the reverse is true, at least for arthritis in the knee.¹²

One tempting theory is the suggestion that the deposition of calcium pyrophosphate with chondrocalcinosis is a cause of osteoarthritis, but studies of hypermobile patients have suggested that it is secondary—at least in those who developed premature osteoarthritis.¹³ These issues are important in considering the biology of osteoarthritis (a subject reviewed in detail recently by Hamerman),¹⁴ and they have obvious therapeutic implications.

The clinician's first concern is to diagnose the condition. When swelling is bony, synovitis is minimal, and acute phase reactants are normal there is little problem. But often life is not so simple. Osteoarthritis with deposition of calcium pyrophosphate may mimic rheumatoid arthritis,¹⁵ and so may acute generalised osteoarthritis in the elderly. Helpful clinical features that support the diagnosis of an inflammatory polyarthritis rather than osteoarthritis are disease affecting the ulnar styloid and synovitis of the radiohumeral, glenohumeral, and second, third, and fourth metatarsophalangeal joints. A raised erythrocyte sedimentation rate or plasma viscosity and raised concentrations of rheumatoid factors are useful pointers to rheumatoid arthritis. Radiological signs of erosion of the metacarpophalangeal and metatarsophalangeal joints and the ulnar styloid would also favour this diagnosis. The synovial fluid of osteoarthritis has characteristics of a transudate rather than an exudate, with a high viscosity and relatively few cells, those present being lymphocytes rather than polymorphs.

The syndrome of osteoarthritis is often the end result of a series of processes, and, as the American experience has

shown, definition is not easy. Yet it is vital if appropriate treatment strategies are to be devised.

V WRIGHT

ARC Professor of Rheumatology,
University Department of Medicine,
General Infirmary,
Leeds LS1 3EX

- 1 Peyron JG. Osteoarthritis: the epidemiologic viewpoint. *Clin Orthop* 1986;213:113-9.
- 2 Wright V, ed. *Clinics in rheumatic diseases. Osteoarthritis*. Eastbourne: Saunders, 1976.
- 3 Sokoloff L, ed. *Clinics in rheumatic diseases. Osteoarthritis*. Philadelphia: Saunders, 1985.
- 4 Altman R, Asch E, Bloch D, et al. Development of criteria for the classification for the reporting of osteoarthritis: classification of osteoarthritis of the knee. *Arthritis Rheum* 1986;29:1039-49.
- 5 Altman RD, Bloch DA, Bole GG, et al. Development of clinical criteria for osteoarthritis. *J Rheumatol* 1987;14(suppl 1):3-6.
- 6 McAlindon T, Dieppe P. Osteoarthritis: definitions and criteria. *Ann Rheum Dis* 1989;48:531-2.
- 7 Wood PHN. Osteoarthritis in the community. *Clinics in Rheumatic Diseases* 1976;2:495-507.
- 8 Adams ID. Osteoarthritis and sport. *Clinics in Rheumatic Diseases* 1976;2:523-41.
- 9 Lawrence JS, Brenner JM, Bier F. Osteoarthritis, prevalence with population and relationship between symptoms and x-ray changes. *Ann Rheum Dis* 1966;25:1-24.
- 10 Mankin HJ, Brandt KD, Shulman LE. Workshop on etio-pathogenesis of osteoarthritis: proceedings and recommendations. *J Rheumatol* 1986;13:1130-60.
- 11 Moskowitz RW. Introduction. In: Moskowitz RW, Howell DS, Goldberg VM, Mankin HJ, eds. *Osteoarthritis: diagnosis and management*. Philadelphia: W B Saunders, 1984:1-6.
- 12 Felson T, Anderson JJ, Naimark A, Walker AM, Meenan RF. Obesity and knee osteoarthritis: the Framingham study. *Ann Intern Med* 1988;109:18-24.
- 13 Bird HA, Tribe CR, Bacon PA. Joint hypermobility leading to osteoarthritis and chondrocalcinosis. *Ann Rheum Dis* 1978;37:203-11.
- 14 Hamerman D. The biology of osteoarthritis. *N Engl J Med* 1989;320:1322-30.

Patient power in mental health

Working for users?

The difference between the provider of health services and the patient has been said to "resemble that between the hen and the pig in the preparation of eggs and bacon: the hen is involved but the pig is committed."¹ One way in which the patient's commitment is being acknowledged is "user participation," a jargon phrase that means that patients lead the way in planning new services. Patients want to participate actively in health care rather than be passive recipients,² and there is evidence that this will improve outcomes.³ But "user participation" has also been advocated for ethical or even political reasons.⁴ Here confusion begins.

Conflicting views of the patient's role in health care have been apparent for over a decade.⁵ Some pressure groups envisage active "consumers" such as those described by Nader.⁶ These groups talk about "empowerment," enhanced self respect derived from having a recognised role in services.⁷ In contrast the consumers described in the first Griffiths report were patients who exercised their influence obliquely by looking at the services on offer, choosing and trying, and then complaining if necessary.⁸ Emphasis on "patient choice" in *Working for Patients* still reflects this view.⁹

The concept of user participation is of particular importance in mental health, where the doctor-patient relationship may be an important part of therapy.¹⁰ A broad range of groups with interests in mental health already exists: charities, support groups for patients and families, and pressure groups. Some accept a biological view of mental disorder, some do not, and some want no contact with formal psychiatry.⁷ Yet across this diversity there is increasing emphasis on what the patient wants. For example, over the past year both MIND (the National Association for Mental Health) and the National Schizophrenia Fellowship have launched groups run by users for users.¹¹

The consumer network of MIND lists over 70 "user groups."¹² In Chesterfield user groups are self run with financial sponsorship from social services.¹³ In Nottingham groups participate in planning mental health services. They