Association. Goya Wannamethee is supported by the Institute of Alcohol Studies. We thank Dr Andrew Phillips and Mrs Mary Walker for their helpful advice in preparing this paper and other members of the department for their comments and suggestions.

- 1 Royal College of Physicians. Obesity. *J R Coll Physicians Lond* 1983;17:5-65.
- 2 Borhani NO, Hector HH, Breslow L. Report of a ten-year follow-up of the San Francisco longshoremen. *J Chronic Dis* 1963;16:1251-66.
- 3 Tibblin G, Wilhelmsen L, Werkö L. Risk factors for myocardial infarction and death due to ischaemic heart disease and other causes. *Am J Cardiol* 1975;35:514-22.
- 4 Society of Actuaries. *Build and blood pressure study*, 1959. Chicago: Society of Actuaries, 1959.
- 5 Lew EA. Mortality and weight: insured lives and the American Cancer Society studies. *Ann Intern Med* 1985;103:1024-9.
- 6 Jarrett RJ, Shipley MJ, Rose G. Weight and mortality in the Whitehall study. *Br Med J* 1982;285:335-7.
- 7 Dyer AR, Stamler J, Berkson DM, Lindberg HA. Relationship of relative weight and body mass index to 14-year mortality in the Chicago people gas company study. *J Chronic Dis* 1975;28:109-23.
- 8 Gordon T, Doyle JT. Weight and mortality in men: the Albany study. *Int J Epidemiol* 1988;17:77-81.
- 9 Sorlie P, Gordon T, Kannel WB. Body build and mortality. The Framingham study. *JAMA* 1980;243:1828-31.
- 10 Waaler HT. Height, weight and mortality: the Norwegian experience. *Acta Med Scand [Suppl]* 1984;679:1-56.

- 11 Vandembroucke JP, Morritz B, Bruin A, Verhessen H, Heide WC. Weight, smoking and mortality. *JAMA* 1984;252:2859-60.
- 12 Rissanen A, Heliovaara M, Knekt P, Aromaa A, Reunanen A, Maatela J. Weight and mortality in Finnish men. *J Clin Epidemiol* 1989;42:781-9.
- 13 Manson JE, Stampfer MJ, Hennekens CH, Walter C, Willett C. Body weight and longevity: a reassessment. *JAMA* 1987;257:353-8.
- 14 Shaper AG, Pocock SJ, Walker M, Cohen NM, Wale CJ, Thomson AG. British Regional Heart Study: cardiovascular risk factors in middle-aged men in 24 towns. *Br Med J* 1982;283:179-86.
- 15 Walker M, Shaper AG. Follow-up of subjects in prospective studies in general practice. *J R Coll Gen Pract* 1984;34:365-70.
- 16 Shaper AG, Weatherall R. Overweight and obesity in middle-aged men. *European Journal of Clinical Nutrition* 1988;42:221-31.
- 17 Simopoulos AP, Van Itallie TB. Body weight, health, and longevity. *Ann Intern Med* 1984;100:285-95.
- 18 Garrison RJ, Feinleib M, Castelli WP, McNamara P. Cigarette smoking as a confounder of the relationship between relative weight and long term mortality. *JAMA* 1983;249:2199-203.
- 19 Rhoads GG, Kagan A. The relation of coronary disease, stroke and mortality to weight in youth and middle age. *Lancet* 1983;i:492-5.
- 20 Hoffmans MDA, Kromhout D, Coulander C. The impact of body mass index of 78 612 18-year old Dutch men on 32 year mortality from all causes. *J Clin Epidemiol* 1988;44:749-56.
- 21 Selzer F. Measurement of overweight. *Statistical Bulletin* 1984;65:20-4.
- 22 Andres R, Elahi D, Tobin JD, Muller DC, Brant L. Impact of age on weight goals. *Ann Intern Med* 1985;103:1030-3.

(Accepted 19 October 1989)

Sympathetic tone and relation between sodium intake and blood pressure in the general population

J/Staessen, C J/Bulpitt, L/Thijs, R/Fagard, J V/Joossens, R/Van Hoof, A/Amery

Hypertension and Cardiovascular Rehabilitation Unit, Department of Pathophysiology, University of Louvain, Louvain, Belgium

J Staessen, MD, lecturer
L Thijs, BS, statistician
R Fagard, MD, professor
J V Joossens, MD, professor
R Van Hoof, MD, research assistant
A Amery, MD, professor

Division of Geriatric Medicine, Department of Medicine, Royal Postgraduate Medical School, Hammersmith Hospital, London
C J Bulpitt, MD, professor

Correspondence to: Dr J Staessen, Klinisch Laboratorium Hypertensie, Inwendige Geneeskunde-Cardiologie, UZ Gasthuisberg, Herestraat 49, B-3000 Leuven, Belgium.

Br Med J 1989;299:1502-3

In a random sample of the population we tested the hypothesis of Poulter *et al* that sodium intake and sympathetic tone interact to determine blood pressure.¹ As in Poulter *et al*'s study, the pulse rate at rest was taken as an index of sympathetic tone and the urinary excretion of sodium as an index of salt consumption.

Subjects, methods, and results

We studied 2450 subjects aged 18 and over who were randomly selected from the population.² Everybody was visited twice at home and on each visit the pulse rate over one minute and five blood pressure measurements were obtained. The results were averaged for analysis.² Twenty four hour urine collections were analysed for sodium and creatinine. People taking antihypertensive drugs ($n=296$) and those who collected more or less urine than required² ($n=73$) were removed from the analysis. As in Poulter *et al*'s study,¹ the remaining 2081 subjects were divided according to sex and the two groups stratified into three according to pulse rate. Significant blood pressure covariates were traced by stepwise multiple regression. Blood pressure adjustments for age included both a linear and a quadratic term. Coincidence of regression lines was investigated by a multiple regression approach with two dummy variables identifying the pulse rate thirds.

After adjustment for sex, age, and body mass index there was a significant curvilinear relation between urinary excretion of sodium over 24 hours and both systolic and diastolic pressure in the upper third of pulse rate ($n=735$) (figure). In the middle third of pulse rate ($n=694$) the correlations between blood pressure and urinary sodium excretion were not significant. In the lowest third of pulse rate ($n=652$) urinary sodium excretion was again curvilinearly related to both systolic and diastolic blood pressure (figure).

The regression lines adjusted for sex, age, and body mass index were not coincident across the pulse rate

thirds ($p<0.001$ for test of coincidence). Indeed, the linear and quadratic regression coefficients for urinary sodium excretion, but not the intercepts, were different in the upper third of pulse rate compared with the two others ($F_{2,2071}=18.4$ and 25.7 for systolic and diastolic pressure respectively; $p<0.001$). Additional adjustments for smoking, consumption of alcohol, and use of the contraceptive pill did not remove these differences.

Curvilinear relations between urinary sodium excretion and systolic and diastolic blood pressures (adjusted for sex, age, and body mass index) in three pulse rate thirds (fast, intermediate, and slow). For each regression line 95% confidence interval for population mean, correlation coefficient (r), F value, and associated p value are given

Comment

In our study less than 2% of the variance in blood pressure was explained by changes in the urinary excretion of sodium. Nevertheless, the finding that a raised blood pressure was associated with the combination of a fast pulse rate and a high urinary excretion of sodium is compatible with the suggestion of Poulter *et al* that a high sympathetic tone is a prerequisite for sodium to act as a pressor substance.¹ Alternatively, a high intake of sodium may facilitate the pressor effect of sympathetic stimulation. In our study smoking and consumption of alcohol were ruled out as confounding factors, but no information was available on the subjects' psychological stress and usual physical activity and therefore a possible influence of these two factors cannot be excluded.

Subjects with a slow pulse rate might have a low sympathetic tone; if their intake of sodium increases from 50 to 300 mmol/24 h sympathetic tone might be inhibited further and the secretion of renin suppressed, resulting in a fall in blood pressure. In contrast, people

with a fast pulse rate might have an enhanced sympathetic tone and less effective baroreflexes; if their intake of sodium increases above 160 mmol/24 h the high sympathetic tone might prevent the inhibition of renin secretion and impair the ability of the kidney to excrete sodium,⁴ thus causing the blood pressure to rise. Increasing the intake of sodium may therefore have opposite effects depending on the prevailing sympathetic tone.

- 1 Poulter NR, Shipley MJ, Bulpitt CJ, Markowe HJ, Marmot MG. Pulse rate and twenty-four hour urinary sodium content interact to determine blood pressure levels of male London civil servants. *J Hypertens* 1988;6(suppl 4):611-3.
- 2 Staessen J, Bulpitt CJ, Fagard R, Joossens JV, Lijnen P, Amery A. Salt intake and blood pressure in the general population: a controlled intervention trial in two towns. *J Hypertens* 1988;6:965-73.
- 3 Floras JS, Sleight P. The lability of blood pressure. In: Amery A, Fagard R, Lijnen P, Staessen J, eds. *Hypertensive cardiovascular disease: pathophysiology and treatment*. The Hague: Martinus Nijhof, 1982:104-17.
- 4 Light KC, Koepke JP, Obrist PA, Willis PWIV. Psychological stress induces sodium and fluid retention in men at high risk for hypertension. *Science* 1983;220:429-31.

(Accepted 15 September 1989)

Quality of life of elderly people after surgery for benign oesophageal stricture

B N M Jayawardhana, K Moghissi, J Knox

Department of Medicine for the Elderly, Kingston General Hospital, Hull
B N M Jayawardhana, MRCP, registrar
J Knox, FRCP, consultant physician

Cardiothoracic Unit, Castle Hill Hospital, Cottingham, Hull
K Moghissi, FRCS, consultant surgeon

Correspondence to: Dr Jayawardhana, Department of Medicine for the Elderly, Barnet General Hospital, High Barnet, Herts EN5 3DJ.

Br Med J 1989;299:1503-4

Benign oesophageal stricture in elderly patients is usually treated by intermittent dilatation and drug treatment to control continuing symptoms of reflux.^{1,2} But this treatment has complications.³ We assessed the results of surgery for stricture in elderly patients by using a validated quality of life scale⁴ in patients in whom endoscopic dilatation was unsuccessful or impossible because of the severity of the stricture.

Patients, methods, and results

We assessed consecutive operations performed between 1971 and 1983 for histologically confirmed benign oesophageal stricture in 124 patients (85 women, 27 men) aged 70 and over (range 70-89, mean 76; 16 patients were over 80). Sixty eight patients were alive, 44 had died, and 12 were excluded from the study because hospital records were untraceable for nine and death records unavailable for three. Surgery was performed by KM and the survivors assessed independently by BNMJ.

Twenty four patients had medical disorders that

were likely to influence the outcome of surgery. Six had a previous history of carcinoma. Forty five had had a laparotomy and 13 repairs of hiatus hernia and gastro-oesophageal reflux. Seventeen were taking non-steroidal anti-inflammatory drugs.

Intraoperative guided dilatation and an antireflux procedure were performed in 79 patients transthoracically. Thirty two required resection of the stricture because it was difficult to carry out the dilatation, or because of deep penetrating ulcer, or because of scarring with considerable oesophageal shortening. A Celestin tube was inserted in one patient when a malignancy was suspected.

Twelve patients died after operation (five from cardiac causes and two from pneumonia); seven had had conservative surgery and five had had resections (not significant 0.5>p>0.1); only one had had previous surgery on the oesophagus. Four further patients were too frail to return home and died in hospital. Twenty eight others died two months to 10 years (mean 42.5 months) after surgery. Five year mortality was 32%.

The state of health of the 68 survivors was assessed from two to 14 years (mean 65.5 months) after surgery according to the index of Fanshel and Bush (table): 41 were in groups S_A, S_B, and S_C, 13 having occasional symptoms. Minor and major disabilities were caused by heart disease, arthritis, old age, or memory impairment. The patient in group S_F had had a myocardial infarction and was expected to return to group S_A. The patients in group S_G were in wheelchairs because of arthritis. The one patient in an institution had dementia. Eight patients continued to receive drugs that were likely to cause local inflammation.

Fifty four of the 68 patients had no dysphagia. Six showed no improvement from the preoperative grade. Only two required further intermittent dilatation. Fifty two had no symptoms of reflux, but six were taking antacids, cimetidine, metoclopramide, or anticholinergic drugs alone or in combination. There was no significant difference in the incidence of dysphagia (p<0.5) or symptoms of reflux (p<0.5) between the conservative and radical procedures. Two patients had symptoms of the gas bloat syndrome. None required further surgery.

Comment

Surgery relieved the symptoms of benign oesophageal stricture in 43 of the 68 patients whose

State of health of 68 survivors of surgery for oesophageal stricture according to Fanshel and Bush index⁴

Quality of life index ⁴	Asymptomatic	Reflux only	Dysphagia only*	Reflux and dysphagia*
S _A : wellbeing	21	3	2 (26)	1 (1)
S _B : dissatisfaction (group that includes most of the population)	5	2	1 (8)	0
S _C : discomfort (minor symptoms that do not appreciably alter efficiency)	2	4	0 (5)	0 (1)
S _D : minor disability (reduced efficiency)	8	1	3 (16)	4
S _E : major disability (severe reduction in expected level of performance)	1	0	1 (2)	0
S _F : disabled but ambulatory (yet able to contribute to society)	1	0	0	0
S _G : confined to home	5	1	1 (4)	0 (3)
S _H : confined and bedridden				
S _I : isolated and in institution	0	0	1 (1)	0
S _J : coma				
S _K : death				
Total No (%)	43 (63)	11 (16)	9 (13)	5 (7)

*No of patients with symptoms before operation in parentheses.