instances six, serotypes. In only two were the maximum titres against the serotypes responsible for the original infection.

Discussion

Our results suggest that the long-term prognosis after the acute renal lesion of leptospirosis is good. Thus in the British patients all the results of renal function tests were normal except for a slightly reduced creatinine clearance in one man; as the serum creatinine level and other results in him were normal the significance of this isolated finding is doubtful. Another man developed hypertension, but renal damage did not appear to be the cause. One group of Gurkha patients had normal creatinine clearances but all did not concentrate urine adequately, while in the others these findings were reversed; since each abnormality was confined to one group it seems likely that these discrepancies reflect incomplete dehydration and incomplete urine collection respectively.

As intravenous pyelograms were performed in only four patients, and renal biopsy in none, we cannot be certain that permanent minor anatomical damage to the kidneys has not occurred; but as the individual results were not influenced by the severity of the initial renal lesion or the time interval before reassessment it is unlikely that there is an active process present causing progressive destruction of renal tissue.

The finding of leptospiral agglutinins in the sera of the 12 patients examined is of interest, though in only two did the serological pattern parallel that of the original infection. Hart (1967), using the sensitized erythrocyte lysis test, has shown that a small proportion of soldiers serving in jungle areas in South-east Asia develop a rising titre of leptospiral antibodies without clinical evidence of infection, and our patients have served in this area on several occasions since their original illness. It is thus conceivable that they have acquired sub-clinical infections during this time and that this explains the serological findings. As, however, all our patients showed significant titres of agglutinins, and none of them had a clinical attack, which would be surprising if they were frequently exposed to infection, this explanation may be too facile. Further investigation of this point is required.

Summary

Renal function was studied in 44 soldiers who had contracted leptospirosis in South-east Asia up to 14 years previously. The findings suggest that renal function is now normal, and individual results do not reflect the severity of the initial renal damage or the time interval before reassessment.

The sera of 12 patients were examined for leptospiral agglutinins. All showed significant titres, but in only two were these against the serotype responsible for the original illness. These results may possibly reflect subclinical leptospiral infections contracted during later periods of service in the same area, but the present evidence is insufficient to allow firm conclusions to be drawn.

We wish to thank Dr. Gordon Smith, Director M.R.E., Porton, and Major-General R. J. G. Morrison, Consultant Physician to the Army, for suggesting and encouraging this investigation; Dr. L. H. Turner for the serological results and advice; Captain K. Hedges, R.A.M.C., for his help in arranging for the investigation of patients under his care; and the Director-General, Army Medical Services, for permission to use case records.

Requests for reprints should be sent to Major B. Simpson.

REFERENCES

- Alexander, A. D., Evans, L. B., Toussaint, A. J., Marchwicki, R. H., and McCrumb, F. R., jun. (1957). *Amer. J. trop. Med. Hyg.*, **6**, 871.
de Brito, T., *et al.* (1965). *Ibid.*, **14**, 397.
Gsell, O. (1952). *Leptospirosen*. Berne.
Hart, R. J. C. (1967). To be published.
Johnson, D. W. (1950). *Med. J. Aust.*, **2**, 724.
Levis, D. G. (1957). *Proc. roy. Soc. Med.*, **50**, 119.
Mackay-Dick, J., and Robinson, J. F. (1957). *J. roy. Army med. Cps.*, **103**, 186.
Penna, D., de Brito, T., Aguiar Pupo, A., Marcondes Machado, M., Ayroza Galvão, P. A., and Soares de Almeida, S. (1963). *Amer. J. trop. Med. Hyg.*, **12**, 896.
Ross Russell, R. W. (1959). *Ann. trop. Med. Parasit.*, **53**, 416.
Spinu, I., Topciu, V., Trinh Thi Hang Quy, Vo van Hung, Chau xuan Luong, and Le van Tuyen (1962). *Arch. roum. Path. exp.*, **21**, 763.
Van Theil, P. H. (1948). *The Leptospiroses*, Leiden.
Trimble, A. P. (1957). *Proc. roy. Soc. Med.*, **50**, 125.
Varley, H. (1962). *Practical Clinical Biochemistry*, 3rd ed., pp. 112, 151, 153. London.

Effect of Medical and Surgical Vagotomy on Intrinsic Factor Secretion

J. F. ADAMS,* V.R.D., M.D., F.R.C.P.ED., F.R.C.P.GLASG.; A. G. COX,*† M.D., F.R.C.S.ED.
ELIZABETH H. KENNEDY,*‡ B.SC.; J. THOMPSON,*§ M.B., M.R.C.P.GLASG.

Brit. med. J., 1967, **3**, 473-476

The recent development of in-vitro techniques for the assay of gastric intrinsic factor has made possible the study of the effects of various pharmacological and surgical manoeuvres on this facet of gastric secretory function. Using one such technique, we have studied the effects of pharmacological vagal blockade—"medical vagotomy"—in a group of patients with proved duodenal ulceration, and in some of these have also studied the effects of section of the vagus nerves.

Material and Methods

Twenty-six patients (23 men aged 22-60 and 3 women aged 42-48) with the clinical and radiological features of chronic duodenal ulceration were studied before operation, and further studies were made in 15 of them after operation. Vagotomy and pyloroplasty were performed on six men and one woman

and vagotomy and gastrojejunostomy on six men and two women. The augmented histamine test was carried out in 26 cases before operation and in 15 after operation. Medical vagotomy was carried out in 25 cases before operation. The insulin test was done in 15 cases after operation. The choice of pyloroplasty or gastrojejunostomy was determined by a randomization procedure. The completeness of surgical vagotomy was confirmed in all cases by an insulin test.

The augmented histamine test (Kay, 1953) was performed after an overnight fast. Gastric aspiration was maintained by

* Southern General Hospital and University Department of Surgery, Western Infirmary, Glasgow.

† Now Senior Lecturer in Surgery, Royal Postgraduate Medical School, London.

‡ Now Scientific Officer, National Engineering Laboratory, East Kilbride, Lanarkshire.

§ Now Lecturer in Dermatology, University and Western Infirmary, Glasgow.

continuous suction. After collection of basal secretion in 15-minute periods for one hour, 100 mg. of mepyramine maleate was injected intramuscularly. Thirty minutes later histamine acid phosphate, 0.04 mg./kg. body weight, was injected subcutaneously and gastric aspiration was continued for a further hour, again in 15-minute collection periods.

The medical vagotomy test (Gillespie and Kay, 1961) was carried out on another day. This differed from the standard augmented histamine test in that a combined intramuscular injection of 50 mg. of hexamethonium bromide, and 0.325 mg. of atropine, was given at the beginning of the test.

The insulin test was performed with a standard dose of 20 units of soluble insulin given intravenously after collection of three 15-minute specimens of spontaneous secretions. After the injection gastric aspiration was continued for two hours. Vagotomy was regarded as complete only if the acid concentration failed to rise 20 or more mEq/l. over the maximal basal level.

The volume and pH of all specimens of gastric juice were measured. The amount of acid was determined by titration to pH 7 with N/10 NaOH, phenol red being used as the indicator. Intrinsic factor assays were performed by the method of Gottlieb *et al.* (1965). Several sera containing antibody to intrinsic factor were used, all being of high potency. When the total binding exceeded 75% of the amount of radioactive cyanocobalamin added (usually 7.5 mμg.) the procedure was repeated with half the normal amount of neutralized gastric juice: if the total binding still exceeded 75% of the amount of radioactive cyanocobalamin added then the procedure was repeated with half the normal amount of neutralized gastric juice and double the normal amount of radioactive cyanocobalamin. For convenience we adopted the convention of expressing intrinsic factor activity in terms of units, one unit being taken as the specific intrinsic factor binding of 1 mμg. of radioactive cyanocobalamin.

The results were analysed by standard statistical methods.

Results

All results are given as hourly values, representing spontaneous secretion ("basal hour"), the augmented histamine response ("augmented histamine hour"), and the response to histamine after medical vagotomy ("medical vagotomy hour").

Effect of Histamine Stimulation

The mean values for volume, acid, intrinsic factor output, and intrinsic factor concentration were all significantly higher ($P < 0.01$) in the augmented histamine hour than in the basal hour (Table I).

Correlation between Acid and Intrinsic Factor Output

An insignificant correlation was found between the mean values for acid and intrinsic factor in the basal hours, both

before and after operation. However, a significant ($P < 0.01$) correlation was found between these two measurements in the augmented histamine hours both before and after operation, and also in the medical vagotomy hour (Table II).

TABLE II.—Correlation and Regression Equations

Series	Period	Regression of y (I.F.) on x (Acid)	No. of Patients	r
All patients	Histamine hour ..	$y = 3.085x + 60.045$	26	0.551
	Medical vagotomy hour ..	$y = 4.773x + 34.629$	25	0.585
Surgical group	Preoperative histamine hour ..	$y = 4.926x + 25.947$	15	0.693
	Medical vagotomy hour ..	$y = 7.419x + 15.0965$	14	0.751
	Postoperative histamine hour ..	$y = 7.759x + 18.652$	15	0.762

Effect of Medical and Surgical Vagotomy

The results are most conveniently considered in two groups. The first relates to the effect of medical vagotomy on the augmented histamine response. The second relates to the effect of surgical vagotomy on basal secretion and the augmented histamine response.

Medical Vagotomy.—The mean values for volume and acid were significantly less ($P < 0.01$) in the medical vagotomy hour than in the augmented histamine hour. Differences in the mean values for intrinsic factor output and concentration did not

TABLE III.—Values for Acid and Intrinsic Factor Secretions After Histamine and After Medical Vagotomy. The Differences Between the Respective Values are Shown as a Percentage of the Value in the Histamine Hour

Case No.	Augmented Histamine Hour		Medical Vagotomy Hour		Percentage Change	
	Acid (mEq)	I.F. (mμg. Units)	Acid (mEq)	I.F. (mμg. Units)	Acid	I.F.
1	32.6	19,110	13.7	9,564	-58	-50
2	23.6	9,170	14.3	6,698	-39.4	-27
3	19.6	10,351	7.0	4,096	-64.3	-60.4
4	21.8	10,875	14.0	4,326	-35.8	-60.2
5	52.6	16,311	34.2	10,523	-35.0	-35.5
6	41.3	15,524	23.1	12,173	-44.1	-21.6
7	73.6	19,023	44.3	14,827	-39.8	-22.1
8	29.9	12,078	4.5	10,875	-84.9	-9.3
9	43.2	10,619	20.2	6,919	-53.2	-34.8
10	39.2	18,195	17.6	5,275	-55.1	-71.0
11	28.3	15,000	13.5	5,300	-52.3	-64.7
12	7.3	10,931	4.3	10,174	-41.0	-6.9
13	48.2	16,593	26.6	17,929	-44.8	+ 8.1
14	21.8	6,233	23.9	11,510	+ 9.6	+ 84.6
15	24.4	12,765	14.5	12,705	-40.5	-52.7
16	16.3	11,223	7.7	7,554	-52.7	+ 13.2
17	20.8	13,984	10.3	13,937	-50.4	-45.9
18	15.0	17,280	13.4	20,613	-10.6	-19.3
19	36.1	35,838	21.1	30,125	-41.5	-42.5
20	56.8	29,796	31.4	13,969	-44.7	+ 1.1
21	43.9	18,589	16.7	9,466	-61.9	-24.8
22	23.9	6,406	20.0	7,270	-16.3	+ 47.8
23	36.2	12,412	18.8	38,280	-48.0	-41.4
24	57.5	45,021	42.4	27,381	-26.2	-14.9
25	24.8	17,485	21.1	6,088	-14.9	+ 56.6
26	34.8	14,802	19.8		-43.1	-58.8
All cases					-42.9	-20.6
All except Case 15					-43.0	-20.6

TABLE I.—Mean Hourly Values for Volume, Acid, Intrinsic Factor and Intrinsic Factor Concentration

	Basal Hour					Post-Histamine Hour					Medical Vagotomy Hour					Postoperative Basal Hour					Postoperative Post-Histamine Hour				
	No.	Vol. (ml.)	Acid (mEq)	I.F. (mμg. units)	I.F./ml.	No.	Vol. (ml.)	Acid (mEq)	I.F. (mμg. units)	I.F./ml.	No.	Vol. (ml.)	Acid (mEq)	I.F. (mμg. units)	I.F./ml.	No.	Vol. (ml.)	Acid (mEq)	I.F. (mμg. units)	I.F./ml.	No.	Vol. (ml.)	Acid (mEq)	I.F. (mμg. units)	I.F./ml.
Surgical All patients	No. Mean	26	126.84	6.29	3,680	31.16	26	299.77	33.60	16,370	55.94	25	181.6	19.36	12,702	71.32									
	S.D.		48.87	3.88	1,783	19.26		98.57	15.44	8,654	24.27		68.93	10.40	8,492	38.33									
Surgical All patients	S.E.	15	9.58	0.76	350	3.78	15	19.33	3.03	1,697	4.76	14	13.79	2.08	1,698	7.67	12					15			
	No. Mean		119.47	5.70	3,406	30.96		276.87	31.19	17,957	64.61		175.57	19.82	16,214	93.36									
Surgical All patients	S.D.		49.72	4.00	1,908	22.87		97.34	15.24	10,838	28.22		70.01	9.77	9,653	37.97									
	S.E.		12.84	1.03	493	5.91		25.13	3.94	2,798	7.29		18.71	2.61	2,580	10.15									

achieve significance, though the mean intrinsic factor output was lower and the concentration higher in the medical vagotomy hour. The detailed results (Table III) show the very marked individual variations in response.

Surgical Vagotomy.—The effect of surgery on basal secretion was a significant reduction in mean values for volume, acid, and intrinsic factor ($P<0.01$) and for intrinsic factor concentration ($P<0.05$). The effect of surgery on histamine-stimulated secretion was a significant reduction in mean values for volume, acid ($P<0.01$), and intrinsic factor output ($P<0.05$), but there was no significant change in mean values for intrinsic factor concentration. Patients having gastrojejunostomy did

Discussion

The object of this study was to determine and compare the effects of medical and surgical vagotomy on gastric secretory function with particular emphasis on intrinsic factor secretion, and discussion of the results is limited to these points.

During the course of the study Bitsch *et al.* (1966) published their findings on the effects of surgical vagotomy on gastric secretory function, and comparison of the two sets of results is desirable. The mean values for volume, acid, intrinsic factor, and intrinsic factor concentration found by us are less than those of Bitsch *et al.*, but the data given in their paper do not

TABLE IV.—Values for Acid and Intrinsic Factor for the Hour After Histamine Stimulation, Before and After Operation, and also During Medical Vagotomy. The Differences are Shown as a Percentage of the Value in the Preoperative Histamine Hour

Case No.	Preoperative Augmented Histamine Hour		Preoperative Medical Vagotomy Hour		Postoperative Augmented Histamine Hour		Percentage Change			
	Acid (mEq)	I.F. (mµg. Units)	Acid (mEq)	I.F. (mµg. Units)	Acid (mEq)	I.F. (mµg. Units)	Acid		I.F.	
							Due to M. Vag.	Due to Op.	Due to M. Vag.	Due to Op.
12	7.3	10,931	4.3	10,174	6.1	5,872	-41.0	-16.4	-6.9	-47.9
13	48.2	16,593	26.6	17,929	20.9	7,677	-44.8	-56.6	+8.1	-53.7
14	21.8	6,233	23.9	11,510	4.1	1,689	+9.6	-81.1	+84.6	-39.4
15	24.4	12,765	14.5	12,705	12.7	16,770	-40.5	-47.9		+31.3
16	16.3	11,223	7.7	12,705	1.2	1,878	-52.7	-92.6	+13.2	-83.2
17	20.8	13,984	10.3	7,554	4.1	4,175	-50.4	-80.2	-45.9	-70.1
18	15.0	17,280	13.4	13,937	8.2	14,497	-10.6	-45.3	-19.3	-16.1
19	36.1	35,838	21.1	20,613	5.8	3,226	-41.5	-83.9	-42.5	-90.9
20	56.8	29,796	31.4	30,125	24.6	24,871	-44.7	-56.6	+1.1	-16.5
21	43.9	18,589	16.7	13,969	19.6	15,961	-61.9	-55.3	-24.8	-14.1
22	23.9	6,406	20.0	9,466	10.8	5,091	-16.3	-54.8	+47.8	-20.5
23	36.2	12,412	18.8	7,270	5.0	4,630	-48.0	-86.1	-41.4	-62.6
24	57.5	45,021	42.4	38,280	14.6	18,166	-26.2	-74.6	-14.9	-59.6
25	24.8	17,485	21.1	27,381	8.7	12,711	-14.9	-64.9	+56.6	-25.0
26	34.8	14,802	19.8	6,088	8.6	11,036	-43.1	-75.2	-58.8	-25.4
All cases							-37.4	-66.8	-11.5	-44.9
All except case 15							-37.5	-67.9	-11.5	-48.7

not differ from those in whom pyloroplasty was employed as the drainage procedure. Detailed results (Table IV) again showed very marked individual variation in response. The pattern of response to histamine after operation was similar to that found during medical vagotomy (see Chart).

Secretory patterns in cases studied before and after surgery. The figures 5-8 signify the four 15-minute periods after the injection of histamine, 1-4 being the four 15-minute periods before the injection.

Comparison of Effects of Medical and Surgical Vagotomy.—Comparison of the mean hourly values after medical vagotomy and after histamine stimulation in the postoperative period showed that the mean values for volume and intrinsic factor did not differ significantly ($P>0.05$) but that the mean values for acid and intrinsic factor concentration were significantly less ($P<0.01$ and <0.05 respectively) after operation than after medical vagotomy.

permit calculation of the significance of these differences. The results can be compared, however, by taking the difference between the values obtained in the histamine hour before and after operation and expressing the difference as a percentage of the preoperative value. Bitsch *et al.* observed a volume reduction of 46.7%, an acid reduction of 62.5%, and a reduction in intrinsic factor output of 43.8% (not 34% as printed in their paper). The corresponding values in the present study are 40.8% for volume, 66.8% for acid, and 44.9% for intrinsic factor. These results are clearly in close agreement and confirm that surgical vagotomy and a drainage procedure effect a significant reduction in the output of gastric juice, acid, and intrinsic factor in response to histamine.

The type of drainage procedure employed does not appear to be important. In our series both groups responded in a similar manner and no significant differences were found between them in regard to volume, acid, or intrinsic factor output after operation. This suggests, but does not prove, that vagotomy is the effective agent.

The effect of medical vagotomy differs sharply from that of surgical vagotomy. The effect of medical vagotomy was a significant reduction in volume and acid output after histamine stimulation, as was expected (Gillespie and Kay, 1961), but an insignificant change in intrinsic factor output. This finding is of interest in relation to the mechanisms which control intrinsic factor secretion. Hoedemaeker *et al.* (1964) have produced evidence from immunological and autoradiographic studies that the site of secretion is the parietal cell, and this view is supported by the close correlation between amounts of acid and intrinsic factor secreted (Ardeman, 1965; Irvine, 1965; Rødbro, *et al.*, 1965). The pattern of acid and that of intrinsic factor secretion differ, however, the difference being most obvious on continued histamine stimulation when the output of acid remains high and that of intrinsic factor falls off rapidly after an initial rise (Irvine, 1966; Lawrie and Anderson, 1967). This indicates either that the two secretions come from different cells or that they come from the same cell, in which case the

secretion of acid is by a process of true stimulation and that of intrinsic factor is one of release or "washout." The importance of vagal control is clearly shown by the effect of surgical vagotomy as found by Bitsch *et al.* (1966) and confirmed here.

That medical vagotomy did not have a quantitatively significant effect on the histamine-stimulated output of intrinsic factor was therefore unexpected but is not necessarily evidence against the concept of vagal control of intrinsic factor secretion. The pattern of intrinsic factor secretion is affected by medical vagotomy in the same way as by surgical vagotomy, suggesting that both procedures exert an effect on the controlling mechanisms even if the effect is quantitatively insignificant with one but not with the other. There are two possible explanations. First the vagal component of acid secretion may be more dominant than the vagal control of intrinsic factor secretion, and since medical vagotomy is less complete than surgical vagotomy the quantitative effect will be correspondingly less. Other influences on intrinsic factor secretion, such as intragastric reflexes, may be more important than the vagal component. Secondly, the medical vagotomy differs from surgical vagotomy in that it fails to block histamine-stimulated washout effect while having a blocking effect on the pattern of active secretion stimulated by histamine.

Summary

The effect of medical and surgical vagotomy on intrinsic factor secretion was studied in patients with duodenal ulceration.

Surgical vagotomy brought a significant reduction in both basal and histamine-stimulated output, but medical vagotomy did not significantly affect histamine-stimulated output. The pattern of intrinsic factor secretion in response to histamine was affected in the same manner by medical and surgical vagotomy.

The results suggest either that the vagal component in the control of intrinsic factor secretion is less dominant than the vagal component in acid secretion or that medical vagotomy differs from surgical vagotomy, the former failing to block a washout type of pattern and the latter doing so.

We are grateful to Professor A. W. Kay for advice and encouragement, to Professor T. Ferguson Rodger for facilities in his department, and to Merck Sharp & Dohme Ltd. for supplies of radioactive cyanocobalamin. We acknowledge with thanks a grant from the Secretary of State for Scotland on the advice of the Advisory Committee on Medical Research.

REFERENCES

- Ardeman, S. (1965). *Proc. roy. Soc. Med.*, 58, 916.
 Bitsch, V., Christiansen, P. M., Faber, V., and Rødbro, P. (1966). *Lancet*, 1, 1288.
 Gillespie, I. E., and Kay, A. W. (1961). *Brit. med. J.*, 1, 1557.
 Gottlieb, C., Lau, K-S., Wasserman, L. R., and Herbert, V. (1965). *Blood*, 25, 875.
 Hoedemaeker, P. J., Abels, J., Wachters, J. J., Arends, A., and Nieweg, H. O. (1964). *Lab. Invest.*, 13, 1394.
 Irvine, W. J. (1965). *New Engl. J. Med.*, 273, 432.
 — (1966). *Clin. exp. Immunol.*, 1, 99.
 Kay, A. W. (1953). *Brit. med. J.*, 2, 77.
 Lawrie, J. H., and Anderson, N. M. (1967). *Lancet*, 1, 68.
 Rødbro, P., Christiansen, P. M., and Schwartz, M. (1965). *Ibid.*, 2, 1200.

Medical Memoranda

Hydatidiform Mole and Coexistent Foetus, both with Triploid Chromosome Constitution

Brit. med. J., 1967, 3, 476-478

Triploid chromosome complement in man has been described in abortions and also in three living subjects, all of whom were mosaics (Böök and Santesson, 1960; Penrose and Delhanty, 1961; Delhanty *et al.*, 1961; Carr, 1963, 1965; Ellis *et al.*, 1963; Ferrier *et al.*, 1964; Thiede and Salm, 1964; Aspillaga *et al.*, 1964; Szulman, 1965).

Atkin and Klinger (1962) and Szulman (1965) have reported a case of mole with an associated foetus in which triploidy was found in the molar tissue. Makino *et al.* (1964) found triploid chromosome complements in three cases of early abortion, where the chorionic villi were swollen and oedematous, forming small vesicles as in hydatidiform mole. Carr (1965) in his later series described a similar case. In none of these molar or molar-like gestations were the clinical features reported.

This paper describes the finding of triploid chromosome complement in a hydatidiform mole and its associated abnormal foetus, the mother showing antepartum and postpartum clinical features of molar pregnancy.

CASE REPORT

A 22-year-old woman developed severe pre-eclampsia at 17 weeks of gestation in her first pregnancy. She had palpitations, severe dyspnoea, and pain in the right iliac fossa when referred to the Royal Women's Hospital seven weeks later. The haemoglobin had

fallen from 13 to 7.1 g./100 ml., though vaginal bleeding had not occurred. The blood pressure was 170/110 and the urine contained 20 g. of protein per litre. The right lower abdomen was tender, but no mass was felt apart from the uterus, which reached to the level of the umbilicus. X-ray examination showed a foetal skeleton commensurate with 18 rather than 24 weeks of development; the plasma uric acid was 4.7 mg./100 ml.; urinary oestriol 1.5 mg./24 hours, and urinary chorionic gonadotrophin 300-400 i.u./ml.

After treatment with sedation and hypotensive drugs artificial rupture of the membranes was performed and 300 ml. of amniotic

FIG. 1.—Malformed foetus together with placenta showing diffuse macroscopic molar degeneration.