the standard retinal photographs have proved of practical worth. Five retinopathic abnormalities have been photographed in four standard degrees of severity, and the photographs under study were graded according to these standards. In this way not only was it possible to define deterioration or improvement, it was also possible to devise a simple quantitative assessment of the extent of the changes. With the exception of venous changes, observer errors in assigning grades was slight and results were found to be reproducible.

TABLE II.—Behaviour of Exudates in Control Groups of Similar Studies (Percentage of Total Eyes)

	Edinburgh (Cullen 1966)	Bristol (Marmion and Harrold, 1966*)	London (King <i>et al.</i> , 1963)	Present Study
Improved ..	16	15	12	14.2
No change ..	56	63	59	68
Deteriorated ..	28	22	29	17.8
Duration (years)	2			1

* Personal communication.

Methandienone was chosen as an agent known to be an effective anabolic steroid given by mouth, and the dosage of 5 mg. twice daily was sufficient to induce an average rise of serum cholesterol of 108 mg./100 ml. as compared with a rise of 43 mg./100 ml. in the control group. In an ailment such as diabetes mellitus, which is known to be associated with vascular disease, this cholesterol elevation must be regarded as an unwelcome effect, especially in the long term.

The effects of methandienone on diabetic retinopathy were compared with those of placebo tablets by means of the familiar double-blind technique in which neither patient nor observer was aware of the nature of the tablets used. Assessment of the changes at the end of 18 months seemed to show an increase of haemorrhages in patients taking methandienone, and though this was not statistically significant ($P > 0.12$), it made it highly unlikely that even over longer periods any benefits would accrue.

Summary

Changes in retinopathy were assessed in two groups of diabetics, one receiving 10 mg. of methandienone daily and the other a placebo. No beneficial effect was observed in the treated group.

Assessment of changes in retinopathy was made by serial retinal photographs. These were compared with sets of standard photographs. This method of assessment proved effective and reproducible and could be used for future trials.

We acknowledge with gratitude the receipt of a grant from the North-West Metropolitan Regional Hospital Board. We are indebted to Dr. Terry Binns, CIBA Laboratories, Horsham, for his help in designing the trial and providing methandienone (Dianabol) tablets and to Mr. Peter Smith, of the M.R.C. Department of Statistics, for assistance in assessing the results. Finally we are grateful to Professor Sir Max Rosenheim for allowing us to include patients attending University College Hospital.

REFERENCES

- Carr, J. J., and Drekter, I. J. (1956). *Clin. Chem.*, **2**, 353.
 Dardenne, U. (1961). *Acta endocr. (Kbh.)*, Suppl. No. 63, p. 143.
 Davis, M. D. (1965). *Arch. Ophthalm.*, **74**, 741.
 Fabrykant, M., Gelfand, M. L., and Rosenberg, A. S. (1964). *Amer. J. med. Sci.*, **248**, 304.
 Harrold, B. P. (1966). M.D. Thesis, University of London.
 Houtsmuller, A. J. (1961). *Acta endocr. (Kbh.)*, Suppl. No. 63, p. 154.
 King, R. C., Dobrce, J. H., Kok, D'A., Foulds, W. S., and Dangerfield, W. G. (1963). *Brit. J. Ophthalm.*, **47**, 666.
 Lee, P., McMeel, J. W., Schepens, C. L., and Field, R. A. (1966). *Amer. J. Ophthalm.*, **62**, 207.
 Leopold, I. H. (1964). In *Diabetes Mellitus: Diagnosis and Treatment*, edited by T. S. Danowski, p. 165. New York.
 Oakley, N. W., Kohner, E., Hill, D. W., Joplin, G. F., and Fraser, R. (1967). *Diabetologia*. In press.

Preliminary Communications

Fibrinolytic Activity in the Clinical Crisis of Sick-cell Anaemia

Brit. med. J., 1967, **3**, 653-654

The natural history of subjects with sickle-cell anaemia is punctuated by recurrent clinical crises. There is rarely any evidence of acute haemolysis in the vast majority of episodes of crisis (Diggs, 1956), and emphasis has been placed on the finding of a variety of thromboembolic manifestations affecting almost every body organ. It is inferred that there is a marked tendency towards spontaneous vascular occlusion in sickle-cell anaemia though the mechanism remains unknown. A generally accepted conclusion is that thrombotic crises arise as a consequence of intravascular agglutination of sickled cells leading to local thrombosis and tissue ischaemia (Allison, 1956). Walters (1958) has suggested, however, that infarction might be precipitated by factors likely to influence the coagulability of the plasma, with subsequent precipitation of a fibrin reticulum within vessels. Abnormalities in coagulation factors have been reported in subjects with sickle-cell anaemia (McKellar and Dacie, 1958; Henstell *et al.*, 1965), but their relation to clinical crisis has not been investigated.

The present study is an attempt to assess whether the fibrinolytic mechanism is impaired in sickle-cell anaemia, both during periods of apparent well-being and during episodes of clinical crisis.

PATIENTS AND METHODS

Children attending the sickle-cell anaemia clinic or admitted to the paediatric wards at Mulago Hospital, Kampala, were studied. All blood samples were obtained between the hours of 11 a.m. and 2 p.m., and a clean venepuncture was carried out with minimum stasis. Dilute blood clot lysis was determined with the technique of Fearnley *et al.* (1957) as modified by Lackner and Goosen (1959). The disintegration of the clot was taken as the end-point in this study, and clots were followed for a maximum of 24 hours. The following groups of patients were studied:

Group A (37).—Hospital inpatients with haemoglobin phenotype AA and with a wide variety of disorders. The duration of their stay in hospital preceding examination ranged from 4 to 42 (mean 9) days.

Group B (77).—Subjects with sickle-cell anaemia attending as outpatients and apparently well.

Group C (7).—Subjects with sickle-cell anaemia attending with mild symptoms attributable to the disease, but not severe enough to necessitate their admission to hospital.

Group D (13).—Subjects with sickle-cell anaemia with symptoms attributable to the disease and severe enough to require their admission to hospital. These children were usually bled within 48 hours of admission.

RESULTS

The distribution of the lysis times obtained in the four groups is shown in the Table. In testing for the significance of differences between any two groups, both the actual lysis time in hours and the logarithmic equivalent were used; differences between groups were equally significant whichever value was employed. The Mann-Whitney U test was used for assessing the significance of differences between the groups (Siegel, 1956).

Distribution of Lysis Time in Hours

Hours	A	B	C	D
1	3	2	1	0
2	6	15	0	1
3	3	19	3	2
4	5	13	3	1
5	2	6	0	0
6	2	7	0	0
7	3	1	0	0
8	2	3	0	0
9	1	3	0	0
10	1	2	0	0
11	0	0	0	1
12	0	0	0	0
13	0	0	0	0
14	0	1	0	0
15	0	1	0	0
16	0	1	0	0
17	2	0	0	0
18	1	0	0	0
19-23	0	0	0	0
24+	3	4	0	7
	37	77	7	13

The most striking finding is that the lysis time in group D—that is, sickle-cell anaemia subjects in clinical crisis admitted to the ward—is far greater than in group A ($P=0.008$), group B ($P=0.008$), or group C ($P=0.006$). There is a suggestion that lysis time in the apparently well sicklers is longer than in those with mild symptoms ($P=0.02$). Lysis time in the children with haemoglobin phenotype AA is slightly longer than in “well” sicklers ($P=0.05$) or in sicklers with mild symptoms ($P=0.04$), but it is very likely that this reflects the effect on fibrinolytic activity of the various illnesses for which these children were admitted to hospital and the medication they were receiving—for example, corticosteroids.

DISCUSSION

Fibrinolytic activity is significantly lower in patients with sickle-cell anaemia in whom symptoms attributable to this disorder were severe enough to necessitate their admission to hospital. If the severe symptoms associated with clinical crisis are due to intravascular thrombosis, it is possible that the fibrinolytic mechanism may be involved in either the development or the maintenance of such vascular occlusion. There is presumed to be a dynamic equilibrium between the deposition of fibrin in vessels and its removal by lysis (Astrup, 1956; Fearnley, 1961), and the marked diminution in fibrinolytic activity during episodes of sickle-cell crisis might either reflect some factor inhibiting normal enzyme activity or may represent the body's response to intravascular thrombosis and tissue damage.

High levels of fibrinogen have been reported in older subjects with sickle-cell anaemia in America, though the finding was

thought to be nonspecific. There appeared to be no difference in fibrinogen levels when patients were studied in or out of crisis (Fenichel *et al.*, 1950). This report of raised fibrinogen levels is commented upon because Blix (1961) asserted that clots made of plasma containing a high concentration of fibrinogen are more resistant to lysis than those formed from a lower fibrinogen concentration. However, Bennett *et al.* (1967) believe that the situation is far more complex than this and that changes in plasminogen activator, as assessed by euglobulin clot lysis, do not simply reflect changes in fibrinogen levels.

While it is possible that the subjects in sickle-cell crisis form a special group in whom there is a persistently reduced fibrinolytic activity, we think that this is very unlikely. It is more probable that activity is lowered during or before a severe clinical crisis. Four of the 13 patients studied in clinical crisis were studied again several months later, when they were apparently well. Of three subjects with long lysis times (24 hours) one remained prolonged and two had fallen to 3.9 and 3.3 hours respectively.

Several of the 13 children admitted in clinical crisis had haemoglobin levels below the level of those attending the clinic and not admitted to hospital. It has, however, previously been shown that the degree of anaemia per se does not affect fibrinolytic activity (Shaper *et al.*, 1967).

SUMMARY

Fibrinolytic activity has been shown to be significantly reduced in children with sickle-cell anaemia in clinical crisis as compared with sickle-cell anaemia subjects with no symptoms or with mild symptoms. Further studies are indicated to ascertain whether a disturbance of the fibrinolytic mechanism has a causal relationship to the signs and symptoms of the clinical crisis in sickle-cell anaemia.

We are grateful to Dr. M. C. Pike (Medical Research Council Statistical Research Unit) for his advice and assistance. This work is supported by grants from the Medical Research Council to A. M. and A. G. S. and by the World Health Organization.

A. MAHMOOD, PH.D., M.R.C.P.ED.,
Honorary Clinical Lecturer.

DORIS M. MACINTOSH, M.B., CH.B.,
Research Fellow.

A. G. SHAPER, M.B., M.R.C.P.,
Research Professor in Cardiovascular
Diseases.

Departments of Paediatrics and Medicine,
Makerere University College Medical School,
Kampala, Uganda.

REFERENCES

- Allison, A. C. (1956). *Trans. roy. Soc. trop. Med. Hyg.*, **50**, 185.
- Astrup, T. (1956). *Lancet*, **2**, 565.
- Bennett, N. B., Ogston, C. M., and Ogston, D. (1967). *Clin. Sci.*, **32**, 27.
- Blix, S. (1961). *Acta med. scand.*, **169**, 495.
- Dacie, J. V., (1960). *The Haemolytic Anaemias: Congenital and Acquired*, part 1, 2nd ed. London.
- Diggs, L. W. (1956). *Amer. J. clin. Path.*, **26**, 1109.
- Fearnley, G. R. (1961). *Lancet*, **1**, 992.
- , Balmforth, G., and Fearnley, E. (1957). *Clin. Sci.*, **16**, 645.
- Fenichel, R. L., Watson, J., and Eirich, F. (1950). *J. clin. Invest.*, **29**, 1620.
- Henstell, H. H., Kligerman, M., and Irwin, L. E. (1965). *Blood*, **25**, 907.
- Lackner, H., and Goosen, C. C. (1959). *Acta haemat. (Basel)*, **22**, 58.
- McKellar, M., and Dacie, J. V. (1958). *Brit. J. Haemat.*, **4**, 404.
- Shaper, A. G., Kyobe, J., and Mwanje, D. K. (1967). *Amer. J. clin. Path.*, **47**, 344.
- Siegel, S. (1956). *Nonparametric Statistics for the Behavioral Sciences*. New York.
- Walters, J. H. (1958). *Proc. roy. Soc. Med.*, **51**, 646.