REFERENCES

- Bach, J. F., Dardenne, M., and Fournier, C. (1969). *Nature*, **222**, 998.
 Beutner, E. H., and Jordon, R. E. (1964). *Proceedings of the Society for Experimental Biology and Medicine*, **117**, 505.
 Beutner, E. H., Jordon, R. E., and Chorzelski, T. P. (1968). *Journal of Investigative Dermatology*, **51**, 63.
 Coe, R. O., and Bull, F. E. (1968). *Journal of the American Medical Association*, **206**, 1515.
 Corley, C. C., jun., Lessner, H. E., and Larsen, W. E. (1966). *American Journal of Medicine*, **41**, 404.
 Ebringer, A., and Mackay I. R. (1969). *Annals of Internal Medicine*, **71**, 125.
 Krakowski A., Covo, J., and Rozanski, Z. (1969). *Archives of Dermatology*, **100**, 117.
 Lever, W. F., and Goldberg, H. S. (1969). *Archives of Dermatology*, **100**, 70.
 Lever, W. F., and White, H. (1963). *Archives of Dermatology*, **87**, 12.
 Muller, S. A., Farrow, G. M., and Martalock, D. L. (1969). *Archives of Dermatology*, **100**, 523.
New England Journal of Medicine, 1968, **278**, 277.
 O'Rourke, R. A., and Eckert, G. E. (1964). *Archives of Internal Medicine*, **113**, 191.
 Wolff, K., and Schreiner, E. (1969). *Archiv für klinische und experimentelle Dermatologie*, **235**, 63.

Medical Memoranda

Case of Ergot Poisoning

British Medical Journal, 1970, **3**, 86-87

Ergotamine tartrate is often prescribed in the treatment of migraine. Complications from the drug are rare but potentially serious, as is illustrated by the following case, where severe lower limb arterial spasm resulted from its use. This provided an opportunity to study the reversible vasoconstrictor effect by means of plethysmography, ultrasound, thermography, and arteriography.

CASE REPORT

A 28-year-old woman complained of intermittent claudication in the left calf of 18 months' duration. The distance at which calf pain occurred varied from 100 yards to 1 mile (90 to 1,600 m.). One month before her admission this distance had on occasions been as short as 25 yards (23 m.), and both calves were affected. On admission she had severe paraesthesiae in the toes and soles of both feet. Her legs felt "icy cold and dead." On direct questioning she admitted to suffering from headaches for several years. These had been previously diagnosed as migraine and became more frequent under stress. They were characterized by pains in the back of the neck radiating over the left temple. There was occasional blurring of vision and vomiting but no warning aura. She was given Migril and Cafergot suppositories to be self-regulated during the attacks. The attacks had been more severe during the preceding few months with the result that two or three times the prescribed dose was taken. She admitted that the claudication was exacerbated by taking ergotamine preparations. Her health was otherwise good.

On physical examination she was fit but anxious. The upper limb pulses were present and equal. There were no ischaemic changes in the fingers and the blood pressure was 120/80 mm.Hg in both arms. There were no pulses palpable below the femorals in either of the lower limbs and there were no bruits. The feet and legs were cold and pale. The capillary return was slow and there was reduced sensation to pin-prick, touch, and vibration over the soles. All the reflexes were present and equal.

She was treated with bed rest. The limbs were kept at room temperature. An intravenous infusion of low molecular dextran (Rheomacrodex), 500 ml. eight-hourly, containing heparin 5,000 i.u. was begun and a total of 2 litres was given. The legs became warmer but she experienced severe burning pain, requiring large doses of analgesics. The cutaneous sensitivity improved and the toes showed reactive hyperaemia. The foot pulses returned on the fifth day. She was discharged on the nineteenth day, when the foot pulses were present and the capillary return was normal, but she still experienced some hyperaesthesia of the soles, aggravated by walking. One month later these symptoms persisted but had diminished.

INVESTIGATIONS

Arteriography.—Bilateral femoral arteriograms were performed on admission and showed severe narrowing of both superficial

femoral arteries which extended throughout their lengths. The calf vessels were very narrow and difficult to identify. Smoothness of the arterial wall was seen at all levels (Fig. 1, right thigh). On the tenth day bilateral femoral arteriograms were repeated and showed good filling of the femoral and popliteal arteries in both limbs (Fig. 2, right thigh). No atheroma or occlusion was demonstrated.

FIG. 1.

FIG. 2.

Plethysmography.—Serial studies were made of blood flow in the calf and foot of both legs by strain-gauge plethysmography. Measurements were made of resting flow and perfusion pressure (the difference between arterial systolic and venous pressure). The peripheral resistance was calculated from the ratio of perfusion pressure to the resting flow.

Ultrasound.—Daily examination of the posterior tibial and dorsalis pedis pulses and indirect ankle systolic pressure measurements were made by means of a Doppler flow velocity meter (Yao *et al.*, 1968, 1969) until the pulses were clinically palpable. Improvement was seen in the change of configuration of the flow velocity patterns, together with gradually increasing ankle systolic pressure.

Thermography.—Serial thermographic pictures of both feet were made with a thermogram (AGA Thermovision) (Fig. 3). Cold areas appear black and hot areas white. Intense hyperaemic flow was seen in the left foot.

FIG. 3.—Thermogram of feet, showing intense hyperaemic flow of left foot.

COMMENT

Peripheral arterial insufficiency due to ergot poisoning, though uncommon, has been reported previously. Glazer *et al.* (1966) and Iuel and Mune (1967) demonstrated the sequential changes of blood flow when studied by plethysmography. These changes were well shown in this patient.

Renal Homotransplant Rejection Associated with Microangiopathic Haemolytic Anaemia

British Medical Journal, 1970, 3, 87-88

The association between disease of the small blood vessels and red cell fragmentation, termed microangiopathic haemolytic anaemia, is well recognized. This report describes a patient in whom this disorder developed after renal homotransplantation.

CASE REPORT

In 1966 a 30-year-old man was admitted to hospital with a minor cerebrovascular accident, malignant hypertension, and chronic renal failure. This was controlled with methyl dopa and Giovanetti diet until 1967, when regular haemodialysis therapy was started. During this period the haemoglobin concentration was maintained at about 6 g./100 ml. by blood transfusion. The blood film showed only anisocytosis and the platelet count was 120,000/cu. mm.

In April 1968 bilateral nephrectomy was carried out and antihypertensive therapy was stopped. Renal homotransplantation with a cadaver kidney was performed on 7 August. The warm ischaemic time for the kidney was 20 minutes. Tissue leucocyte typing showed the donor and recipient to differ in one major antigen. At this time the haemoglobin was 8.2 g./100 ml. and the platelet count 150,000/cu. mm. (see Chart). Postoperative renal function

The narrowing of the superficial femoral arteries extended throughout both thighs, and these appearances had reverted to normal in the arteriograms repeated 10 days later. The plethysmographic and thermographic studies showed the presence of hyperaemic flow, which is commonly seen after the return of foot pulses and is experienced as a burning sensation (St. Anthony's fire). Plethysmographic studies also showed that vasoconstriction in the distal vessel bed may persist for some days after the disappearance of major vessel spasm. The flow velocity pattern and pressure studies by the ultrasound technique firmly established its value in monitoring the progress of arterial disease, especially when the distal pulses are not palpable at the outset (Yao *et al.*, 1968, 1969).

The diagnosis of ergot poisoning is easily made when a clear history of administration is obtained. This should be sought in all young patients presenting with arterial insufficiency in the limbs. In the present case the recovery was rather slow, but this was complete after withdrawal of the drug.

We would like to thank Dr. T. Csato for referring the patient; Professor W. T. Irvine for permission to use the blood flow laboratory of the surgical unit; Dr. D. Sutton for performing the arteriograms; and Dr. C. Gourmos and Mr. T. N. Needham for their assistance.

S. T. YAO, M.D.,

Lecturer in Surgery.

D. P. GOODWIN, M.B., B.Sc., F.R.C.S.,

Senior Surgical Registrar.

J. R. KENYON, B.S.C., CH.M., F.R.C.S.,

Consultant Surgeon.
St. Mary's Hospital, London W.2.

REFERENCES

- Glazer, G., Myers, K. A., and Davies, E. R. (1966). *Postgraduate Medical Journal*, 42, 562.
Iuel, J., and Mune, O. (1967). *Vascular Diseases*, 4, 159.
Yao, S. T., Hobbs, J. T., and Irvine, W. T. (1968). *British Medical Journal*, 4, 555.
Yao, S. T., Hobbs, J. T., and Irvine, W. T. (1969). *British Journal of Surgery*, 56, 676.

Haematological results and therapy after homotransplantation.