to light through the work of the team at the Imperial Cancer Research laboratories¹ and others.

It is probable that the enormous social convenience of the pill—and the first true emancipation of women that has occurred—are clouding our judgement to the true risks. It is striking that the possible induction of carcinoma is barely mentioned in the countless discussions on the pill that can be read in the non-medical press. Have the sociologists (and the drug houses) won the argument yet again before the problem has even been properly formulated, let alone solved?—I am, etc.,

PAUL STRICKLAND.

Regional Radiotherapy Centre,
Mount Vernon Hospital,
Northwood, Middx.

REFERENCE

- ¹ Imperial Cancer Research Fund, *Annual Reports*, London, 1964/69.

Diagnosis of Pulmonary Embolism

SIR,—I was interested in the Clinical Progress article (27 June, p. 773) on the diagnosis of pulmonary embolism. I refer particularly to the final paragraph on the mainstays of diagnosis “in an acutely ill patient.”

I have recently been rather worried by some six cases of acute pulmonary embolism in young, fit patients, four of whom have walked down to the surgery complaining only of a little pain in the chest. Only one out of the six showed any tachypnoea. All were of sufficient severity to be treated with anticoagulants. None of them had leg or abdominal signs.—I am, etc.,

S. B. TWIVY.

Dunstable, Beds.

House-dust and Asthma

SIR,—The connexion between house-dust and asthma (30 May, p. 501) was noted as long ago as 1698 when Sir John Floyer referred to it in his “Treatise on Asthma.” During the past six years evidence has accrued that the major antigenic component of house-dust is derived from the mite *Dermatophagoides pteronyssinus*,¹ whose main source of food is shed human epidermal scales. In this connexion bed-making is notorious for provoking allergic respiratory symptoms (sneezing, asthma). Maunsell *et al.*² found that the mite content of house-dust varied according to its size of collection, being greatest in dust from mattresses.

It is interesting, therefore, to see what observations Ramazzini, the father of occupational medicine, made on the effect on the lungs of dust from old mattresses and clothes. Bernardino Ramazzini, born in 1633, was professor of medicine at Modena from 1671 to 1700, and thereafter held the chair at Venice. In 1700 the first edition of his *De Morbis Artificum* (*Disease of*

Workers) was published. The following extract is taken from the 1713 edition, the translation from the Latin being slightly modified from that of Wright.³

“It is the custom, in Italy at least, for Jews to remake wool mattresses in which the wool has become packed by several years of daily use on beds, so that they are too hard. They lay the wool on hurdles of wicker-work, beat it with sticks, and shake it thoroughly; thus they make the mattresses softer and more comfortable to lie on. . . . In shaking and combing it they take in by the mouth a great deal of foul dust. This causes serious ailments, a harassing cough, dyspnoea, and a disordered stomach. . . . I consider that it is not this dust from old wool that is so deadly as that which rises from the impurities left in it by the bodies of those who have lain on the mattresses. . . .”—I am, etc.,

BERNARD J. FREEDMAN.

Dulwich Hospital,
London S.E.22.

REFERENCES

- ¹ Voorhorst, R., Spijksma-Boezen, M. I., and Spijksma, F. T., *Allergie und Asthma*, 1964, 10, 329.
² Maunsell, K., Wraith, D. G., and Cunningham, A. M., *Lancet*, 1968, 1, 1267.
³ Ramazzini, B., *De Morbis Artificum*, 1713, translated by Wright, W. C., Chicago, University of Chicago Press, 1940.

Who Should Do Psychotherapy?

SIR,—In asking this, and other pertinent questions, your leader writer (13 June, p. 617) refers to the survey of the outcome of referring patients to psychiatric units from my practice over a five-year period,¹ and in doing so appears, probably inadvertently, to be using what I described as the “somewhat disappointing” results as evidence to support the suggestion that psychotherapy is not of any great value.

I am pleased to be recognized as “a by no means unfriendly critic”, but would be grateful for a little space in which to emphasize that the results of my survey did nothing to support Eysenck’s view that psychotherapy is of no value—but they do show up very clearly the appalling shortage of staff attached to the psychiatric units and the shamefully poor administration of this sector of the National Health Service. That many of my patients did so badly was because of the disgracefully long wait for their initial consultation, and the further six to nine months delay before they could expect to be put in the psychotherapeutic group recommended for them. This is not a criticism of my consultant psychiatric colleagues, but of the conditions under which they have to work.

As your leader-writer states, there is a growing volume of published evidence on the usefulness and value of properly organized psychotherapy in a variety of different groups of patients, so that Eysenck’s “bomb” is turning out to be little more than a damp squib. Indeed, as pointed out in the paper on psychiatric referrals,¹ the results even of the superficial psychotherapy that can be expected from a family doctor are far better in many patients than can be expected from traditional medical treatment, including all the most up-to-date tranquillizers. Follow-up of

several of those patients whose results were “somewhat disappointing” showed that they benefited from properly given psychotherapy either by myself, or by psychotherapists consulted privately. This leads on to the suggestion, quoted by you and favoured in the U.S.A., that this type of treatment should be given by psychologists and trained counsellors. This could well be the answer for many patients.

However, it is still my view that the majority of patients requiring treatment by one of the several psychotherapeutic methods would best be treated by their own family doctors, if only proper training could be given to, and the necessary skills learned by, these doctors. They have a feeling of inability to deal with a large proportion of the suffering they meet in their patients, so that that terrible term “trivia” is applied to much of their patients’ illnesses, and where once a doctor would have talked about his enjoying a busy practice he now talks about his heavy “work-load”.

Balint’s work on this subject has done much for many of us, as is now shown by the development in Belgium, France, and Switzerland—and recently in Great Britain—of medical societies of Balint groups, whose primary aims include the encouragement of further research into, as well as providing a forum for further discussion of, all those problems associated with the patient-doctor relationship. In this way perhaps we may find ways and means of improving the quality of our medical care for the large majority of our patients who require some form of psychotherapy as part of their treatment.

There is no doubt in my mind, after applying these ideas in my practice for nearly 20 years, that large numbers of patients with psychosomatic disorders, as well as those with frank psychoneurotic problems who ordinarily would not dream of consulting a psychiatrist or psychotherapist, can and do benefit enormously from simple or minor psychotherapy that is well within the competence of the family doctor. This not only helps the patient, but also makes the doctor’s work more satisfying. It also reduces considerably the need for constantly having to refer patients to hospital out-patient clinics for innumerable and unnecessary investigations into their illnesses which are based not on pathological changes but on emotional disorders.

I am convinced that if we could improve our standard of medical practice in this way we would save the Health Service millions of pounds each year by avoiding so much unnecessary hospital investigation—and at the same time improve the status of the family doctor in the eyes of the public.—I am, etc.,

PHILIP HOPKINS.

London N.W.3.

REFERENCE

- ¹ Hopkins, P., and Cooper, B., *British Journal of Psychiatry*, 1969, 115, 1163.

Gold Therapy in Rheumatoid Arthritis

SIR,—Reading the medical memorandum about the nephrotic syndrome and gold therapy (27 June, p. 772) I notice that the dosage of gold aurothiomalate given to the patient