of the findings remain to be published.

I have never been an enthusiastic advocate of immunofluorescence in the general diagnosis of virus disease,¹ but it may encourage those who have struggled to identify virus antigen in smears of inflamed tissue or deposits of degenerating cells to know that the identification and titration of specific antibody by straining a known virus antigen is an easier and a much more exact technique. The identification of IgM and other components is a new and significant factor.

The man without means to buy a high-speed centrifuge or one without skill in gradient-centrifugation should try fluorescent antibody. I predict that a clean preparation of rubella-infected cells, specific antisera to human immunoglobulins from Wellcome Reagents Ltd., an inexpensive, but efficient Reichert Fluorpan microscope, and patients' sera will provide a rapid, reliable, and beautiful test which will give the ultra-centrifuge a run for its money now, and will replace it quite soon.—I am, etc.,

K. B. FRASER.

Department of Microbiology,
The Queen's University of Belfast.

REFERENCE

- ¹ Fraser, K. B., in *Fluorescent Protein Tracing*, Ed. R. C. Nairn, 3rd edn. Edinburgh, Livingstone, 1969.

Infantile Gastroenteritis

SIR,—Your leading article on this subject (4 July, p. 2) leaves a lot to be desired. "The cure of the illness lies, therefore, in the prevention or treatment of dehydration—usually a simple matter." This is an oversimplification of the situation, and begs the main question about the prevention of the disease itself. One imagines that your remarks are mainly aimed at the doctor practising in the community, because the treatment and management of the more severe cases admitted to hospital is always rather more than a "simple matter," and I for one would not wish to follow your advice of avoiding antibiotics. Some of us can remember the treatment of gastroenteritis before suitable antibiotics were available and remember the main problem then was the relapsing nature of the disease. This problem was almost abolished by the use of oral non-absorbed antibiotics such as streptomycin and later neomycin. I prophesy that if your advice is followed about the non-use of antibiotics, we shall see this situation again in the hospital wards. Many of us think that not all the pathogenic strains of *E. coli* and other bacteria which can cause infantile gastroenteritis have yet been identified, and many of us think that not all laboratories are equally skilled in identifying the strains already known. Reliance upon the result of faeces culture etc. for starting antibiotic treatment may therefore be misleading from the point of view of identification of the causative organisms and dangerous from the point of view of delay in a seriously ill infant. The main use of faeces culture is to monitor the sensitivity of the organisms to antibiotics, and here again some of us would feel that the *in vivo* sensitivity of the organism is a different matter from the *in vitro* sensitivity reported by the laboratory, which in any case is very rarely

given in a quantitative manner so as to be really meaningful in the clinical situation.

In the case of the carrier stage of food poisoning organisms I think there is need for strong stricture against the misuse of antibiotics—especially one after the other through the long (two to three months) carrier stage—but the advice in your leading articles of discouraging the use of antibiotics in acutely ill infants with gastroenteritis, and a few weeks ago with whooping cough (13 June, p. 619) is, I think, most unfortunate. One can still remember the number of children with bronchiectasis which followed complicated whooping cough and measles in the days before antibiotics were used.

I am delighted to see that Dr. T. S. Matthews (18 July, p. 161) has corrected your failure to mention breast feeding as the most important means of preventing infantile gastroenteritis altogether. Short of the ideal of universal breast feeding of babies, I have been using a method of prevention of infantile gastroenteritis in hospital over the past seven years which seems to be effective. This method involves the administration of a culture of *Lactobacillus acidophilus* (Uniflor) $\frac{1}{2}$ to 1 teaspoonful of powder in alternate feeds to all babies in such wards. The preparation Enpac, in which there is the same organism made resistant to all the main antibiotics, is used if an infant is receiving antibiotic treatment. I also insist on a policy of non-admission to a general ward of any infant with diarrhoea, and immediate isolation of any child or infant who develops loose stools. Infantile gastroenteritis is one of the most infectious of all diseases, and we can ill afford to neglect any means of preventing it.—I am, etc.,

ROBERT WIGGLESWORTH.

Kettering, Northants.

SIR,—Your statement (4 July, p. 2) that antibiotics should not be prescribed in the routine treatment of infantile gastroenteritis cannot pass unchallenged. There have been few major controlled trials in the use of antibiotics in this disease in this country. As the pattern of the disease has not changed in the intervening years, the results of such a controlled trial carried out at the Queen Elizabeth Hospital for Children in 1952 are still relevant.¹

One hundred and nineteen patients with infantile gastroenteritis, under the age of one, were divided into three treatment groups and given three identical looking preparations labelled A, B, and C. These were dispensed in strict rotation in a blind trial. Mixture A contained an inert substance. Mixture B contained chloramphenicol, prepared to supply 75 mg./lb. of body weight. Mixture C contained sulphadiazine prepared to supply 125 mg./lb. of body weight, in the amount to be given daily. The patients were further subdivided into two severity groups, mild and severe, depending on whether or not intravenous fluids were required within the first 24 hours of admission. Special type *E. coli* were isolated from 40 of the 119 patients. Supportive treatment was standardized as far as possible. Comparisons of progress were made and the following criteria used.

Patients fully recovered by the tenth day were regarded as having made straightforward progress. All others, including patients who did not tolerate the standard regimen of regraded feeding or whose condition deteriorated during treatment were classified as making delayed progress. Patients whose clinical condition became extremely grave were taken out of the trial. These "failures" were then given whatever treatment or antibiotic which was considered to be most effective. Thus the basis of comparison between trial and control groups was the relative number making straightforward or delayed progress and the relative number of "failures".

The results of the trial were:

(1) Of the 42 patients treated with chloramphenicol 40 made straightforward progress to recovery in an average of 8.2 days.

(2) Of the 39 patients treated with sulphadiazine 29 made straightforward progress to recovery in an average of 8.5 days.

(3) Of the 38 control cases 22 made straightforward progress to recovery in an average of 12.1 days.

The two patients who failed to respond to chloramphenicol were severe cases, which persistently excreted *E. coli* 055 resistant to chloramphenicol but sensitive to chlortetracycline.

These results showed that antibiotics have a definite place in the treatment of *E. coli* gastroenteritis. While it is not disputed that the majority of patients with gastroenteritis will make straightforward progress to recovery with supportive treatment alone, I would urge that an effective antibiotic should be administered under the following circumstances:

(1) Patients failing to make straightforward progress in spite of adequate rehydration and supportive therapy.

(2) Bacteriological results showing the presence of a type-specific *E. coli* sensitive to a specific antibiotic, then this antibiotic should be given.

(3) In epidemic outbreaks in nurseries or children's units especially among infants under the age of one, type-specific *E. coli* must be suspected as being the aetiological factor. It has been shown frequently in the past that there are many additional pathogenic strains of *E. coli* which are not covered by the standard antisera and which may not, as yet, be identified. Every effort must, therefore, be made with the assistance of the bacteriologists to see if such an organism can be implicated. Treatment with antibiotics must be given in the meantime without necessarily waiting for the bacteriological results.—I am, etc.,

L. HAAS.

City Hospital,
Exeter.

REFERENCE

- ¹ Haas, L., M.D. Thesis, Cambridge University, 1953.

Surgery of Fallopian Tubes

SIR,—I ventured to suggest (21 March, p. 761) that the surgery of the Fallopian tubes was far behind the modern standards of surgery, and that a transplant should be developed for certain cases of atresia. This has now been started at Monash University,