Ronald Bodley Scott for an expert haematologist's opinion. His reports to me confirmed the frequent incidence of thrombosis occurring in those cases of prostatic carcinoma undergoing hormonal therapy.

Later I became alerted by the thrombocytic tendency in the female towards the higher oestrogen content in the contraceptive pill. Now having lost a patient, aged 60 years, not due to his low-grade prostatic malignancy, but to extensive arterial thrombosis following months of intensive hormonal therapy, I have decided to continue future treatment under anticoagulant treatment subjected to adequate control.

Relatively speaking, prostatic carcinoma is a slow progressive disease afflicting usually the aged. The mortality rate is slow compared with cardiovascular disease, to which most succumb in the end. So rather than precipitate the end why not incorporate a therapeutic measure to combat both?—I am, etc.,

E. SUTHERLAND-RAWLINGS.

London W.2.

REFERENCE

- ¹ *British Medical Journal*, 1968, 2, 132.

Profile Analysis

SIR,—In spite of the fact that Professor T. P. Whitehead is an old friend for whose opinion I usually have the greatest respect, I feel it incumbent upon me to contradict a statement attributed to him in your leading article (22 August, p. 417).

To say that "clinical chemistry is not medicine, it is a service to medicine," is all very well from the point of view of Professor Whitehead, who is not medically qualified. It is, however, my considered opinion that the subject in the hands of the medically qualified practitioner is undoubtedly an important branch of medicine, in the same sense as any other branch of clinical pathology. To say it is not medicine would mean that it would no longer attract medical graduates, and this would be a catastrophe of the highest order.

You point out in your leading article that Professor Whitehead and his colleagues in Birmingham have recently reported that profile analysis carried out on patients in general practice led to new diagnoses in 16.9% of patients. To quote from their article (7 March, p. 620) "iron-deficiency anaemia was the major finding . . . Diabetes, renal disease, and one patient with myxoedema accounted for the remaining diagnoses." To diagnose iron-deficiency anaemia from a serum iron and haemoglobin level alone is perhaps a little unwise, and I am surprised that in apparently only one of these cases was an obvious cause for the anaemia found, but in any case, in the list of altered diagnoses given, most of the profile tests were hardly necessary.

I fully approve of the Birmingham work from a research point of view, and if the general practice which is reported is representative of general practice as a whole then one can only conclude that the large battery of biochemical investigations was not necessary. In fairness to the Birmingham workers they do point out that "The choice of tests should remain flexible

in the light of further experience. . . ."

A final thought occurs to me. In what percentage of patients not showing abnormal chemical profiles would an altered diagnosis have been made if all of these had been referred to hospital?—I am, etc.,

A. L. LATNER.

Department of Clinical
Biochemistry,
University of Newcastle upon Tyne.

E.S.R. Surveys

SIR,—Dr. H. Dale Beckett (15 August, p. 408) asks can E.S.R. surveys replace mass miniature radiography surveys when the latter are discontinued? The answer from experience is a qualified "yes."

In about 1940 E.S.R. surveys were used successfully at West Park Hospital, Epsom, to identify patients likely to be suffering from tuberculosis. Shortly after being appointed to the Mental Hospitals Group Laboratory at Epsom, I made E.S.R. surveys available to hospitals in the L.C.C. Mental Health service, Cane Hill Hospital being one of them. These surveys continued till mass miniature radiography became available to the hospitals.

E.S.R. surveys are relatively inexpensive, and with adequate skill, equipment, and organization can be carried out quickly. An average of 100 venepunctures an hour is easily attained (our best was 120 in 54 minutes). The disadvantage is the disadvantage of the E.S.R.—lack of specificity. About 20% of abnormal results—or even more—can be expected, and all patients with high rates need clinical physical examination and careful selection for full x-ray examination. Nevertheless, E.S.R. surveys can be made to work and be clinically valuable.—I am, etc.,

W. W. KAY.

Greenfield,
Bedford.

Thromboembolic Disease and the Pill

SIR,—In a recent paper by Dr. W. H. W. Inman and his colleagues (25 April, p. 203) an analysis is shown of 1,305 reports of thromboembolism occurring during the time period 1 January 1965 and 30 June 1969 in women using oral contraceptives. The conclusion made was "the data . . . leave no doubt that there is a positive correlation between the risk of thromboembolism and the dose of oestrogen in oral contraceptives."

It is the purpose of this letter to point out reasons why this relationship cannot be proved by the data and methods used by the authors.

The study group was made up of cases of thromboembolism obtained from voluntary reports from the doctors to the Committee on Safety of Drugs. As estimated by the authors this sample constituted less than 10% of the total number of cases that occurred during the time period. A 10% sample would be more than adequate if it is either random or representative of the whole. This sample is obviously not random and questionably representative. Voluntary reports are known to be subjected to biases of all sorts. In fact, the sample appears to be biased in the direction of new drugs, since

the Committee instruct the doctors to report all the reactions to new products and only the serious reactions to the established ones.

No patient control data were used in this study. Instead, the expected number of occurrences of thromboembolism were determined from market research estimates of sales, under the assumption of equal risks for all the products. This departs from the acceptable procedure used in retrospective studies where, in addition to using patient control data, attempts are made to match them to the study group by characteristics such as predisposing conditions, age, parity, etc.¹⁻³ Additional uncertainties and biases arise from the use of market research estimate of sales as substitute for control data: patient change from product to product, time lag between sales and occurrence of a disease, etc.

The significance of the relation between thromboembolism and oestrogen dose levels was proved using a χ^2 test with one degree of freedom. This test results from a modification of the test of significance of a slope under the assumptions that sampling errors in the expected frequencies are relatively negligible and that the total number of thromboembolism is small when compared with the total sales.⁴ Normally when the expected frequencies are not known the χ^2 test can still be applied with a reduced number of degrees of freedom. This obviously does not apply here with only one degree of freedom available. It is not clear what the effect of this uncertainty would be in the results obtained.

Four oestrogen dose levels: 50 μ g, 75/80 μ g, 100 μ g, and 150 μ g were used in testing significant trends. The last dose level, represented by one out of the 15 brands used in the study, had a strong effect in the result. It is fair to point out that if this dose level is eliminated no significant trend exists. For instance, when all the groups are combined, the ratio of observed/expected for the dose levels of 50 μ g, 75/80 μ g, and 100 μ g are equal to 1.07, 0.78, 0.98, respectively, which are not significantly different. These values were obtained after adjustments were made in order to have the total and expected frequencies equal to each other.

The authors also show that no significant differences exist between mestranol 100 μ g and ethinylestradiol 100 μ g ($\chi^2=2.7$) and between mestranol 50 μ g and ethinylestradiol 50 μ g ($\chi^2=2.5$). It is interesting to note that following the same procedure it can be shown that mestranol 150 μ g does not differ from ethinylestradiol 100 μ g ($\chi^2=0.33$) and that mestranol 75/80 μ g does not differ from ethinylestradiol 50 μ g ($\chi^2=0.89$). These latter results appear to be related to estimates available of the relative potencies of the two oestrogens.

The discrepancies shown above are a direct result of the lack of patient control data. To prove the possible existence of a relationship between thromboembolism and oestrogen dose levels, a study should be based on a large, unbiased sample of thromboembolic cases and on carefully matched patient control data.—I am, etc.,

LUIS F. NANNI.

Rutgers University,
New Brunswick, New Jersey, U.S.A.

REFERENCES

- ¹ Sartwell, P. E., Masi, A. T., Arthes, F. G., Greene, G. R., and Smith, H. E., *American Journal of Epidemiology*, 1969, 90, 356.
- ² Inman, W. H. W., and Vessey, M. P., *British Medical Journal*, 1968, 2, 193.
- ³ Vessey, M. P., and Doll, R., *British Medical Journal*, 1969, 2, 651.
- ⁴ Armitage, P., *Biometrics*, 1955, 11, 375.