entry to the medical club so technically rigid that valuable new members are excluded—drugs which could have enormous potential in helping with the geriatric crisis.—I am, etc.,

C. G. ELLIOTT

East Hoathly,
Sussex

Treatment of Myasthenia

SIR,—May I add my support to Dr. S. Wolf's letter (19 June, p. 710).

For many years it has been the practice in this unit^{1,2} to stop all cholinesterase inhibitor drugs following thymectomy until such time as there was clinical evidence of the need for these drugs. Then, and only then, were the drugs restarted, and invariably in a much reduced dosage.

Since the adoption of this procedure the incidence of cholinergic crisis has been reduced to negligible proportions.—I am, etc.,

M. J. LANGE

Endocrine Department,
New End Hospital,
London N.W.3.

¹ Lange, M. J., *British Journal of Surgery*, 1960, **48**, 285.

² Lange, M. J., *Proceedings of the Royal Society of Medicine*, 1968, **61**, 751.

Unnecessary X-rays

SIR,—Mr. J. Charnley's reply (1 May, p. 279) to Mr. D. F. Thomas (10 April, p. 105) is too pessimistic. He rightly stresses the value of two separate clinical examinations as a means of avoiding unnecessary x-rays but he assumes that these must be made by the same doctor. This would be true if clinical records were not kept.

If there is no convincing clinical evidence of a fracture in a limb then the patient should be told to come back in three days for re-examination and x-ray if necessary. If the need to avoid unnecessary x-rays is explained to the patient and the proposed procedure is noted on the case card, then even if a fracture is subsequently discovered, the casualty officer can be assured on three points: (1) The effect of delay in treatment of the fracture will not be detrimental to the final result. (2) The doctor will not lose face with the patient nor will he be liable for damages for negligence. (3) The doctor will not lose face if his colleague diagnoses a fracture in three days' time.

In the previous correspondence on this subject last year (7 February 1970, p. 362) I advocated a three-day delay in radiography of doubtful fractures of limb bones. This provoked a rejoinder from the Medical Protection Society warning us that patients might even be able to bring a claim based on three days' delay in the treatment of a minor fracture (28 February 1970, p. 564).

In the recent correspondence Mr. Thomas rightly stressed that the dangers of not taking an x-ray must be balanced against the possible harmful effects of taking too many x-rays. Once again this provoked a warning, this time from the Medical Defence Union, to the effect that failure to take an x-ray could be regarded as ipso facto negligent (1 May, p. 278).

It is therefore becoming apparent that the main obstacle to a rational policy to avoid unnecessary x-rays is not the legal profession but our own profession. The legal profession

is entirely dependent upon medical evidence to establish negligence. Therefore unless the medical advisers to the protection societies and the medical witnesses that are called for the plaintiff are convinced of the need for balanced judgement in ordering x-rays then there is little hope of improvement in the present position.

Both groups of correspondents included a neurosurgeon warning about the pitfalls in the diagnosis of skull fractures. For this reason it is vital that clear instructions should be issued to casualty officers on the principles that should guide their discretionary policies in ordering x-rays, as I suggested in my earlier letter.—I am, etc.,

A. W. FOWLER

Bridgend General Hospital,
Bridgend

Thalidomide

SIR,—In a letter on the subject of thalidomide (19 October 1968, p. 190) I wrote "After the lapse of time it may be reasonable to inquire if there has been any survey of the type of mothers who required or demanded the exhibition of thalidomide in the first three months of pregnancy. Here, surely, there is some abnormality."

Apart from inquiries for reprints from America, France, and Australia no interest was shown.

However, a qualified medical social worker in the North East of England in the local newspaper of 8 June is reported very fully on this subject in an interview headed "Thalidomide: A Cry for Help."

She refers to a high proportion of her cases being single-parent families, many of whom are divorced or separated. In many instances this is caused by stress resulting from the thalidomide tragedy. These families have very low incomes or live on Social Security. "There is a tremendous feeling of guilt felt by many parents for having taken thalidomide and a tremendous amount of social work help is as a result needed by those families."

The article refers to injustice, or considered injustice, by the Social Services towards parents, the fear of antagonism between afflicted children and parents, the need of money, the fight for financial compensation not forthcoming without expensive legal costs, and the legal safeguard that the children's money must be held in trust until they are 18.—I am, etc.,

RICHARD BELL

Haydon Bridge,
Hexham

¹ *Newcastle Evening Chronicle*, 8 June 1971.

The Forgotten—An Occupational Centre

SIR,—The occupational centre described in the article "The Forgotten: VIII—An Occupational Centre" (1 May, p. 272) can provide a most valuable social setting and means of encouragement and support for the severely disabled. But such centres should be regarded as a means to an end and not necessarily an end in themselves.

It is important that disabled people attending these centres are re-assessed regularly by a suitably qualified person so that each individual undertakes activities

suitable for his changing physical and mental abilities. If his progress so indicates he should be able to transfer to another type of centre—possibly a medical rehabilitation centre, an industrial rehabilitation unit, a government training centre, or a sheltered work shop—eventually perhaps even to open employment.

This is what rehabilitation is all about. But progress is sometimes impeded because these different types of centres are the responsibility of different government departments—for example Health and Employment, or local authorities, or indeed voluntary organizations. A greater degree of co-ordination is much to be desired.

Occupational therapists are employed by the Department of Health in the rehabilitation of patients in hospital and rehabilitation centres and by the local authorities to advise on aids and adaptations to the home and to supervise occupation day centres. Their work brings them into close contact with medical practitioners, social workers, and the Department of Employment in the person of the disablement resettlement officer.

I would respectfully suggest that greater use is made of the skills of the occupational therapist who is in a position to liaise and is trained to assess the physical and mental capacity of the disabled in relation to the requirements of personal independence, social activities, and work situations.—I am, etc.,

ELIZABETH GROVE

Chairman,
Association of Occupational Therapists
Wolfson Medical Rehabilitation Centre,
Atkinson Morley's Hospital,
London S.W.20

Migraine Research

SIR,—I read the articles on migraine (26 June, pp. 751-756) with interest. I was surprised to find no reference to the possible aetiological influence of errors of refraction and ocular muscle balance. I took particular interest in this subject years ago, largely on account of the views held by my then colleague at Guy's, Sir Arthur Hurst. I quote one or two sentences from his Savill lecture of 1924: "Very small errors (of refraction) are quite sufficient to give rise to migraine: the most accurate possible correction of every error is required: muscle balance defect may by itself cause migraine; the most important part of treatment is the prevention of eyestrain." His enthusiasm was such that he arranged lectures, jointly delivered by himself and myself, to propagate the thesis.

While my conviction by no means equalled his, it was natural that I should take special interest in the subject. One result of this was an analysis of 100 consecutive private patients who had either been referred to me for migraine or whose trouble was diagnosed as such on arrival. Of this series 14 were rejected as not true cases of migraine; unfortunately no report was obtained for 70; seven were unimproved, but nine of the 86 were improved or completely cured by the correction of errors of refraction or of muscle balance. Even this number, which is certainly inaccurately small, justifies careful examination and accurate correction by an ophthalmologist in every case.

A further omission in these articles which

I noted was mention of treatment by urea. I have had a few quite dramatic successes with this drug in otherwise resistant cases, and feel that it should remain on the list of possible remedies.—I am, etc.,

FRANK W. LAW

London W.1

¹ Hurst, A. F., *Lancet*, 1924, 2, 1.

² Law, F. W., *Transactions of the Ophthalmological Society of the United Kingdom*, 1956, 76, 153.

SIR,—I found myself saddened to read Dr. J. M. Blau's comments in relation to psychotherapy in his article on "Migraine Research" (26 June, p. 751).

I do not intend to go into any counter arguments about the importance of psychological factors in migraine, but merely to comment on his reasons for minimizing them. Surely Dr. Blau does not believe that people with psychological problems can be recognized on sight. He must know that someone with psychological problems is not necessarily mad. Many neurotic people attain high professional status and function very well in their work. The fact that pleasant occasions can precipitate migraine in some people only supports the importance of psychological factors. These people may have great difficulty in being happy. The fact that migraine occurs after prolonged sleep is an argument in favour of psychopathology rather than against it, because the mind is not asleep in sleep.

In the frenzied search for organic causes in migraine and indeed in other illnesses, let us not lose sight of the importance of psychological factors, even if these are more difficult for us to comprehend and to treat.—I am, etc.

RAYMOND GLEDHILL

London W.1

Medical Presentations of Choriocarcinoma

SIR,—I really enjoyed reading the article "Medical Presentations of Choriocarcinoma" (12 June, p. 633). I think I have learned a lesson from this article that a pregnancy test of urinary gonadotrophin assay should become a routine in postnatal clinics as well as in abortion and terminations follow-up clinics. The intensive follow-up of diagnosed cases of trophoblastic tumours is now well known to all gynaecologists.—I am, etc.,

N. G. MUSSALLI

The Manor Hospital,
Walsall, Staffs

Hydatidiform Mole and Hyperthyroidism

SIR,—I was very interested in your leading article (12 June, p. 606) on the relationship between hydatidiform mole and hyperthyroidism and would like to quote a case of my own.

A 38-year-old Kikuyu woman presented at hospital outpatients with florid thyrotoxicosis and in cardiac failure. On admission she was found to have a mass in the lower abdomen equivalent to a 20-week pregnancy and a history of three months' amenorrhoea, but with intermittent small losses of blood per vaginam.

The human chorionic gonadotrophin titre was raised and a diagnosis of hydatidiform mole was made. Once her cardiac state had been stabilized an evacuation was carried out. Almost immediately her cardiac state began to improve and the thyrotoxicosis disappeared.

I followed her up for 18 months before leaving Kenya, and she had no recurrence of thyrotoxic symptoms or cardiac failure, and there was obviously a direct causal relationship between the two conditions.—I am, etc.,

A. ROBSON

Selby, Yorks

Encouraging Recruits to Medicine

SIR,—Your columns have recently contained a fascinating set of letters dealing with medical students. A main focus of interest has been medical students' grants. I should like to assure Mr. A. R. W. Forrest (1 May, p. 281) that although medical students may not have "the backing of a large and experienced union" or "public sympathy" the British Medical Students' Association has been waging a most vigorous campaign over the last 18 months on his and other medical students' behalf. Surveys were undertaken, the support of deans and vice-chancellors was enrolled, and M.P.s were lobbied intensively leading to a question in the House. I hope that this positive action will encourage Mr. R. D. Clegg (17 April, p. 168) not to fear excessively for the future of the medical profession on the basis of medical students' "self-pity".

Even more important, I trust that Dr. D. J. MacMyn for his part will not dismiss the B.M.S.A. as "smart-Alec, cynical and materialistic" (1 May, p. 281). We feel as much as Mr. J. A. Lourie (15 May, p. 402) "that to be a student of medicine is a very great privilege" and here one passes to the second main focus of your correspondents' concern—recruits to medicine. The two subjects are of course not easy to separate.

Mr. A. D. Shroot (5 June, p. 591) questioned Dr. John McKee's inclination "to accept all doctors' offspring with reasonable standards, as at least they should have a true knowledge of the type of life ahead." (15 May, p. 397). It cannot be denied that there are advantages in favourably considering applications from members of medical families, but clearly the prime duty of selection committees is to ensure that places are offered to candidates possessing that hard-to-define mixture of qualities which goes to make the "good doctor." One hopes that the medical profession no longer bars anyone displaying the necessary academic ability, common sense, dedication, kindness, and sense of humour; what is tragic is that official grants are not available at all for those graduates who may be eminently suited to becoming superb doctors, even when they have gained places at medical schools. Among these people is to be found a supreme sense of vocation that would undoubtedly warm Mr. Clegg's heart and that "humanity and interest in patients as people" the apparent absence of which so rightly concerns Dr. McKee.

A glance at the letters in a B.M.S.A. file in my possession describing various graduates' financial struggles might perhaps convince Dr. MacMyn of the need for all medical students to fight for enough money for them to enjoy the great privilege of

becoming doctors without the one obsessive thought of "Who is going to pay for my medical education?"—I am, etc.,

RICHARD WESTCOTT
Vice-President B.M.S.A.

Oxford

Radiological Training

SIR,—The article by Dr. A. G. W. Whitfield and others (29 May, p. 514) will be read and seriously considered by many others besides all thoughtful radiologists. It is unfortunate, therefore, that there are some inaccuracies in the paper, and a strange statement relating to the policy of the Faculty of Radiologists which is a misconception—strange in that none of the four authors have anything to do with the Faculty's policy and only one is a member of the Faculty.

They state "It is understood that the Faculty of Radiologists have it in mind to reduce the total training period required for their Fellowship in Radiodiagnosis to three years . . . this will obviously lower its academic status." A document which will be published in June 1971 and which has already had confidential prior circulation to certain individuals, including one of the authors, states in relation to a new proposed examination and training structure "It will be seen, therefore, that a candidate could obtain the F.F.R. in a three-year training period. In view of the fact that it is the intention to maintain the same standard in the F.F.R. as at present, it seems likely that many candidates will not present themselves for examination until they have achieved a four-year experience in radiology." It also seems likely that the majority of heads of departments will organize the programmed experience of trainee radiologists working for the F.F.R. to cover a minimum of four years as at present. The intention of the faculty is to provide the opportunity for the outstanding candidate to sit the examination earlier if he should so wish.

If the need (or lack of need) for the F.F.R. is such as is implied by the authors in their paper it is strange that the numbers entering for the examination are still increasing. In April 1971 the highest number ever, 71 candidates in radiodiagnosis, entered the examination. It is equally interesting to review the figures for England and Wales for 1970. In that year 39 consultant appointments in radiodiagnosis were made and of those appointed 26 held the F.F.R. In Scotland, of six appointments made two had the F.F.R., and in Northern Ireland three were appointed all holding the F.F.R. (total for U.K. 48 appointments, 31 holding the F.F.R.).

Dr. Whitfield and his colleagues state: "Nine senior registrars posts in the (Birmingham) Region are vacant at the moment." At the time of the faculty's annual census of radiological posts on 1 January 1971 there were seven vacant senior registrar posts in the Birmingham Region, but only a further seven vacant senior registrar posts in the remainder of England and Wales, and of these latter, five had been vacant less than eight weeks and were presumably in process of being filled. The paper states that those attending the D.M.R.D. course in the Birmingham Region " . . . are given paid supernumerary registrar posts . . . " In this they differ from the remainder of England and Wales. In other training