abdominal pressure of between 4 and 8 cm H₂O (see Chart). These increases were accompanied by significant rises in central and femoral venous pressures and by smaller (but still significant) rises in intrathoracic and peak airway pressures. There was an increase in heart rate and mean blood pressure of the order of 10%. End-tidal CO₂ concentration rose by 0.39% ($\approx$ 2 mm Hg). No cardiac dysrhythmias occurred.

Discussion

Our results show that peritoneal insufflation of CO₂ causes a rise in central venous pressure (about one-third as great as the rise of intra-abdominal pressure) accompanied by a smaller rise in intrathoracic pressure, suggesting an increase in right atrial filling pressure. This would be expected to produce an increase in cardiac output (Guyton, 1963) which would be accentuated by the simultaneous increase of arterial PCO₂ (Prys-Roberts *et al.*, 1967). The observed increases in pulse rate and mean arterial blood pressure support this contention.

We attempted to measure cardiac output in these patients by dye dilution with indocyanine green and a photoelectric earpiece, and though interpretation of the dye curves was confounded by stasis of blood in the microcirculation of the ear pinna, our findings are consistent with an increase in cardiac output of the order of 20%. Hodgson *et al.* (1970) found a 25% increase in cardiac output in one patient under similar conditions.

Desmond and Gordon (1970) postulated that increased intra-abdominal pressure might cause a reduction in cardiac output by impedance of "venous return," analogous to that which occurs in the supine hypotensive syndrome of pregnancy (Holmes, 1960). We found no evidence for this at intra-abdominal pressures up to 20-25 cm H₂O, and our results suggest that cardiac output may be increased by this procedure. However, our unpublished observations in a smaller number of patients in whom the intra-abdominal pressure was increased above 30 cm H₂O suggest that under these circumstances blood may be dammed back in the legs with a consequent reduction of central venous pressure and presumably of cardiac output.

There are several possible anaesthetic techniques for laparoscopy. The one used in this investigation would seem to be generally satisfactory, since our results show that the patients whom we studied were able to compensate adequately for the stress imposed on their cardiovascular system by laparoscopy. The patients did not receive inhalational anaesthetic agents, such as halothane, which are known to cause cardiovascular depression. They were artificially ventilated so that CO₂ retention was not a problem. They were also well relaxed so that good visualization of the pelvic contents could be obtained without a large increase in intra-abdominal pressure. With this technique the high intra-abdominal pressures used by other workers—for example, up to 50 cm H₂O (Baratz and Karis, 1969), and up to 40 cm H₂O (Steptoe, 1967;—are unnecessary.

Further studies of the cardiovascular effects of peritoneal insufflation of CO₂, including measurements of cardiac output, are continuing.

We thank Dr. A. M. Michie, medical superintendent, Aberdeen Royal Infirmary, for supplying the indocyanine green dye used in this work. The study was aided in part by a grant from the Scottish Hospitals Endowment Research Trust (to G.R.K.).

References

- Arthure, H. (1970). *British Medical Journal*, 4, 492.
- Baratz, R. A., and Karis, J. H. (1969). *Anesthesiology*, 30, 463.
- Desmond, J., and Gordon, R. A. (1970). *Canadian Anaesthetists' Society Journal*, 17, 378.
- Guyton, A. C. (1963). *Cardiac Output and its Regulation*. Philadelphia, Saunders.
- Herrera-Llerandi, R. (1971). *British Medical Journal*, 2, 47.
- Hodgson, C., McClelland, R. M. A., and Newton, J. R. (1970). *Anaesthesia*, 25, 382.
- Holmes, F. (1960). *Anaesthesia*, 15, 298.
- Jacobaeus, H. C. von (1910). *Münchener medizinische Wochenschrift*, 57, 2090.
- Loveday, R. (1971). *British Medical Journal*, 1, 348.
- Milic-Emili, J., Mead, J., Turner, J. M., and Glauser, E. M. (1964). *Journal of Applied Physiology*, 19, 207.
- Muldoon, M. J. (1971). *British Medical Journal*, 1, 51.
- Prys-Roberts, C., Kelman, G. R., Greenbaum, R., and Robinson, R. H. (1967). *British Journal of Anaesthesia*, 34, 533.
- Seed, R. F., Shakespeare, T. F., and Muldoon, M. J. (1970). *Anaesthesia*, 25, 223.
- Steptoe, P. C. (1965). *Journal of Obstetrics and Gynaecology of the British Commonwealth*, 72, 535.
- Steptoe, P. C. (1967). In *Laparoscopy in Gynaecology*, p. 32. Edinburgh, Livingstone.
- Steptoe, P. C. (1969). *Proceedings of the Royal Society of Medicine*, 62, 439.

MEDICAL MEMORANDA

Adipsia and Hypothermia after Subarachnoid Haemorrhage

S. G. SPIRO, J. S. JENKINS

British Medical Journal, 1971, 3, 411-412

The virtual inaccessibility of the diencephalon to direct study has allowed detailed knowledge of the functions of the hypothalamus in man to accumulate only slowly. Understanding of the pathogenesis of human hypothalamic disease has therefore

heavily depended on experimental analysis in animals. During recent years such animal work has resulted in the definition and understanding of several classical problems of the hypothalamus. Jenkins *et al.* (1969) produced biochemical evidence of hypothalamic-pituitary-adrenal disturbance after subarachnoid haemorrhage. The present case showed clinical features attributable to a lesion of the anterior hypothalamus.

Case Report

A 52-year-old woman with no known history of hypertension had a subarachnoid haemorrhage on 15 March 1969. On admission to hospital she was comatose and had a blood pressure of 180/120. Examination showed no focal neurological signs but she had two large subhyaloid haemorrhages in the left fundus and scattered small haemorrhages in both fundi. Lumbar puncture produced blood-stained cerebrospinal fluid.

Bilateral carotid angiography showed a moderately-sized irregular anterior communicating artery aneurysm filling spontaneously from the left. Bilateral subclavian angiograms were negative. The aneurysm was clipped through a right frontal craniotomy. When

St. George's Hospital, London S.W.1

S. G. SPIRO, B.Sc., M.R.C.P., Medical Registrar
J. S. JENKINS, M.D., F.R.C.P., Consultant Physician

discharged she was fully conscious but mentally very lethargic, slow, and unemotional, lacking any spontaneous conversation. Her left visual acuity was reduced to finger counting only.

She was admitted to hospital 10 months later after three further admissions to Atkinson Morley's Hospital during the previous eight months because of her continuing confusion, disorientation, and deterioration in mental ability. Further specialized neurological investigations confirmed obliteration of the aneurysm and the absence of any other lesion, such as another aneurysm or hydrocephalus.

INVESTIGATIONS AND MANAGEMENT

The outstanding clinical observation was that the patient had no spontaneous desire to drink, and yet she ate her meals spontaneously and fully. It was consistently found that when her fluid intake fell below 1,500 ml/24 hours for more than a few days her serum sodium concentration rose from levels of 140-144 mEq/litre to 150-160 mEq/litre. This would be associated with a corresponding decrease in urinary output and an increase in urinary osmolality. When she was dehydrated her confusion increased, with further disorientation in time and space. Throughout the investigation her oral temperature ran between 34.8° and 35.8°C.

The following tests on the hypothalamic pituitary axis were made. (1) Diurnal rhythm (plasma cortisol, µg/100 ml): 9 a.m. 14.5 (normal 5.5 to 23.2); 12 midnight 18.5, reversed rhythm; after 2 mg of dexamethasone at 12 midnight, 4.4 at 9 a.m. (normal). (2) Metyrapone test (urinary 17-oxogenic steroids): control, 15.4 mg/24 hours; change after 24 hours of metyrapone -6.5 mg/24 hours (normal range + 8.5 to 55.0); the resting level was normal but there was no increase after metyrapone. (3) Insulin hypoglycaemia: control cortisol 16.5 µg/100 ml, maximum increase 10.0 µg/100 ml (normal range + 7.5 to 30.3). (4) Intravenous pyrogen: control (a) 16.4 µg (b) 17.2 µg/100 ml; maximum increase 3.8 µg/100 ml (normal range + 8.6 to 45); deficient responses. (5) Vasopressin test: control 26.3 µg/100 ml, change -4.2 µg/100 ml (normal range + 8.3 to 32); absent response. (6) Twenty-four-hour urinary aldosterone excretion 4.8 µg/24 hours (normal range 4 to 18). (7) P.B.I. 4.8 µg (normal range 4 to 8). (8) Four-hour iodine uptake 18% (normal range 10 to 33%).

Comment

The hypothalamus lies close to the circle of Willis, and Crompton (1963) found that 61% of patients dying shortly after a ruptured berry aneurysm had ischaemic or haemorrhagic lesions in the anterior hypothalamus. Jenkins *et al.* (1969) showed that many patients immediately after a subarachnoid haemorrhage had evidence of a disturbance in hypothalamic-pituitary-adrenal function though none showed any clinical evidence of hypopituitarism. Of 18 of their patients 11 had an abnormal diurnal rhythm within the first 27 days of their bleed, but during a further 28-day period all but two patients had reverted to a normal rhythm. Our patient still had a reversed rhythm after one year.

All but four of the patients in the series reported by Jenkins *et al.* showed a normal dexamethasone response—as also did the present patient—and 3 out of 15 also showed a similar unresponsiveness to metyrapone two months after their haemorrhage.

The exact anatomical pathways concerned in maintaining a normal diurnal rhythm of plasma cortisol are not fully established, but Krieger and Krieger (1966) reported that these disturbances in rhythm were seen only in patients whose lesions were limited to the hypothalamus, prethalamus, and temporal

lobe. As both metyrapone and dexamethasone depend for their effect on the integrity of feedback controls of corticotrophin release, whose site is widely believed to be in the hypothalamus (Reichlin, 1963), the abnormal response to metyrapone suggests a disruption of at least one of these pathways.

Hypoglycaemic Stress.—When subjected to the stress of insulin-induced hypoglycaemia the patient showed a normal hypothalamic-pituitary-adrenal axis response. Similarly, Krieger *et al.* (1968) found a normal response in all seven patients with hypothalamic disease who were studied and who had an abnormal diurnal variation. The failure to react to the stress of intravenous pyrogen is not surprising, as its effect is produced almost certainly from a different hypothalamic centre from that which responds to insulin. Evidence for the initiation of the pyrexia produced by pyrogens on the central thermoregulators in the anterior hypothalamus was produced by Cooper *et al.* (1967) by the direct injection of leucocyte pyrogen into the hypothalamus in animals. The most rapid and sensitive response was from the preoptic and anterior areas.

Body Temperature.—The homeostasis of body temperature seems to be the function of the anterior hypothalamus and the adjacent preoptic areas (Cooper, 1966). Hence damage to this area may produce a subnormal body temperature—as in the case studied by Fox *et al.* (1970). Their patient had a temperature of 34.4° to 36.7°C during the three years studied, and at necropsy the only significant abnormality was glial scarring with astrocytic hypertrophy and neural loss in the anterior hypothalamus, optic chiasma, and tracts. The patient also had an abnormal diurnal rhythm and a deficient response to vasopressin, as in the present case.

Thirst.—The secretion of vasopressin is determined by the supraopticohypophyseal neurones, their endings forming the neural lobe of the pituitary. The present patient showed normal plasma and urine osmolalities and water conservation responses to the stress of dehydration, indicating a normal vasopressin mechanism. Urinary aldosterone excretion was also within normal limits. Andersson and McCann (1956) showed in dogs that lesions produced in the anterior hypothalamus caused a state of hypodipsia, and damage to the "thirst centre" will produce unresponsiveness to dehydration. This appears to be separate from the supraoptic centre, where a lesion can cause diabetes insipidus.

The disorganization of this patient's response to thirst and to pyrogen and her subnormal body temperature can therefore be explained by damage to the anterior and preoptic areas of the hypothalamus. Also her responses were preserved to insulin-induced hypoglycaemia, hunger, water homeostasis, and dexamethasone—supposedly controlled by the more ventral and posterior hypothalamic nuclei.

References

- Andersson, B., and McCann, S. M. (1956). *Acta Physiologica Scandinavica*, **35**, 312.
- Crompton, M. R. (1963). *Brain*, **86**, 301.
- Cooper, K. E. (1966). *British Medical Bulletin*, **22**, 238.
- Cooper, K. E., Cranston, W. I., and Honour, A. J. (1967). *Journal of Physiology*, **191**, 325.
- Fox, R. H., Davies, T. W., Marsh, F. P., and Urlich, H. (1970). *Lancet*, **2**, 185.
- Jenkins, J. S., Buckell, M., Carter, A. B., and Westlake, S. (1969). *British Medical Journal*, **4**, 707.
- Krieger, D. T., Glick, S., Silverberg, A., and Krieger, H. P. (1968). *Journal of Clinical Endocrinology and Metabolism*, **28**, 1589.
- Krieger, D. T., and Krieger, H. P. (1966). *Journal of Clinical Endocrinology and Metabolism*, **26**, 929.
- Reichlin, S. (1963). *New England Journal of Medicine*, **269**, 1182.