incidence of infertile marriage which he said had been compiled by me. If Dr. Brown is correctly reported then his statement, with the implication that it indicates the present state of affairs, is misleading.

The statistics he quoted (incompletely and inaccurately in one respect) were apparently taken from my book.¹ They are to be found in a paragraph which makes it clear that they apply not to the present or the immediate past but to the decade 1900-1909. This is before my time.—I am, etc.,

NORMAN JEFFCOATE

Department of Obstetrics and Gynaecology,
The University,
Liverpool

¹ Jeffcoate, T. N. A., *Principles of Gynaecology*, 3rd edn., Butterworth, London, 1967.

Mechanism of Action of Amantadine

SIR,—It has been well established that amantadine is a moderately effective drug in patients with Parkinsonism.^{1,2} Little is known of its mode of action, and reports of side effects of hallucinosis, confusion, insomnia, and dizziness are so non-specific as to shed little extra light on the level or site of central action.

A 68-year-old lady with severe rigid and akinetic Parkinsonism, previously untreated, was given amantadine 100 mg b.d. Her Webster score improved from 18/30 to 14/30 in one week, and six weeks later her score was 10/30 and she was mobile and much less disabled. She complained, however, that "I can't keep my tongue still in my mouth," and examination confirmed a gross lingual-facial dyskinesia, precisely mimicking that seen with levodopa.

Amantadine has been shown in animal studies to potentiate catecholaminergic (including dopaminergic) transmission; this may be due to inhibition of neuronal uptake across the synaptic cleft, or it may release catecholamines from the small extragranular depots.³ Locomotor behaviour is increased to a much lesser extent than with amphetamines in animals, but of interest is the report⁴ that it potentiates stereotyped behaviour (licking and gnawing) in rats and mice. This pattern of stereotyped motor behaviour has been related to dopamine activity,⁵ and the case summarized here may represent the first parallel to this observation in the human subject. The evidence suggests that the augmentation of dopaminergic transmission is more likely to be due to release from extragranular stores than from the larger granular stores, since amantadine has no effect on raising cerebrospinal fluid levels of homovanillic acid derived from dopamine breakdown.—I am, etc.,

JOHN PEARCE

Combined Neurological Service,
Hull Royal Infirmary,
Hull, Yorks

¹ Parkes, J. D., et al. *Lancet*, 1971, 1, 1083.
² Rao, N. S., and Pearce, J., *Practitioner*, 1971, 206, 241.
³ Offermeier, J., *Parkinson's Disease*, Ed. G. Birdwood, S. Gilder, and C. A. S. Wink, London. Academic Press, 1971, p. 85.
⁴ Simon, P., Malatray, J., and Borssier, J. R., *Journal of Pharmacy and Pharmacology*, 1970, 22, 546.
⁵ Ernst, A. M., and Smelik, P. G., *Experientia (Basel)*, 1966, 22, 837.

Back to Vesalius, 1970

SIR,—A great deal has been written in recent years about the importance of impact, relevance, and the integrative approach in medical education especially in that Cinderella of the basic sciences, gross anatomy. It is often argued that familiarity with the appearance of the parts of the body in the dissecting room is of little help in visualizing these structures in the living or for their identification at operation.

The interest and enthusiasm shown by the University of Iowa freshmen medical students at the routine demonstrations of fresh human placenta as part of their integrated course in gross anatomy and human embryology caused some surprise, since the interest seemed out of proportion to the importance of the gross structure of the organ in question. It was therefore decided, in the interest of the fresh approach as well as in the ever continuing quest for the new and the unusual, to go back to Vesalius.¹

The subject was a 65-year-old male, whose previously deeded body became available when he died instantaneously after impact with a telegraph pole. He was received, unembalmed, very shortly after death, without a previous postmortem examination, and was immediately placed in cold storage. As the students had already completed dissection of upper limb, head and neck, thorax, and the greater part of abdomen, it was considered a suitable time for the demonstration, both from their anatomical knowledge and the familiarity with which they regarded embalmed bodies. The subject had been in cold storage for four days when dissection began. The experiment was terminated after 2½ days, because at the end of this period (despite cold storage when the class was not present) the odour rose above the nuisance level. We are not prepared to comment whether the factor responsible for a tolerable dissecting period, considerably shorter than that customarily employed by our ancestors, was a winter dissecting room temperature of 75° F. (24° C.) or merely a simple lack of intestinal fortitude.

The brunt of demonstration on this cadaver was carried out by the medically qualified members of the gross anatomy teaching group (four out of eight). However, all members of the group (who are always present during scheduled gross anatomy sessions) were available to demonstrate and discuss the unembalmed body during the entire time it was on display. Any student who wished to examine the specimen alone, or in company with his fellows without an instructor, was encouraged to do so. An unlimited supply of disposable surgical gloves was provided for this purpose.

Student comments, derived from a questionnaire were, on the whole, favourable, and 81% of the 146 students felt that the demonstration was valuable, while 44% would have liked the experimental scope enlarged to include all regions of the body. Thirty-five per cent. expressed interest, but felt that they had insufficient time to examine the subject fully, owing to upcoming exams and the pressure of work, while only 5% found the unembalmed body "objectionable" (a similar percentage were "upset" by post mortems in the second semester of the freshmen year) and a final 3%, though not bothered by aesthetic considerations, found it to be "of no value."

These comments suggest that separate regional demonstrations synchronized with student dissections might be an effective method for future presentation.—We are, etc.,

N. F. METCALF
W. K. METCALF
D. J. MOFFATT

Department of Anatomy,
University of Iowa,
Iowa City, Iowa,
U.S.A.

¹ Ball, J. M., *Andreas Vesalius: The Reformer of Anatomy*, p. 75. St. Louis, U.S.A. 1910.

Echovirus Type 4 Infections

SIR,—Further to the current outbreaks in Tees-side and in Northern Ireland recently referred to (10 July, p. 124),^{1,2} it may be noted from the *Weekly Communicable Diseases (Scotland) Reports* that this virus is currently also active in Scotland. In this area it was first observed in 1963³ when it produced an outbreak with an older age distribution than the more common enteroviruses.⁴ Except for a few isolations in 1964 it was not encountered again until 1968, since when only a few strains have been isolated each year until 1971, with 27 isolations reported to date.

In early April, three strains were isolated in Inverness from one family in Ross and Cromarty, but the main focus has been in Lanarkshire, particularly in the Coatbridge/Airdrie area, with 16 isolations. Other isolations have been reported in recent weeks in Glasgow (five cases) and in Ayrshire (three cases), indicating further spread of infection. In addition, three other cases from Glasgow have been diagnosed by a four-fold rise in antibody titre. In 25 instances, the virus has been isolated from faeces, in one of which isolation was also reported from respiratory secretions. In two cases the virus was obtained from cerebrospinal fluid, in one of which isolation was also reported from a throat swab.

With the exception of three cases, all have suffered from aseptic meningitis, although in every instance as elsewhere recovery has been almost complete within 48 hours and no after effects reported. Of the 30 cases, there have been 16 males and 14 females with a peak incidence in the same 5-20 age groups as in Tees-side and Northern Ireland.

This report was compiled from information provided by contributors to the *Weekly C.D.S. Reports*.

—I am, etc.,

J. C. M. SHARP

Communicable Diseases (Scotland) Unit,
Ruchill Hospital,
Glasgow N.W.

¹ *Lancet*, 1971, 2, 52.
² Donaldson, R. J., *Lancet*, 1971, 2, 108.
³ Bell, E. I., *Lancet*, 1964, 1, 1195.
⁴ Landsman, J. B., Bell, E. J., and Grist, N. R., *Health Bulletin*, 1966, 24, 38.

Prophylaxis of Venous Thrombosis

SIR,—Your opening sentence of the leading article "Prophylaxis of Venous Thrombosis" (6 February, p. 305) is, to say the least, timely, and the need of prevention of deep vein thrombosis of the leg and possible fatal thromboembolism extremely urgent.

The second part of the same sentence states that "no one can yet offer a practical solution." You refer to the method advocated by myself and G. DeBlasio,¹ and conclude that "trials with the new method of assessment are not yet available." At the time of your writing you could not be aware of a trial of our method now in progress under the direction of Mr. J. Lannon of the Lady Dudley Hospital in Johannesburg, South Africa, for he has as yet not published his results of the past five years.

In a letter dated 2 July 1971 he states that as of that date he had used the method now in 749 operative patients all elderly with "not a single case of pulmonary embolism or leg vein thrombosis." He also states that he will not be without heparin prophylaxis in performing surgery. His experience parallels ours with the same success attained by us now, in 926 usually at risk patients. Before the use of heparin prophylaxis Mr. Lannon had indicated he had as we had experienced a 4% incidence of fatal thromboembolism in his operative patients.

I have no knowledge at this writing of others who presently use our prophylactic method and what their results are. I would also be happy to see a report of Mr. Lannon's experience but to-date he has not found the time to publish.

So perhaps you may find Mr. Lannon's assessment of our heparin prophylaxis as offering a "practical solution" to the prophylaxis of venous thrombosis for the operative patient at least.—I am, etc.,

J. GEORGE SHARNOFF

Department of Pathology,
Mount Vernon Hospital,
New York, U.S.A.

¹ Sharnoff, J. G., and DeBlasio, G., *Lancet*, 1970, 2, 1006.

Aplastic Anaemia and Leukaemia

SIR,—A case resembling those presented by Drs. I. W. Delamore and C. G. Geary (26 June, p. 743) has been reported by Gmyrek and others.¹ An 8-year-old girl with Fanconi type of aplastic anaemia received prednisone and an anabolic steroid and developed after a period of three years acute leukaemia of myelomonocytic type. Although Fanconi's anaemia might be potentially more leukaemic than aplastic anaemia of other type, there are only a few reports of this occurrence.—I am, etc.,

M. I. SAARNI

Department of Medicine,
Central Hospital,
Kotka,
Finland

¹ Gmyrek, D., Witkowski, R., Syllm-Rapoport, L., and Jacobasch, G., *Deutsche medizinische Wochenschrift* 1967, 92, 1701.

Gastric Ulcer and Gastritis

SIR,—Your leading article (10 July, p. 62) draws attention to the association of gastric ulcer and atrophic gastritis and cites evidence suggesting that the latter precedes the former. Whatever the chronological connexion between the two conditions, the observation is of interest for the pathogenesis of gastric ulcer because it is not long ago that both duodenal and gastric ulcers were classified as peptic ulcers. Hyperacidity whether of

nervous or hormonal origin was believed to play an important part in their pathogenesis.

It is timely to go further and to suggest that not only have duodenal ulcers and the primary gastric ulcer a different pathogenesis but a clear distinction should also be made between the primary (classical) gastric ulcer and those which occur in association with lesions in the pyloric antrum or the duodenum (pyloric channel syndrome). While in the former a normal or more often a low gastric acidity is found, the same does not apply to gastric ulcers in the pyloric channel syndrome.

There is a distinct difference in the age distribution between the types of gastric ulcer. Bateson¹ found that duodenal ulcer reaches a peak in patients between 30 and 60 years, concomitant gastric and duodenal ulcers between 40 and 60 years, and solitary gastric ulcer between 50 and 70 years. My own series² of 350 cases with primary chronic gastric ulcer shows likewise that these ulcers are predominantly seen in late middle and in old age. It is therefore difficult to ascertain the significance of atrophic gastritis or of pyloric incompetence in the pathogenesis of these ulcers, for both conditions are frequently manifestations of physiological involutionary processes.

An outstanding feature in chronic primary gastric ulcer is the high incidence of calcified atherosclerotic lesions in the central blood supply of the stomach. Though atherosclerotic lesions in the abdominal aorta are a common finding in advancing years, I found³ that while 65% of patients with the classical gastric ulcer show calcified lesions of various degrees in the abdominal aorta between the ages of 50 and 60 years, these lesions are seen in only 23% of control cases of the same age group. Moreover, a detailed radiological study of the distribution of the calcified atheroma in gastric ulcer cases often reveals their location not only in the proximal portion of the abdominal aorta but also in the coeliac axis.³ Thus the classical chronic gastric ulcer seems to be a complication of atherosclerosis like coronary, internal carotid, or renal artery disease, where arterial insufficiency and obstruction are less often caused by involvement of the small arteries immediately supplying the affected organ but rather by atheromatous plaques in the more central arterial supply.—I am, etc.,

A. ELKELES

London N.W.8

¹ Bateson, E. M., *British Journal of Radiology*, 1969, 42, 598.

² Elkeles, A., *American Journal of Roentgenology*, 1964, 91, 744.

³ Elkeles, A., *Journal of the American Geriatric Society*, 1970, 18, 450.

Reflux and Hernia

SIR,—In your leading article entitled "Reflux and Hernia" (24 July, p. 205) you draw attention to recent work by Cohen and Harris¹ and comment that some of their views will cause consternation among the traditionalists.

This may, I suppose, be true; but they will not, I am sure, cause consternation among many surgeons with experience of operating on cases of gastro-oesophageal reflux. The occurrence of reflux without herniation has, of course, been found by many observers to be not so very uncommon,

even if less common than Cohen and Harris's findings might seem to suggest. And after conventional repair operations with restoration of normal anatomical relationships, and excluding cases with hernia recurrence, reflux symptoms, and perhaps demonstrable reflux, persist in many cases. In my own experience this occurred in some 66% of such cases, and, though symptoms were often much improved, in over 20% they were little, if at all, improved.

You quote Cohen and Harris as challenging the rationale of surgical repair of hiatus hernia. Their findings certainly challenge the rationale of the conventional type of repair. But I regard them as providing support for those of us who for some time have held that restoration of normal anatomy may not be followed by restoration of normal antireflux function and who, because of disappointing results from conventional repair operations, have adopted one of the gastroplication procedures. I hold that these procedures work so well because they provide a completely new antireflux mechanism—a valve mechanism. It is with interest, therefore, that I note Cohen and Harris's finding that surrounding pressure does not increase sphincter strength. In the plication operation I carry out, in fact, the oesophago-gastric junction is not over-reduced, but fixed in its normal position, as with conventional repair operations, and the new antireflux plication arrangement is constructed beyond the cardia. To date, this appears to be completely effective in controlling reflux.

By carrying out this procedure on cadaveric stomachs, and testing them at the laboratory bench it can readily be demonstrated that the flow of fluid from the stomach into the oesophagus is prevented. No part can be played in this by sphincteric action or by intra-abdominal pressure. It would seem that it must be due to the production of a new antireflux mechanism—a valve mechanism.

The conclusion to be drawn would seem to be that when gastric reflux causes troublesome symptoms or oesophageal damage, whether associated with a hernia or not, what is required is the creation of a new antireflux mechanism. Fortunately this can be done.—I am, etc.,

J. A. W. BINGHAM

Royal Victoria Hospital,
Belfast

¹ Cohen, S., and Harris, L. D., *New England Journal of Medicine*, 1971, 284, 1053.

Pupillary Paralysis after Tranquillizer

SIR,—I would like to report on an unusual side effect apparently due to pimozide (Orap, Janssen Pharmaceuticals). The patient was a 24-year-old female, diagnosed as suffering from simple schizophrenia. She was treated with phenothiazines for three weeks (initially thioridazine, followed by trifluoperazine), without improvement in her clinical state. She was started on pimozide 6 mg in the morning, and after two weeks the dosage was changed to pimozide 4 mg in the morning, 2 mg at night. There was some improvement in her symptoms, and after three weeks the dosage was increased to pimozide 4 mg b.d. Six days later she complained of mild Parkinsonian tremor of the hand and leg and immobile facies. This responded to