field testing also gives an unacceptably high number of false positives. Examination of the fundus by a skilled ophthalmologist seems to be the most reliable single test for glaucoma, but this is impractical as a screening procedure.

At the BUPA Medical Centre we have in the last year carried out just over 1,000 routine analyses of visual fields in middle aged men, using a Friedmann visual field analyser.⁴ Three patients were found to have significant defects: one due to myopia, one due to hypertensive exudates, and one due to a local vascular lesion (these diagnoses were made by our consultant ophthalmologist). In addition, nine isolated peripheral lesions were found, which are almost certainly of no significance.

We found the instrument easy to use and acceptable to staff and patients. However, as no case of glaucoma was found, the test is clearly not worth using routinely for screening purposes. We intend continuing the survey, restricting the test to patients aged 65 or over, or those with a positive family history of glaucoma.

Other studies using this or similar instruments have found yields of 0.7%¹ and 0.6%⁵ and of 0.7%² using tonometry. The total absence of chronic simple glaucoma in our population is most surprising. It is possible that they have been missed owing to technical error; however, as three cases of non-glaucomatous field defects were subsequently confirmed this seems unlikely. As the condition is asymptomatic at this stage any form of conscious selection is unlikely. The possibility remains that glaucoma may be less common in our social class 1 and 2 population than in the country as a whole.—I am, etc.,

G. PINCHERLE

The Medical Centre,
London N.1

¹ Graham, P. A., *Proceedings of the Royal Society of Medicine*, 1966, 59, 1215.

² Banks, J. L. K., Perkins, E. S., Tsolakis, S., and Wright, J. E., *British Medical Journal*, 1968 1, 791.

³ Graham, P. A., and Holows, F. C., in *Glaucoma*, Ed. L. B. Hunt, P. 103, Edinburgh, Livingstone, 1966.

⁴ Bedwell, C. H., *Ophthalmic Optician*, 1967, 7, 566 and 616.

⁵ Linfield, P. B., *Ophthalmic Optician*, 1970, 10, 1044.

Sleep and a Bedtime Beverage

SIR,—The critical letter from Miss Aviva Petrie (10 June, p. 652) following the paper by Dr. Vlasta Březinová and myself (20 May, p. 431) raises principles of research strategy. We had used electrophysiological techniques to compare sleep after 250 ml of hot milk, containing 32 g of Horlicks powder, with sleep after an inert capsule. An indistinguishable control was not feasible, and people today are attuned to pills as sleep improvers. There is, actually, no evidence that suggestion will improve sleep when measured electrophysiologically. The sleep of poor sleepers has been reported not altered by blank sleeping pills.¹ Miss Petrie says we should have used hot water and milk as additional variables, evidently thinking nutrient control less important.

It would indeed be nice to use the additional controls she suggests, as well as other proprietary food drinks, bread and cheese, fish and chips, black pudding, and other bedtime snacks, with varying doses of each. In high-virgin territory, however, demands

must be simple. Too complex an initial design merely wastes resources. We asked whether we could find any evidence to support a report that a concentrated, palatable, liquid food before sleep could make sleep less restless. We used two conditions. Had we used four, that would have meant not 18 experimenter-months and 60 miles (96 km) of recordings, but 3 years and 120 miles (200 km).

Miss Petrie, a statistician, would probably say she could have offered us a neat design embodying four conditions and no additional nights, but at the cost of reduced nights per condition. Then the study would have drawn a blank. We found again, as we have often found before, that sleep under a first laboratory condition is not the same as when the individual has had the same condition repeatedly.

No, only when one has got a simple first answer, and learned from practical experience and inevitable discussion and rumination, can one justifiably commit expensive resources to answer the new questions that present themselves.—I am, etc.,

IAN OSWALD

Department of Psychiatry,
University of Edinburgh

¹ Kales, J., Kales, A., Bixler, E. O., and Slye, E. S., *Clinical Pharmacology and Therapeutics*, 1971, 12, 691.

Febrile Convulsions in Early Childhood

SIR,—I would like to take up two points from your leading article "Febrile Convulsions in Early Childhood" (10 June, p. 608).

The statement that "continuous prophylactic antiepileptic treatment should not be given: it is unnecessary and unhelpful" is too dogmatic. No satisfactory trial has yet been described. Millichap and others¹ compared continuous prophylactic phenobarbitone given only during febrile episodes and found no difference, but their sample was small and there was no control group. One would not recommend continuous prophylaxis after one or two febrile convulsions, but for the child who is having frequent convulsions, often occurring before anticonvulsants can be administered, continuous prophylaxis may be effective.

The desire to exclude the possibility of pyogenic meningitis should not require resort to lumbar puncture in every case. If one raises the age limit for "febrile convulsions" from 6 months to 10 months, then there should be no difficulty in differentiating pyogenic meningitis clinically. Convulsions in meningitis are usually due to cerebral oedema and one would not expect such a child to recover from a convulsion in the way that a child with a simple "febrile convulsion" does. In fact, a careful search of the memories of two senior consultant paediatricians revealed no instance of pyogenic meningitis presenting as a "febrile convulsion." It is mandatory, however, to perform a lumbar puncture if there is the slightest doubt.—I am, etc.,

B. W. LEWIS

Children's Department,
London Hospital,
London E.1

¹ Millichap, J. G., Aledort, L. M., and Madsen, J. A., *Journal of Paediatrics*, 1960, 56, 364.

SIR,—The leading article (10 June, p. 608) describing febrile convulsions in early child-

hood was informative, but the advice on treatment was a little inconsistent with the facts given earlier.

If a child is unwell before the convulsion, the fit does not last more than 10 minutes, and is not followed by postepileptic hemiplegia then it fits the diagnosis of a febrile convulsion. If the cause of the pyrexia is obvious—for example, otitis media, and clinical signs of meningitis are absent why should the child be sent immediately to hospital? Tepid sponging and antibiotics are available at home. It would be interesting to know how many paediatricians would perform a lumbar puncture in such a case as the one described above.

As febrile convulsions almost always occur only once in each febrile episode even the "traditional" paraldehyde may be a case of "shutting the stable door."—I am, etc.,

DONALD MORTON

Middlesbrough,
Teesside

SIR,—Your recent leading article on febrile convulsions (10 June, p. 608) though excellent should not pass without comment.

In the final paragraph are two statements which I would question. Surely there are occasions, for instance where attacks are frequent, when it is undoubtedly best both for the child itself and the family, if the parents, with of course medical help, learn to deal with the problem at home. Secondly, is paraldehyde, unpleasant stuff even in the old days, really still the drug of choice for acute convulsions? I have not seen a glass syringe in years and I cannot believe that many doctors in Britain carry them these days.

No drug is completely without danger, but the popular and very effective treatment in Denmark seems to be intramuscular or intravenous diazepam with which many would combine intramuscular phenobarbitone to avoid the danger of the sometimes rather short therapeutic action of the former drug given alone.—I am, etc.,

SIMON SPRAGGE

Children's Department,
Centralsygehuset,
Slagelse, Denmark

Mode of Action of Verapamil

SIR,—I have read with interest the correspondence concerning verapamil and the heart. Dr. L. Schamroth and others (11 March, p. 660) discussed the possible modes of action of this drug and concluded its action to be uncertain. They were criticized in this by Drs. B. Livesley and S. Oram (1 April, p. 50), who felt that verapamil could act by augmenting vagal tone on the atrioventricular specialized tissue of the heart.

However, in making this suggestion they give no indication as to their source of reference regarding the vagal innervation of ventricular specialized tissue. To the best of my knowledge, morphologic evidence of direct efferent vagal supply to the atrioventricular bundle and its ramifications in the human heart is lacking. This was certainly the conclusion drawn by Vassalle,¹ who stated that while vagal stimulation produced ventricular standstill there was no evidence that this was due to vagal inhibition

of the specialized tissue. He felt that for inhibition to occur it should be shown that the vagus nerve innervated the entire ventricular specialized network, and that acetylcholine should be shown to suppress spontaneous activity of ventricular Purkinje fibres. It would seem that these two conditions also require to be met with regard to verapamil and the human heart before the claims of Drs. Livesley and Oram can be justified.—I am, etc.,

R. H. ANDERSON

Department of Anatomy,
University of Manchester

¹ Vassalle, M., *American Journal of Cardiology*, 1971, 28, 245.

Epidemic Ataxia in Western Nigeria

SIR,—Dr. B. O. Osuntokun (3 June, p. 589) states, "I have found no evidence of viral infection as an aetiological factor [of epidemic ataxia]." It is true that efforts to isolate a virus have so far been unsuccessful, and Dr. Osuntokun is undoubtedly referring to the work of the Virus Research Laboratory of the Faculty of Medicine, University of Ibadan, although he did not say so. In collaboration with the Wesley Guild Hospital, Ilesha, Western Nigeria, we have been investigating the possible viral aetiology of this condition, and the first report is at the moment being prepared for publication. Experience dictates that we should not close our minds to the possibility of viral aetiology, as it is a known fact that in the developing countries, where laboratory facilities are relatively limited, a negative result does not permit one to pronounce a categorical verdict. A "no find" may well be due to a "no extensive" or "no exhaustive" examination. A classical example is the case of dengue viruses in Africa. Physicians and indeed scientists have for a long time concluded that the continent of Africa was a dengue-free zone. But in 1970 from our laboratory dengue viruses were isolated and fully documented.¹

In clinical virology, while there is the danger to look upon viruses as a diagnostic dump heap, there is also the equally dangerous risk of returning a negative verdict too early in the operation. An open mind will help open prospects for future closer examination.—I am, etc.,

TAM S. DAVID-WEST

Virus Research Laboratory,
Faculty of Medicine,
University of Ibadan,
Ibadan, Nigeria

¹ Carey, D. E., Causey, O. R., Reddy, S., and Cooke, A. R., *Lancet*, 1971, 1, 105.

Dressler's Syndrome

SIR,—Dr. M. S. T. A. Lawrence and Professor Ralph Wright (11 March, p. 665) report a case of Dressler's syndrome complicated by tamponade unassociated with haemorrhage into the pericardial space. I would like to report a further case.

A 62-year-old man originally presented in May 1971 with an acute myocardial infarction. At follow-up two months after the first attack he was complaining of tiredness, dyspnoea, and shoulder and back pain. His chest x-ray showed an enlarged cardiac shadow with no sign of pulmonary con-

gestion. He was treated with diuretics and practolol without symptomatic or clinical benefit.

In November 1971 he was first seen at this hospital. At this time his jugular veins were filled to his ears, liver edge palpated 5 cm below the costal margin, 15 mm Hg arterial paradox, but no sign of fluid in his lungs. His chest x-ray showed a large globular heart shadow but no pulmonary oedema. He was admitted for further investigations, which, apart from Hb 12.3 g/100 ml and E.S.R. of 43 mm, were unremarkable.

The diagnosis seemed clear: post-myocardial infarction (Dressler's) syndrome, complicated by tamponade. An attempt was made to aspirate the pericardial fluid but was unsuccessful. He was started on prednisolone 60 mg a day with a brisk improvement in his condition. His jugular venous pressure fell to normal, and his cardiac shadow returned to a normal size and shape. Steroids were reduced to a maintenance dose of 10 mg a day. He has now been well for six months, apart from one episode of pericardial pain and a friction rub occurring a month after discharge, which responded to a brief increase in his steroid dosage. His steroids have now been tailed off.

This man seems a typical example of a postmyocardial infarction syndrome presenting two months after the initial infarct with tiredness, anaemia, a raised E.S.R., and signs of pericardial disease. He had physical signs of cardiac tamponade with x-ray confirmation. It seems unlikely that this could have been due to haemorrhage as he was not on anticoagulants and because he responded so well to steroid therapy. It seems likely that this situation can be complicated by pericardial effusion, severe enough to give tamponade and which responds very well to steroid therapy.

I would like to thank Dr. Turner for his permission to report his case.

—I am, etc.,

B. J. HUTCHCROFT

Edgware General Hospital,
Edgware, Middx

Food and Cyanide

SIR,—I would like to comment on two specific aspects raised in your leading article (27 May, p. 481) which have been of particular interest to me since our original studies on the aetiology of tobacco amblyopia¹ and retrobulbar neuritis in pernicious anaemia.²

Firstly, it is certainly important to remember that, besides being present in tobacco smoke and alcohol, cyanide is widely distributed in the plant kingdom. In this connexion, I contended that retrobulbar neuritis in patients with vitamin B₁₂ deficiency who are non-smokers is the result of exposure to cyanide from some other source than tobacco.³ Abnormally raised plasma-cyanocobalamin levels have been found in some patients with the so-called "West Indian amblyopia"⁴ and in Nigerian patients with tropical amblyopia and ataxic neuropathy.⁵ The ophthalmic findings, as reported, are consistent with the strict criteria we specified and adopted in the diagnosis of tobacco-amblyopia visual fields,¹ and add further support to the hypothesis that exposure to cyanide from the dietary source

of cassava, or other roots or vegetables containing cyanogenic glycosides, contributes to the pathogenesis of tropical amblyopia.

If the indiscriminate dumping of industrial cyanide waste continues unchecked, with the inherent risk of pollution of soil and water supplies, there may well come the time when more widespread chronic cyanide neurotoxicity occurs from a dietary source in persons with a genetic or acquired error of cyanide or vitamin B₁₂ metabolism.

Secondly, I consider that your statement that "although it has been suggested that vitamin B₁₂ has a primary role in the detoxification of cyanide, this is not finally proved" is open to misinterpretation. It is certainly well recognized that cyanide is also converted to thiocyanate by an independent pathway via the enzyme rhodanese, which is present in vivo in high concentration in the liver.⁶ Clinical studies from many centres have shown conclusively that hydroxocobalamin, but not cyanocobalamin, is a powerful cyanide antagonist. Because confusion is likely to persist in the profession over the differences between various forms of vitamin B₁₂ commercially available, we have drawn attention to the superiority of hydroxocobalamin, and the possible adverse effect of cyanocobalamin in neuro-ophthalmological disorders, and submitted a case for the complete withdrawal of cyanocobalamin in favour of hydroxocobalamin for therapeutic use.⁷ In our view, the only indication for the continued availability of cyanocobalamin is its use in the performance of the Schilling test of vitamin B₁₂ absorption.

Looking to the future, it is possible that other cyanide antagonists may become available for our therapeutic use with the obvious advantage of oral administration. In such circumstances, I would strongly urge that all patients with proved tobacco amblyopia and patients with optic atrophy, myelopathy, or neuropathy of obscure origin should be very carefully screened for evidence of vitamin B₁₂ deficiency before embarking on any therapy other than hydroxocobalamin.—I am, etc.,

A. G. FREEMAN

Princess Margaret Hospital,
Swindon, Wilts

- 1 Heaton, J. M., McCormick, A. J. A., and Freeman, A. G., *Lancet*, 1958, 2, 286.
- 2 Freeman, A. G., and Heaton, J. M., *Lancet*, 1961, 1, 908.
- 3 Freeman, A. G., *Lancet*, 1959, 2, 963.
- 4 Wilson, J., Linnell, J. C., and Matthews, D. M., *Lancet*, 1971, 1, 259.
- 5 Osuntokun, B. O., and Osuntokun, O., *American Journal of Ophthalmology*, 1971, 72, 708.
- 6 Smith, A. D. M., *Lancet*, 1961, 1, 1001.
- 7 Foulds, W. S., Freeman, A. G., Phillips, C. I., and Wilson, J., *Lancet*, 1970, 1, 35.

Neuromuscular Involvement in Pituitary Gigantism

SIR,—Dr. P. D. Lewis (27 May, p. 499) describes two cases of pituitary gigantism with neuromuscular involvement. I would like to report a case of similar neuromuscular involvement in a man aged 69 with acromegaly of at least eight years duration.

This man first presented in a Dublin hospital in 1964 with left foot drop; acromegaly was noted for the first time. The acromegaly has not apparently progressed since then, and clinical indices of activity have been negative over the past two years. His feet, however, have got steadily worse.