I would like to see a method instruction sheet enclosed with every condom that is sold. I realize that there would be those who would not read the instructions, and also people who read the instructions but not follow them, but I think there would also be a large number of responsible people who would both read and follow the instructions, and thereby achieve the maximum safety possible with this type of birth control, and reduce the number of unwanted pregnancies.—I am, etc.,

KATHLEEN HUNTINGTON

London S.W.3

Infection in Rheumatoid Disease

SIR,—Your leading article entitled "Infection in Rheumatoid Disease" (3 June, p. 549) included the statement (unsupported by reference to published data), "Similarly, these patients are thought to have an increased incidence of urinary infections and chronic pyelonephritis. . . ." The relevant literature does not contain any data to support this contention. Indeed, Mowat¹ in a controlled study of bacteriuria in patients with rheumatoid arthritis showed that the incidence of urinary tract infection was similar in the patients and control groups, and this result has been confirmed in a larger group of 247 patients with rheumatoid arthritis who were compared with 129 control subjects matched for age, sex, and duration of disease (unpublished). Furthermore, in the latter study there was no evidence that analgesic consumption altered the susceptibility to renal infection, though patients treated with systemic corticosteroid drugs showed a small but statistically significant increase in prevalence of bacteriuria.

The confusion in this matter may arise from reports of an increased incidence of "pyelonephritis" in post-mortem studies of patients with rheumatoid arthritis.²⁻⁴ Since there is no evidence of an increased incidence of urinary tract infection during life, it is very unlikely that the pathological features noted are the result of infection. The term pyelonephritis, with its usual clinical connotation of an infective process, seems inappropriate in these circumstances, and substitution of the more non-committal term "chronic interstitial nephritis" might avoid confusion in the future.—I am, etc.,

H. C. BURRY

Keat's House,
Guy's Hospital,
London S.E.1

¹ Mowat, A. G., Hothersall, T. E., and Gould, J. C., *Annals of the Rheumatic Diseases*, 1970, **29**, 143.

² Lawson, A. A. H., and Maclean, N., *Annals of the Rheumatic Diseases*, 1966, **25**, 441.

³ Duthie, J. J. R., Brown, P. E., Truelove, L. H., Baragar, F. D., and Lawrie, A. J., *Annals of the Rheumatic Diseases*, 1964, **23**, 193.

⁴ Clausen, E., and Pedersen, J., *Acta Medica Scandinavica*, 1961, **170**, 631.

Coal Workers' Pneumoconiosis

SIR,—I am sorry to prolong the correspondence on this subject but the last letter by Dr. J. P. Lyons and others (10 June, p. 654) is so obscure that clarification is required. In my first letter (6 May, p. 353) I simply made the point, subsequently made by Professor I. T. T. Higgins (17 June, p. 713) that the evidence they presented in their paper is

inconsistent with their conclusion that emphysema in their cases was due to pneumoconiosis (which I take to imply a process in the lung due to dust). I am not interested in their "repeated assertions" that the emphysema was due to dust, I am only interested in quantitative evidence. How can the severity of a process due to dust bear no quantitative relationship to the amount of dust in the lung as demonstrated by the x-ray category? The explanation given in the authors' letter is that many of the cases they classified as category 0 had previously been classified as category 2 or more. Since regression of x-ray categories is almost invariably due to observer error it is most likely that there was as much observer error in the categorization of their cases classified as category 2 or 3 and these errors should cancel each other out. The best way of overcoming such errors would be to present mean emphysema counts in relation to mean category of all recent films read independently by two or more observers. If this showed a significant correlation between emphysema count and x-ray category, the authors would have convincing evidence that emphysema is quantitatively related to the amount of dust. This would be consistent with (if not proof of) their assertion which at present remains in conflict with the evidence they have hitherto presented.

Finally, a protest must be registered against their suggestion that cases with irregular opacities were "excluded" from the epidemiological surveys whose findings are in such close agreement with their own observations. Such cases would have been classified as category 0, but this did not affect the low mean dust content found in lungs thus classified.¹—I am, etc.,

C. M. FLETCHER

Department of Medicine,
Royal Postgraduate Medical School,
London W.12

¹ Rossiter, C. E., (1972) *British Journal of Industrial Medicine*, 1972, **29**, 31.

Plantar Warts

SIR,—I was surprised that in your leading article (24 June, p. 725) no mention was made of magnesium sulphate ionization, using the anode as the active electrode, in the treatment of multiple warts. It is safe, easy to apply, and effective.—I am, etc.,

CLIVE SHIELDS

London S.W.3

SIR,—I refer to your leading article on "Plantar Warts" (24 June, p. 725). For treatment you appear to commend a variety of local applications as a first choice. I would submit that these treatments are often tedious, messy, prolonged, and frequently ineffective, and freezing with carbon dioxide or liquid nitrogen usually requires a hospital visit.

The removal of warts by curette gives relief from discomfort, as I can myself testify, within 24 hours. The objection to this method, you imply, is the pain of infiltration of the local anaesthetic. But there is a simple way of achieving anaesthesia for the sole of the foot—namely, blocking of the medial and lateral plantar nerves at the ankle. The technique was shown to me by my friend Dr. R. Bryce-

Smith of the department of anaesthetics at the Radcliffe Infirmary some four years ago. Since then I have employed it for more than 200 patients ranging in age from 4 to 45 years with only two failures to achieve satisfactory anaesthesia.

The method is to inject from 3 ml to 6 ml, according to age, of 1% lignocaine in 1/200,000 adrenaline into the ankle at a point midway between the most prominent part of the medial malleolus and the margin of the Achilles tendon. Angle the needle forward so that it strikes the tibia and does not pass between it and the Achilles tendon. Before injecting withdraw the plunger to ensure the needle is not in one of the posterior tibial vessels. Wait for ten minutes. (If this is being done in a general practitioner's surgery the procedure so far can be done in a side room while instruments are set to boil and another patient is being seen.)

The sole of the foot is tested for anaesthesia. The skin is cleaned with a suitable agent. The wart is curetted with a small spoon while the thumb and index finger of the free hand exert firm pinching pressure on either side of the cavity to prevent bleeding. This pressure is maintained while the margin of the cavity is trimmed with scissors and a silver nitrate stick is applied. Pressure can then be relaxed and bleeding should not ensue. A dry gauze dressing secured by adhesive plaster should be kept on for five days.

The sole of the foot may remain numb for about one hour and patients must be warned against stumbling and car drivers should not be allowed to drive in this condition. It is obvious that both feet should not receive this treatment at the same session.

This nerve block is also very useful in the treatment of lacerations of the foot or removal of foreign bodies. It is markedly less painful than local infiltration.—I am, etc.,

W. G. R. M. LAURIE

Oxford

SIR,—I read with interest your leading article on plantar warts (24 June, p. 725). However, I noticed that you omitted cautery as a means of treating these lesions. Cautery has the advantage of causing little or no bleeding and I believe that the burning process has a better chance of ensuring that the virus particles are all destroyed.—I am, etc.,

STUART SANDERS

London W.1

Stress and the Schizophrenias

SIR,—I am rather surprised that your leading article (3 June, p. 550) should regard severe stress and a favourable outcome in schizophrenias as paradoxical. It surely indicates that it needs more stress to precipitate the illness in those of better personality than their weaker brothers. I have always considered in psychiatry in particular that environment $\times$ constitution = a constant. In other words a slight wind will topple the weak, a strong wind is needed to knock down the strong.—I am, etc.,

W. LINDSAY NEUSTATTER

London W.1