This case is reported to draw attention to this false-positive phentolamine test which might be found in other patients being investigated for hypertension while taking ethambutol.

I am indebted to Mr. Howard G. Hanley for permission to report this case. The plasma catecholamines were estimated by Dr. Malcolm Carruthers, of the Institute of Ophthalmology

References

- Litchfield, J. W., and Peart, W. S. (1956). *Lancet*, 2, 1283.
 Place, V. A., and Thomas, J. P. (1963). *American Review of Respiratory Diseases*, 87, 901.
 Sheps, S. G., Tyce, G. M., Flock, E. V., and Maher, F. T. (1966). *Circulation*, 34, 473.
 Strauss, I., and Erhardt, F. (1970). *Chemotherapy*, 15, 148.

Unusual Case of Paratyphoid Fever

N. TODD

British Medical Journal, 1972, 3, 333

Salmonella paratyphi C is not normally isolated in Canada or the U.K. The present report concerns a case of paratyphoid that occurred and was treated in British Columbia. Investigations failed to trace the source of infection.

Case Report

A 14-year-old white schoolboy with no previous illnesses of any note became ill with a fever which lasted about two weeks. He had fainted once during the course of this time but had not had any abdominal pain, vomiting, or diarrhoea. There were no similar cases in the area at the time.

Examination showed no abnormalities except a temperature of 39.4° C and a sinus tachycardia of 110/min. He was admitted to hospital and the following investigations were done: chest x-ray picture negative; urine, routine analysis and culture negative six times; E.C.G. normal on each of three times performed; anti-streptolysin O titre 50 units; throat swab negative; Paul-Bunnell test negative twice; haemoglobin 14 g/100 ml. Blood measurements for days 1, 6, and 8 are given in the Table.

Haematological Picture during Treatment

Day	W.B.C. (/mm ³)	Poly- morphs	Stab Cells	Lympho- cytes	Mono- cytes	E.S.R. (mm/hr)
1	8,000	54%	13%	29%	4%	35
6	39,000	27%	49%	19%	5%	42
8	7,000	44%	3%	48%		104

PROGRESS

While in hospital his temperature fluctuated between 38° and 40°C until he was started on intravenous medication. Blood cultures were taken immediately on admission but grew nothing for the first few days.

Day 3.—He developed a discharge in the posterior nasopharynx which was somewhat haemorrhagic. It was not readily moved by swabbing. Sinus x-ray films were negative. A culture from the posterior nasal discharge grew *Diplococcus pneumoniae*.

Day 4.—His posterior nasal discharge was still present and he had a number of small shallow ulcers on the nasal septum. He also developed two large scrotal infarcts, one about 1 in (2.5 cm) long and the other about ½ in (1.3 cm). The skin became rapidly dark and black and then sloughed in these areas. At that time the question of diphtheria arose and it was noted that he had been immunized against diphtheria, but consultation with an E.N.T.

specialist and an internal medicine physician was obtained largely to try to exclude this seemingly remote possibility. Both physicians felt confident that this was not a case of diphtheria although the diagnosis was obscure at the time. The possibility of infectious mononucleosis was again raised and consideration of Wegener's granuloma, septicæmia possibly due to pseudomonas, and also a vasculitis due to periarteritis nodosa were discussed.

Day 6.—By this time it seemed evident that some sort of septicæmia was involved and intravenous ampicillin 500 mg and cloxacillin 500 mg every four hours was instituted, since the organism was unknown.

Day 9.—There was noticeable clinical improvement with a fall in the fever and the mucosal and skin lesions started to heal. On this day the blood culture taken from day 1 was read as positive and a probable salmonella organism was identified.

Day 10.—The cloxacillin was stopped and ampicillin continued.

Day 17.—The organism was identified as *Salm. paratyphi* C (1). He was clinically a great deal better and the intravenous ampicillin was changed to 500 mg every six hours by mouth.

Day 24.—Stool cultures grew the same *Salm. paratyphi* C (1) three times. He was feeling well but developed a new lesion in the mouth. It appeared to be a simple vesicle but was approximately 4 mm on its longest side. Since there had been earlier oral mucosal lesions and it was difficult to be certain about the cure of this condition it was elected to discontinue the ampicillin and he was started on co-trimoxazole in a dose of two tablets twice a day. This had just become available in the hospital on an experimental basis.

Day 26.—It was obvious that this was a simple vesicle in the mouth and he was discharged to continue with the co-trimoxazole for two weeks at home.

He was seen in the course of the next two weeks as an out-patient and made a smooth recovery with healing of his scrotal lesions. Three stools were tested at about six weeks after his discharge and they were reported as negative.

Comment

Systemic salmonella infections are increasing in incidence. Focal lesions have been described in numerous organs and tissues in such cases but I have seen no record of involvement such as we have had in this case with lesions of the nasopharynx and nasal septum and actual infarcts of the skin of the scrotum. The importance of early blood cultures in such cases is obvious and the long period of incubation (nine days) of the paratyphoid organism is noteworthy. The way in which these lesions are produced is not clear, but since they seemed to start about two weeks after the onset of the illness it raises the possibility of circulating antigen-antibody complexes being responsible.

I wish to thank Dr. D. Klassen and Dr. R. W. Van der Flier for their help with this case.

Reference

- Christie, A. B., *British Journal of Hospital Medicine*, 1971, 5, No. 5.

Chilliwack General Hospital, Chilliwack, B.C., Canada
 N. TODD, B.Sc., M.B., Active Staff, General Practitioner