kept constantly under review. Time must be set aside for regular consultation with the patient, his family, and their general practitioner. A psychiatrist who becomes well known to the staff should be available so that he may contribute in his now traditional role as leader of the team concerned with mental health.—We are, etc.,

R. C. B. AITKEN
C. G. BARROW

University Department of Psychiatry,
Royal Edinburgh Hospital,
Edinburgh

¹ Report of the Joint Committee on Maintenance Dialysis and Transplantation in the Treatment of Chronic Renal Failure. London. Royal College of Physicians, 1972.

Pulmonary Oedema of Mountains

SIR,—Your leading article, "Pulmonary Oedema of Mountains" (8 July, p. 65), carries the recommendation that "if qualified medical help is available, it seems generally agreed that morphine as well as frusemide should be given . . ." May I introduce a caveat?

In 1961 our case ¹ went into the mountains for a second time following a first attack of acute pulmonary oedema of altitude. He was given a syrette containing 10 mg morphine sulphate with the advice that "this may do you some good if you need it." At 11,500 ft (3,444 m) he developed nausea, vomiting, cough, shortness of breath, and pulmonary rales. After he had walked as far as he could out of the mountains he sat down to rest at about 7,500 ft (2,280 m). Following failure to improve on several hours' rest propped against a tree the morphine sulphate 10 mg was given intravenously. There was an immediate and profound cyanosis, noticeable to the patient (a physician) as well as to those around him and a subjective lowering in his expectation of survival, rather than a "relief of anxiety" (Singh's phrase²).

Unless this was a unique experience it might be wise for physicians in the mountains to avoid giving morphine to patients with acute pulmonary oedema of altitude unless they are prepared to give frusemide and oxygen, as Singh generally did.³—I am, etc.,

TALCOTT BATES

Monterey,
California, U.S.A.

- ¹ Fred, H. L., Schmidt, A. M., Bates, T., and Hecht, H., *Circulation*, 1962, 25, 929.
² Singh, I., et al., *New England Journal of Medicine*, 1969, 280, 175.
³ Singh, I., Kapila, C. C., Khanna, P. K., Nanda, R. B., and Rao, B. D. P., *Lancet*, 1, 229.

E.C.T. and CAMP Excretion

SIR,—I should like to support albeit in a preliminary way the findings of Dr. K. Hamadah and others (19 August, p. 439). In May 1971 we decided to run a small pilot investigation on the effect of electric convulsion therapy on urinary levels of 3'5' cyclic adenosine monophosphate. Four men suffering from endogenous depression were selected. Their ages ranged from 45 to 62. All were short-term inpatients. Cyclic AMP levels were measured by the saturation analysis method described by Brown and colleagues.¹

In the first experiment urine samples were

collected one hour before and two hours after E.C.T. was administered. In the second experiment 12-hour specimens of urine were collected before and after E.C.T. The levels of cyclic AMP in the first experiment ranged from 0.9 to 2.1 $\mu\text{mol}/24\text{ hr}$ before E.C.T. and there was no apparent change in the levels after E.C.T. However, when 12-hour specimens were analysed in three patients the following results were obtained:

	Before E.C.T.	After E.C.T.
Patient A	2.0-2.3 $\mu\text{mol}/24\text{ hr}$	9.2-9.3 $\mu\text{mol}/24\text{ hr}$
Patient B	0.5-0.8 $\mu\text{mol}/24\text{ hr}$	3.5-3.8 $\mu\text{mol}/24\text{ hr}$
Patient C	2.9-3.3 $\mu\text{mol}/24\text{ hr}$	3.1-3.3 $\mu\text{mol}/24\text{ hr}$

There appeared to be a rise in some patients after E.C.T., and therefore the investigation was thought worth pursuing under more rigorously controlled conditions. An application was therefore made for support for such a study to the Medical Research Council and a grant for technical assistance has been forthcoming. The work is to be undertaken in Hull University Department of Biochemistry in association with Broadgate Hospital, Walkington.—I am, etc.,

ISABEL C. A. MOYES

Beverley, Yorks

¹ Brown, B. L., Albano, J. D. M., Ekins, R. P., Sgerzi, A. M., and Tampion, W. *Biochemical Journal*, 1971, 121, 561.

Rosette Inhibition Test in Management of Transplantations

SIR,—The report from Mr. M. Bewick and others (26 August, p. 491) is of great interest as it describes a serious attempt to prescribe immunosuppressive drugs on a rational basis. It is unfortunate, however, that there has been no statistical analysis which might allow a better appreciation of the results. A major disappointment, as the authors point out, was the large number of patients that died (28%).

Immunosuppressive drugs obviously remain as dangerous as ever and the giving of high doses needs careful consideration. Until safer methods of immunosuppression are available it is probably unjustified (as the authors have intimated) to prescribe on the results of this test alone.—I am, etc.,

J. R. SALAMAN

Cardiff Royal Infirmary,
Cardiff

Oral Contraceptives and Serum Amylase

SIR,—We were interested in the report by Dr. H. Adlercreutz and others (26 August, p. 529) of elevated serum α -amylase values in women taking oral contraceptives.

We reported a case¹ with high serum amylase and clinical features suggestive of pancreatitis in a young woman who had been taking oral contraceptives for two years. The acute abdominal pain, vomiting, and epigastric tenderness developed two weeks after a myocardial infarction. Other causes of acute abdominal pain associated with elevated serum amylase such as cholecystitis, perforated peptic ulcer, and intestinal obstruction were excluded. Serial electrocardiograms and enzyme studies did not show fresh myocardial damage. Though the epigastric tenderness subsided in four days the serum amylase remained raised and was 291 Somogyi units per 100 ml six weeks after the attack of abdominal pain.

We have been unable to find reference in the literature to the occurrence of pancreatitis in women taking oral contraceptives. It would be of interest to know if there was any relationship between the oestrogen content of the oral contraceptives and the raised serum amylase levels in the cases investigated by Dr. Adlercreutz and his colleagues.—We are, etc.,

D. P. ATUKORALE
N. J. WALLOPPILLAI

General Hospital,
Colombo,
Sri Lanka

¹ Atukorale, D. P., and Walloppillai, N. J., *Ceylon Medical Journal*, 1971, 16, 171.

Contraception and Infertility

SIR,—I am sure that all practitioners who utilize their own retrospectroscope with a balance between self-criticism and self-confidence have more than one instance such as Brenda's problem so carefully documented by Dr. K. L. Oldershaw and Mr. Michael Brudenell in "Second Opinion, Please" (9 September, p. 637), where they are not satisfied that they have done as well as they might.

What does puzzle me is that Dr. Oldershaw should criticize himself by writing "My prescription of oral contraceptives for three years would appear to be a good example of a doctor providing bad treatment with the best of motives." Even with the benefit of the findings of this wonderful instrument I fail to see why my colleague accuses himself of providing bad treatment. Nowhere in his discussion is one led to believe that his treatment was bad, and, perhaps in my ignorance, I cannot see that it was. It might have been inappropriate to prescribe contraceptives for a sterile patient, but bad treatment for an unknowable complaint—surely not. My compliments to Dr. Oldershaw on a nicely-presented case history, but he doesn't deserve his own criticism.—I am, etc.,

P. C. ARNOLD

Sydney,
N.S.W., Australia

SIR,—Under the heading "Second Opinion, Please" (9 September, p. 637) Dr. K. L. Oldershaw and Mr. Michael Brunenell gave an excellent detailed account of an unfortunate patient who is sterile and likely to remain so. Dr. Oldershaw states: "My prescription of oral contraceptives for three years would appear to be a good example of a doctor providing bad treatment with the best of motives." It would be nice if he would explain why providing good contraceptive advice and treatment for a newly-married girl who subsequently turns out to be sterile is bad treatment.—I am, etc.,

MICHAEL SMITH

Family Planning Association,
London W.1

Sexual Permissiveness

SIR,—Dr. Violet Anderson's "Personal View" (26 August, p. 524) undoubtedly expressed many of the unspoken thoughts and fears of many doctors. The case described by Dr. K. L. Oldershaw and Mr. Michael Brudenell (9 September p. 637) is surely a perfect example of the final fear of distant