CORRESPONDENCE

Correspondents are asked to be brief

Severely Malformed Children J. Lorber, F.R.C.P. 46	Hepatitis-associated Antigen in Chronic Hepatitis C. Del Vecchio-Blanco, and others 49	Mini-clinics in General Practice P. H. Brunyate, M.B. 51
Antibiotic-sensitivity Tests R. Blowers, F.R.C.P., and others 46	Androgens and Exercise M. H. Briggs, D.Sc., and others 49	Coeliac Disease and Schizophrenia F. C. Dohan, M.D. 51
Ampicillin for Sore Throat H. Pullen, M.R.C.P.E.D.; D. C. Turk, F.R.C.PATH. 47	Isocycline in Enteric Fever and Brucellosis A. Z. Shafei, M.R.C.P., and W. S. Luka 50	Haemolytic-Uraemic Syndrome with Pericarditis and Pleurisy M. H. Winterborn, M.R.C.P., and others 52
Drugs for Angina Pectoris G. Nyberg, M.D. 47	Psychosurgery P. K. Bridges, M.D., and J. R. Bartlett, F.R.C.S. 50	Congenital Hemihypertrophy G. R. Clarke, M.R.C.P. 52
Potassium Deficiency during Treatment with Brinaldix K R. T. Taylor, M.R.C.P., and M. J. T. Peaston, M.R.C.P. 48	American Medicine I. L. Gregory, M.D. 50	Antibiotic Sensitivity of Klebsiella Eunice Lockey, F.R.C.PATH., and M. W. Casewell, M.B. 52
Fenfluramine and Growth Hormone Release D. L. F. Dunleavy, M.B., D.P.M. and others 48	Haemophilus influenzae Apparently Resistant to Trimethoprim S. R. M. Bushby, PH.D. 50	Epidemiology of Simple Hypospadias H. Campbell, M.B., F.F.C.M., and others 52
Transmission of Trichuris trichiura Elizabeth A. Green, M.B., D.P.M., and others 48	Oestrogen Treatment in Carcinoma of the Prostate R. J. Shearer, F.R.C.S., and others 51	Better Aids for the Deaf H. Hillman, PH.D., M.B. 53
Gram-negative Bacilli on Hands S. Selwyn, M.D. 49	Communication Between Psychiatrists E. A. Burkitt, M.R.C.PSYCH. 51	Hospital Social Workers C. C. Slack, F.R.C.S.ED. 53
		Payment for Contraceptive Services L. M. Ellis, M.B. 53
		Representation of Hospital Doctors Pamela Wray, M.D. 53

Severely Malformed Children

SIR,—It is necessary to comment on Mr. R. B. Zachary's interpretation (26 May, p. 482) of the tape recorded discussion on severely malformed infants (5 May, p. 284). There is no "strange anomaly," as Mr. Zachary asserts, in the fact that though untreated babies with spina bifida in first-class centres have all died within a few months, surgeons and paediatricians "are seeing a number of untreated cases for salvage from other hospitals." The vital omission from Mr. Zachary's comment is that the "100% success" rate of no-treatment policy relates to those infants selected for no treatment because of the severity of their condition and not to all babies born with spina bifida. Since I advocated certain definite and strict criteria for selection¹ this policy has been largely accepted in Britain and in most other countries.

Between May 1971 and January 1973, 38 newborn infants have been referred to me, and of these 25 were not treated. All died within eight months (18 before three months of age); 12 of the 13 treated are alive and only one is badly handicapped. We were certainly and rightly horrified to see a few untreated older children who had gross and unnecessarily gross handicaps. These were usually those whose condition at birth was more favourable and who should have been treated, with a good chance of an acceptable quality of life. It is my conviction that all infants born with spina bifida should immediately be seen by an expert in this field, who, on the basis of his experience and with full explanation to the parents, should make the recommendation to treat or not. Occasional mistakes will be made, but if the initial assessment is correct, then the rare survivor in the non-treated group will not suffer if subsequently he is given the "total care" which those treated from the first day should normally receive.

It is very unfair to point to the occasional tragedy of a severely affected untreated infant without mentioning that for every one of these we have some 40 just as severely

handicapped, with mental retardation, gross paralysis, incontinence, kyphos, and other handicaps, in spite of all the operations and other forms of therapy they had from a few hours of age onwards, from the days when we offered treatment to all infants irrespective of their condition at birth.²—I am, etc.,

JOHN LORBER

Children's Hospital,
Sheffield

¹ Lorber, J., *Developmental Medicine and Child Neurology*, 1971, 13, 279.

² Lorber, J., *Archives of Disease in Childhood*, 1972, 47, 854.

Antibiotic-sensitivity Tests

SIR,—During trials by the Public Health Laboratory Service Quality-control Committee, which includes some members from N.H.S. hospital laboratories, the results of antibiotic-sensitivity tests have been studied in relation to the methods used. These studies are continuing, but directors of laboratories who cannot at present be accommodated in the scheme may wish to know of two significant results that have so far emerged.

Cephalosporin-sensitivity Test. A strain of *Staphylococcus aureus*, resistant to methicillin, was tested by disc diffusion methods for sensitivity to cephaloridine in 20 laboratories. Ten laboratories reported it sensitive and 10 reported it resistant, confirming the difficulty

of testing sensitivity of staphylococci to cephalosporins by the disc method. Disc diffusion tests of cephalosporins against staphylococci may result in misleading reports being sent to clinicians, and the committee recommends that such tests should not be used for determining the sensitivity of staphylococci to this group of drugs. In practice it can be assumed that all methicillin-resistant strains of *Staph. aureus* are resistant to cephalosporins and that all methicillin-sensitive strains are sensitive to cephalosporins.

Ampicillin Content of Discs. High-content discs are often used for tests of urinary pathogens because high concentrations of ampicillin can be attained in urine. At present we have no comments to make on the validity of this practice. But when high-content ampicillin discs are used for testing pathogens from sites other than the urinary tract misleading reports may be issued, as shown by the following results from recent trials:

There is no statistically significant difference between our results with 2- μ g and 10- μ g discs for the *Bacteroides* nor between any of the discs for the *Proteus*, but 10 μ g is probably the best compromise for use with most organisms from sites other than the urinary tract. When a 25- μ g disc is used, diminished zone size indicating resistance may be recognized if it is compared with the zone of a fully sensitive control organism but the difference in zone size is likely to be more obvious with a 10- μ g disc. The committee recommends, therefore, that discs containing 10- μ g of ampicillin should be used

Ampicillin content of disc	Resistant <i>Bacteroides</i> Spp. (M.I.C. = 32 μ g/ml)			Sensitive <i>Proteus mirabilis</i> *		
	No. of reports	Errors		No. of reports	Errors	
		No.	%		No.	%
2 μ g	23	1	4	50	7	14
10 μ g	14	1	7	36	3	8
25 μ g	23	7	30†	57	1	2

*tested twice by most laboratories † P < 0.05

for pathogens isolated from sites other than the urinary tracts.—We are, etc.,

ROBERT BLOWERS

Chairman,
P.H.L.S. Quality-control Committee
Clinical Research Centre,
Harrow, Middlesex

E. JOAN STOKES

Department of Clinical Pathology,
University College Hospital,
London W.C.1

J. D. ABBOTT

Secretary,
P.H.L.S. Quality-control Committee
Public Health Laboratory,
Withington Hospital,
Manchester

Ampicillin for Sore Throat

SIR,—Dr. A. A. Robertson's conclusion (9 June, p. 614) that the use of ampicillin must be considered in the treatment of acute sore throats is based on the following premises: (1)—that *Haemophilus influenzae* is frequently isolated from the throats of patients with acute sore throats; (2)—that where this organism is isolated it is responsible for the acute sore throat; and (3)—that ampicillin is the antibiotic of choice in *H. influenzae* infections.

The first premise is true, but it is also true that *H. influenzae* is frequently isolated from the throats of normal people,^{1,2} so that the truth of the second premise is to be questioned. With regard to the third premise it is true that on average ampicillin is about twice as active as penicillin against *H. influenzae*³ and this "resistance" of *H. influenzae* to penicillin and "sensitivity" to ampicillin is frequently stressed (especially in advertisements for ampicillin). However, it should be remembered that many strains of *H. influenzae* are as sensitive to penicillin as to ampicillin, and most *H. influenzae* will have their growth at least inhibited by levels of penicillin which should readily be achieved by correctly given oral phenoxymethylpenicillin.

In life-threatening infections with *H. influenzae* such as acute epiglottitis, septicaemia, and meningitis, it would, of course, be imperative to give the most active antibiotic, and ampicillin would be more appropriate than penicillin in these situations. However, in relatively trivial infections such as acute sore throats, which will almost invariably settle spontaneously within a few days (unless due to infectious mononucleosis) other factors, such as the cost of the drug and the risk of adverse factors such as monilial superinfection and drug rash, have to be taken into account.

If one considers all the factors mentioned above, it is doubtful if ampicillin is ever indicated in the treatment of acute sore throat, even if *H. influenzae* has been isolated. When one considers also the predominance of haemolytic streptococci in bacterial sore throats, surely the routine use of ampicillin in preference to penicillin in acute sore throats is never justified.—I am, etc.,

H. PULLEN

Department of Infectious Diseases,
Seacroft Hospital,
Leeds

¹ Wilson, G. S., and Miles, A. A., eds., *Topley and Wilson's Principles of Bacteriology and Immunity*, 5th edn., vol. 2, p. 2473. London, Arnold, 1964.

² Turk, D. C., and May, J. R., *Haemophilus influenzae: Its Clinical Importance*, p. 13. London, English Universities Press, 1967.

³ Ibid, p. 73.

SIR,—Dr. A. A. Robertson (9 June, p. 164) states that "penicillin was not the antibiotic of choice in over one-third" of 143 patients with sore throat. He apparently bases this statement, and his preference for ampicillin as the drug for treating sore throats, on the fact that 49 of the patients "yielded swabs positive for *Haemophilus* species." Failure to find a penicillin-sensitive potential pathogen may constitute an adequate reason for regarding penicillin as inappropriate; but the findings of *Haemophilus* species in the throats of one-third of the patients is irrelevant to the issue. It is not uncommon to find *H. influenzae* in the throats of 50% or more of the members of a healthy population.¹ I know of no evidence that the ordinary non-capsulated *H. influenzae* strains of the upper respiratory tract cause sore throats (though capsulated type b strains can occasionally cause tonsillitis—and also, of course, epiglottitis, which is hardly to be included under the term "sore throat").² There is some circumstantial evidence that haemolytic haemophili (which are mostly of the species *H. parahaemolyticus*) may cause pharyngitis^{3,4} but this is by no means proved.—I am, etc.,

D. C. TURK

Department of Bacteriology,
Radcliffe Infirmary,
Oxford

¹ Turk, D. C., and May, J. R., *Haemophilus influenzae: Its Clinical Importance*, pp. 13-23. London, English Universities Press, 1967.

² Addy, M. G., Ellis, P. D. M., and Turk, D. C., *British Medical Journal*, 1972, **1**, 40.

³ Salgado Correia, M. J., and Torres Pereira, A., *Arquivos do Instituto Bacteriológico Câmara Pestana*, 1966, **11**, 87.

⁴ Branson, D., *Applied Microbiology*, 1968, **16**, 256.

Drugs for Angina Pectoris

SIR,—Dr. Brian Livesley and others (17 February, p. 375) recently reported a five-drug trial in angina pectoris, comparing verapamil in two dose levels, propranolol, and isosorbide dinitrate. I find it pertinent to make a few methodological observations on this study.

The order of treatment was always the same, though different patients started at different parts of the drug cycle. This meant that drug E, which was isosorbide dinitrate, in 80% of the patients came after at least one other drug. Consequently, all possible carry-over effects of this drug longer than one month will be reflected in the results ascribed to isosorbide dinitrate. The same consequence is true for all the other drugs. A complete randomization of the order of these five treatments would have obviated this difficulty. The authors state that only the last two weeks of each four-week period were analysed in order to avoid carry-over effects. They have missed the point, however, of analysing whether there were any carry-over effects at all. If the analysis of all four weeks yields the same results as for only the last two weeks there are either carry-over effects longer than four weeks or none at all.

The authors give significance limits for differences between various treatments using the non-parametric Wilcoxon signed ranks test, which no doubt is a relevant test since nothing is known about the distribution of data. However, no average or median values are given to show how large these differences

were. Moreover, six comparisons are presented. Dunnnett¹ thinks that if repeated paired testing is performed in a multi-treatment system like the one reported not more than $N-1$ comparisons are allowed, where N = no. of treatments, and furthermore, that one of the treatments is not to be used more than twice. If all treatments are to be compared an analysis of variance should be performed. Assuming that the recorded variables are not normally distributed, an assumption with which I would agree, the type of test that should be used is the Friedman two-way analysis of variance.² This involves ranking of the different treatments for each patient. In doing so, propranolol comes out with the lowest rank sum (= best results) in the 16 patients having all five treatments. The control or placebo values come out with the highest rank sum. However, a Friedman analysis of variance does not yield any statistical significance—that is, the results of these five treatments differ with a confidence limit higher than 5%. If the average values of the work done for these 16 patients are calculated it will be seen that verapamil 120 mg three times daily increases the work (compared with the average control value) by about 33%, verapamil 80 mg three times daily by about 14%, propranolol 100 mg three times daily by about 21%, and isosorbide dinitrate 20 mg three times daily by about 49%. It is readily seen that the high value for isosorbide dinitrate is due to one patient having a tremendous increase of work on this drug, and the results just reported bear out another point worth mentioning: the characterization of a sample which is not normally distributed is better done by the median rather than the mean. The median values of the work done (in kg/sec) for these 16 patients are respectively (approximately) 9,500 for control, 12,500 for verapamil 120 (+32% compared with the median control value), 9,500 for verapamil 80 ($\pm 0\%$), 13,000 for propranolol (+37%), and 11,000 for isosorbide dinitrate (+21%).

Maybe the best and most relevant way of expressing the results is to state which drug was the best for each patient. It is then seen in the 16 patients mentioned that verapamil 120 was best in four patients, verapamil 80 in three patients, and verapamil 120 and 80 was equally good and the best drug in one patient. Propranolol was the best drug in one patient and isosorbide dinitrate in one patient. In one of the patients best on verapamil 120 the result was equal with isosorbide dinitrate. Six patients showed practically no difference between any of the drugs. All these values are with reference to the control values. This analysis would indicate that in the 16 patients going through all five treatments the best drug was in the majority of cases verapamil, whereas propranolol and isosorbide dinitrate were equal in terms of "best drug." On the average, however, there was no significant difference between any of the drugs in terms of increased work done, and the results do not point to very important differences between the drugs. At all events, it seems unwarranted to state specifically for isosorbide dinitrate that it is as ineffective as placebo.

It seems as if the original proprietary tablets were not used in this trial. Nevertheless, the authors state in the summary that Cordilox, Inderal, and Vascardin were used. A clari-