- Levin, J., and Zak, B. (1964). *Clinica Chimica Acta*, **10**, 381.
- Mosbach, E. H. (1972). In *Bile Acids in Human Diseases*, ed. P. Back and W. Gerok, p. 89. Stuttgart, Schattauer.
- Nilsson, S., and Schersten, T. (1970). *European Journal of Clinical Investigation*, **1**, 109.
- Piper, J., and Orrid, L. (1956). *American Journal of Medicine*, **21**, 34.
- Press, M., Miller, P., Shimoyama, T., Kikuchi, H., and Thompson, G. R. (1972). *European Journal of Clinical Investigation*, **2**, 301.
- Sarles, H., Chalvet, M., Ambrosi, D., and D'Ortoli, G. (1957). *Semaine des Hôpitaux de Paris*, **33**, 3424.
- Sarles, H., et al. (1969). *American Journal of Digestive Diseases*, **14**, 531.
- Scherstén, T., Alling, C., Cahlin, E., Jonssen, J., and Nilsson, S. (1972). Abstract of 9th International Congress on Gastroenterology.

- Seitz, W., and Brand, V. (1961). *Klinische Wochenschrift*, **39**, 891.
- Senior, J. R. (1969). In *Bile Salt Metabolism*, ed. E. Schiff, J. B. Carey, jun., and J. M. Dietschy, p. 56. Springfield, Illinois, Thomas.
- Small, D. M. (1971). *New England Journal of Medicine*, **284**, 214.
- Thistle, J. L., and Schoenfield, L. J. (1971). *New England Journal of Medicine*, **284**, 177.
- Thorbjarnson, B. (1969). *Archives of Surgery*, **98**, 372.
- van der Linden, W., and Norman, A. (1967). *Acta Chirurgica Scandinavica*, **133**, 307.
- Vlahcevic, Z. R., Bell, C. C., jun., Buhac, I., Farrar, J. T., and Swell, L. (1970). *Gastroenterology*, **59**, 165.
- Wheeler, M., Hill, S. L. L., and Laby, B. (1970). *Gut*, **11**, 430.
- Wilson, J. D. (1972). *Archives of Internal Medicine*, **130**, 493.

PRELIMINARY COMMUNICATIONS

Some Actions of Growth Hormone Release Inhibiting Factor

Aa. PRANGE HANSEN, H. ØRSKOV,
K. SEYER-HANSEN K. LUNDBÆK,

British Medical Journal, 1973, **3**, 523-524

Summary

Preliminary studies indicate that somatostatin (SRIF, somatostatin release inhibiting factor) can suppress exercise-induced plasma growth hormone rise in normal subjects as well as in diabetics.

Introduction

Three years ago we proposed the hypothesis that growth hormone overproduction is a casual factor in the development of diabetic angiopathy. It is a working hypothesis based on the facts that the plasma growth hormone level is very high and fluctuates throughout life in diabetic patients and that surgical removal of the source of growth hormone production inhibits the further development of vascular changes in severe long-term diabetic angiopathy (Lundbæk *et al.*, 1970; Hansen, 1972).

The practical conclusion of this working hypothesis is that attempts at growth hormone suppression by pharmacological means in diabetic patients should be pursued, particularly since hypophysectomy is not an ideal procedure for many reasons. The exercise test developed in our laboratory (Hansen, 1972) has been used in this search. Exercise is a normally occurring stimulus to growth hormone secretion in daily life. In a study with a practical therapeutic aim this test is to be preferred to artificial stimuli such as insulin-induced hypoglycaemia or the administration of arginine, levodopa, or other compounds.

Using the exercise test we have assessed many commercially available drugs acting in the hypothalamic area, especially various psychotropic drugs, on the assumption that one or more of these compounds might influence the assumed

growth hormone releasing factor in the hypothalamus. Only amitriptyline seems to have any effect, but it is much too slight. Alpha-adrenergic blocking agents suppress growth hormone during exercise in diabetic patients (Hansen, 1971) but only when given intravenously.

Recently a naturally occurring hypothalamic growth hormone release inhibiting factor—somatostatin or SRIF (somatostatin release inhibiting factor)—has been found and isolated as a tetradecapeptide by Guillemin and his co-workers (Brazeau *et al.*, 1973). Preliminary results obtained with this compound in seven normal subjects and in nine diabetics are presented here. The non-diabetics were all doctors from the department. The diabetics and the controls were all normal weight, men, age range 25-39 years. As little is known about possible effects of somatostatin the drug was tested on several of the controls before it was given to the diabetic patients. The only side effects observed were a slight feeling of oppression in the chest of some normal and diabetic subjects immediately after the injection of somatostatin and in all cases a rise in blood pressure of 10-15 mm Hg accompanied by a fall in pulse rate of 10-20 beats/min lasting for less than five minutes. All the subjects were fully informed of the experimental nature of the investigation and of the possible side effects. Only patients who had had diabetes for less than five years and had normal blood pressure were investigated.

Methods and Results

The exercise test consisted in a control period, an exercise period of 450 kg/min for 40 minutes on a bicycle ergometer, and a post-exercise period. It was done once with somatostatin and one without.

In most experiments somatostatin was given as a rapid intravenous injection (two minutes) of 250 µg 10 minutes before beginning exercise, followed by a continuous infusion of 250 µg for the next 60 minutes. In four experiments—all on non-diabetics—somatostatin was given in six short injections every 10 minutes from 10 minutes before to 40 minutes after starting exercise. The results were not appreciably different in these two experiments.

Plasma growth hormone was determined with a single antibody radioimmunoassay employing wick-chromatography (Ørskov *et al.*, 1968). A Wilhelmi preparation (HS 1216 C) was used for growth hormone standards.

The average curve for the seven non-diabetics is shown in fig. 1. The average rise was lower when somatostatin was given. The suppression is statistically significant when using the paired *t* test at the beginning of the work period; after 20, 30, and 40 minutes of work; and 10, 50, and 60 minutes after cessation of work (2 P < 0.05).

Second University Clinic of Internal Medicine, Kommunehospitalet, Aarhus, Denmark

Aa. PRANGE HANSEN, M.D., Senior Registrar in Medicine
H. ØRSKOV, M.D., Director, Laboratory of Experimental Medicine
K. SEYER-HANSEN, M.D., Reader in Medicine
K. LUNDBÆK, M.D., Professor of Medicine

FIG. 1—Exercise-induced growth hormone (GH) curve produced by seven normal subjects with and without somatostatin.

The results for the nine male diabetics are shown in fig. 2. Somatostatin also caused suppression in this group. This effect was statistically significant 20 and 30 minutes after beginning exercise (2 $P < 0.05$ and < 0.025 respectively). The experiments without somatostatin show that fasting growth hormone values were higher and the rise during exercise occurred earlier and was more pronounced in the diabetics. This is in accordance with earlier findings (Hansen, 1972).

It should be emphasized that the metabolic state of the diabetics was similar in the experiments with and without somatostatin (average fasting blood glucose 138 mg/100 ml in control experiments and 130 mg/100 ml in the somatostatin experiments).

Before therapeutic experiments with somatostatin are feasible it is necessary to develop a mode of long-term suppression. One possibility is to produce a delayed absorption compound. Another is to try somatostatin or some modification of the molecule given by mouth. If a preparation which can suppress the growth hormone hypersecretion of diabetics in a clinically acceptable way is found it should be used in con-

FIG. 2—Exercise-induced growth hormone curve produced by nine male diabetics with and without somatostatin.

trolled clinical trials to observe its effect on the development of diabetic angiopathy.

The somatostatin used in this study was prepared and characterized as described by Sarantakis and McKinley (1973).

References

- Brazeau, P., *et al.* (1973). *Science*, **179**, 77.
- Hansen, Aa. P. (1971). *Journal of Clinical Endocrinology and Metabolism*, **33**, 807.
- Hansen, Aa. P. (1972). *Danish Medical Bulletin*, Suppl. No. 1.
- Lundbaek, K., *et al.* (1970). *Lancet*, **2**, 131.
- Ørskov, H., *et al.* (1968). *Nature*, **219**, 193.
- Sarantakis and McKinley (1973). *Biochemical and Biophysical Research Communication*. In press.

MEDICAL MEMORANDA

Acute Pancreatitis in Coxsackie B Infection

BO URSING

British Medical Journal, 1973, **3**, 524-525

The aetiology and pathogenesis of acute pancreatitis still challenge research. Gall stones and the abuse of alcohol are thought to be common casual factors (Edlund, 1970; Lundh, 1970). Infectious agents are rarely responsible (Everett, 1969; Kunz, 1971), though mumps is a well known cause, sometimes leading

to diabetes mellitus (Melin and Ursing, 1958). Pappenheimer *et al.* (1951) showed that injection of material from a patient with meningoencephalitis and pleurodynia damages the pancreas in mice. In generalized Coxsackie virus infection in infants virus and lesions are found in the pancreas, liver, myocardium, and meninges (Kibrick and Benirschke, 1958; Fechner *et al.*, 1963). Reports of Coxsackie virus as a probable cause of acute pancreatitis in adults are rare (Murphy and Simmul, 1964; Nakao *et al.*, 1964).

In 1972 acute pancreatitis was diagnosed in two adults with Coxsackie B5 infection. Their cases are reported here.

Case 1

A woman aged 27 who had been previously healthy and had no history of gall bladder or liver disease fell ill on 4 September with pyrexia, arthralgia, and pain in the right chest. On 9 September she had exanthema and severe abdominal pain and was admitted to hospital. During the following week she developed a fulminant picture of acute pancreatitis, with high-grade fever, hypotension, and tachycardia. She vomited and developed paralytic ileus of seven days'

University Hospital, S-221 85 Lund, Sweden

BO URSING, M.D., Assistant Professor, Department of Infectious Diseases