- ¹ Kannel, W. B., Castelli, W. P., Gordon, T., and McNamara, P. M., *Annals of Internal Medicine*, 1971, 74, 1.
- ² Martins, F. M., et al., in *Proceedings of the 6th European Congress of Cardiology*, Madrid, 1972.
- ³ Watson, D., *Clinica Chimica Acta*, 1960, 5, 637.
- ⁴ Eggstein, M., and Kreutz, F. H., *Klinische Wochenschrift*, 1966, 44, 262.
- ⁵ Canal, J., and Ricquois, M., *Feuillets de Biologie*, 1971, 12, 53.
- ⁶ Glueck, C. J., Heckman, F., Schoenfeld, M., Steiner, P., and Pearce, W., *Metabolism*, 1971, 20, 597.
- ⁷ Miguel, M. J. P., et al., *Annales de Biologie Clinique*, 1973, 31, 147.

Psychiatric Nurse as Therapist

SIR,—Dr. I. M. Marks and his colleagues are to be congratulated on their well-designed paper (21 July, p. 156) which raises a number of topical issues, especially for those engaged on similar ventures.

At this unit we have, since 1961, operated an 18-month postgraduate course for nurses who hold as their basic qualification at least the S.R.N. or R.M.N. The regional hospital board allocates four bursaries (equivalent to a staff nurse's salary) and, in conjunction with the Portsmouth Polytechnic, a special diploma is awarded to candidates after successful completion of the training. Despite obvious differences in clinical approach and setting, our aims are not unlike those outlined by Dr. Marks—for example, the use of behaviour modification through the repeated coupling of two individuals¹ or small "transactional" groups for mothers and pre-school children, involving social comparison processes, especially modelling.² Our curriculum seeks to impart a substantial body of theoretical knowledge, authenticated by continuous practical involvement and the sharing of learning experience with other team members by means of study groups, tutorials, and seminars. There is increasing emphasis on extramural work, follow-ups, and crisis intervention.

Because much of our enterprise deals with multidimensional family systems rather than individuals, it is harder to apply "accurate measurements" to a comparison of therapeutic effectiveness. Frequently a nurse therapist replaces absent or "invalid" parents; or he or she might have to feed back specific activity cues into the home of an autistic child. Nonetheless, we would agree with all the general comments made about the value and functional aspects in the training of nurse therapists. A recent cost analysis points to a possible reduction in drug expenditure; and a prospective study of "breakdown recurrence," now under way, suggests no significant difference between patients treated by nurses and other therapists. The problem of grading on the Salmon scale seems to us of paramount importance. If so much effort is being devoted to training and skilled nurse therapists become ultimately sidetracked into administration, one might perhaps question if this constitutes "the most rational deployment of N.H.S. personnel."—I am, etc.,

CHRISTOPHER HAFFNER

Wessex Unit for Children and Parents,
Portsmouth, Hants

- ¹ Haffner, C., *Deutsche Schwesternzeitung*, 1968, 10, 601.
- ² Mundy, L., *Journal of Child Psychotherapy*, 1972, 3, 77.

SIR,—I found the paper by Dr. I. M. Marks and others (21 July, p. 156) very interesting.

I am aware of a somewhat similar attempt in Canada which was not too successful. However, in this instance attempts were made to utilize the Canadian equivalent of state enrolled assistant nurses, probably because they were the only staff readily available at the time. I note, too, that Dr. Marks and his colleagues were able to select their five trainees out of 80 applicants.

Secondly, the authors say that "qualified therapists need to be linked either to a psychiatrist or to a psychologist." I wonder how they would perform if they found themselves in a post where the psychiatrist or psychologist was either not as enthusiastic as the authors for utilizing nurses in this way or, possibly more important, was not as well informed in behaviour therapy principles and practice as the authors (or possibly even the nurse-therapist)? I am now a little out of touch with the situation in Britain, but certainly in Canada there are still many areas where knowledge of these is somewhat limited. I am not entirely clear whether the authors feel these nurse-therapists could function adequately in such a situation. In other words, could they function where the psychiatrist acted purely as a referrer of patients, or do they require that either he or the psychologist act in a supervisory capacity?

In view of recent evidence suggestive of the advantages in some conditions of in vivo (or in practice) exposure over the use of fantasy (or imagination), the use of some such therapists becomes even more important, particularly on account of the time factor involved. I think, therefore, that it would be a pity if what potentially seems to be a very useful development were to fall into disrepute through misuse.—I am, etc.,

MICHAEL J. C. THOMSON

Department of Preventive Medicine,
Dalhousie University,
Halifax, Nova Scotia

Drug-induced Antiplatelet Antibodies

SIR,—Thrombocytopenia is a well-recognized complication of treatment with phenylbutazone and generally results from marrow hypoplasia. However, in a recent case in this hospital a specific platelet antibody was also present, and as far as we are aware this has not previously been recorded.

The patient, a 58-year-old Caucasian woman, was admitted with one week's history of bruising. Six weeks before admission she had been prescribed phenylbutazone 100 mg thrice daily for low back pain, which she had continued up to the time of admission. Otherwise she had had no recent medication of any kind, and her job as a finisher in tailoring had not exposed her to any unusual chemicals.

On admission she was found to have widespread bruising on the limbs, but there was no evidence of bleeding elsewhere and the spleen was not palpable. The initial blood count showed: haemoglobin 10.9 g/100 ml, W.B.C. 2,800/mm³ (neutrophils 330/mm³), platelets 4,600/mm³, E.S.R. 39 mm in one hr. The blood film was normocytic and normochromic and the reticulocyte count was 1.4%. Routine investigations showed normal blood urea and electrolyte levels, liver function tests, and chest x-ray and negative rheumatoid factor, latex and antinuclear factor tests. Serum proteins were 6.2 g/100 ml (albumin 3.7 g/100 ml, globulin 2.5 g/100 ml); protein electrophoresis demonstrated an increase in the α -globulin and quantitative

immunoglobulin estimations an increase in IgM (IgG 510 mg/100 ml, IgA 165 mg/100 ml, IgM 430 mg/100 ml). Plasma haptoglobin level was 200 mg/100 ml and the direct Coombs test negative. The sternal marrow aspirate showed some reduction in cellularity; erythropoietic and granulopoietic activity was normal, but there was a relative decrease in the number of mature granulocytes. Lymphocytes and plasma cells appeared normal, but the number of megakaryocytes was reduced.

Investigations for platelet and leucocyte antibodies were carried out. For platelets the saline agglutination test at 4°, 20°, and 37°C was negative, complement fixation test at 37°C was positive in the presence of phenylbutazone in a concentration of 1 mg/ml, and the Coombs consumption test at 20°C gave a negative result. For leucocytes the saline agglutination test at 4°C, 20°, and 37° was negative and the complement fixation test at 37°C was negative.

The patient was treated with prednisolone, 60 mg/day for 3 weeks when most of the bruising had subsided and she was discharged taking 30 mg/day, the dosage being slowly tailed off over the next four months. There was a gradual improvement in all parameters of the blood count, and at four months the haemoglobin, W.B.C., and platelet count had returned to the normal range. The patient has been entirely symptom-free since her discharge from hospital.

Though drug-induced thrombocytopenia has been described following a large variety of drugs, in only a few has a specific platelet antibody been described.¹ In this patient the general pancytopenia and the hypocellular marrow suggest that the major effect of phenylbutazone was direct marrow toxicity, and this is supported by the slow haematological response despite the use of steroids. However, both direct marrow suppression and platelet destruction by specific antibodies are recognized in the thrombocytopenia seen with thiazides.^{2,3} We are postulating a similar mechanism for the thrombocytopenia caused by phenylbutazone in our patient, and suggest that antiplatelet antibodies should be sought for in all cases of this type.

We would like to thank Dr. Maxwell Telling for permission to report this case.

—We are, etc.,

C. DAVIDSON
S. M. MANOHITHARAJAH

General Infirmary,
Leeds

- ¹ Aster, R. H., in *Haematology*, ed. W. J., Williams, E., Bentler, A. J., Erslev, and R. W., Rundles, New York, McGraw-Hill, 1972.
- ² Rodrigues, S. U., Leikin, S. L., and Hiller, M. C., *New England Journal of Medicine*, 1964, 270, 881.
- ³ Eisner, E. V., and Crowell, E. B., *Journal of the American Medical Association*, 1971, 215, 480.

Infectious Mononucleosis

SIR,—I have read with interest the article by Dr. H. Pullen (12 May, p. 350). Infectious mononucleosis still represents a rather frequent haemopathy with peculiar features and some obscure nosological aspects. In Dr. Pullen's article the description of the clinical symptomatology fails to include two signs which may be very useful in practice.

The first is eyelid oedema, which may cause narrowing of the palpebral fissure. According to Hoagland and Wintrobe¹ this symptom occurs in 36% of cases. But its diagnostic importance increases because of its early appearance, as confirmed by my recent observations.

The second feature missing from the article is the pain caused by the swollen lymph nodes. In certain cases of lymphaden-

opathy the pain in the neck is so severe that the patient cannot rest his head on the pillow. This second sign, when present, is very useful in the differential diagnosis. In fact, the pain from swollen lymph nodes is not present unless there is a suppurative process.

I agree with Dr. Pullen's remarks on the treatment of infectious mononucleosis. Apart from the fact that cortisone and antibiotics are practically useless in this disease and that antibiotics may carry the added danger of toxic manifestations, we must definitely avoid cortisone for another important reason—that is, the fact that it may influence the haematological picture in cases of acute leukaemia. It is possible that acute leukaemia in a child may be misdiagnosed as infectious mononucleosis and treatment with cortisone may result in prolonging the wrong diagnosis by several weeks or even longer.—I am, etc.,

NEVIO QUATTRIN

Department of Haematology,
Ospedale Riuniti di Napoli,
Naples

¹ Hoagland, R. J., in Wintrobe, M. M., *Clinical Haematology*, p. 1230. London, Kimpton, 1967.

Pollen Count and Asthma

SIR,—Dr. S. A. Copeland refers (21 July, p. 175) to two young adults, both with long histories of hay fever, having their first attack of asthma after a downpour of rain, when the pollen count had fallen from 130 to 23/m³. Rainfall provokes an increase in the number of certain mould spores, namely, sporobolomyces and basidiospores, and it is possible that his patients had acquired a sensitivity to these. It is noteworthy also that whereas grass pollen sensitivity rarely causes rhinitis and asthma, mould spore sensitivity is more likely to cause asthma than rhinitis.

While on the subject of histories misleadingly suggestive of grass pollen sensitivity it may be mentioned that allergic respiratory symptoms due to mowing grass may be due to throwing the spores of *cladosporium* into the air and this of course may or may not be in the grass pollen season.—I am, etc.,

B. J. FREEDMAN

Dulwich Hospital,
London S.E.22

Medically Oriented Language Courses

SIR,—Following Britain's entry into the European Economic Community with the increasing potential mobility of doctors from Britain to other countries in Europe, the attention of the Board of Science and Education has been drawn to the need for medically oriented language courses.

The board's Medical Education Panel agrees that communication should be facilitated—for example, in relation to the technical aspects of the increasing range of medical specialties. Before any decisions are made concerning the most appropriate agencies to organize such courses and related questions the panel would like to have some indication of the demand. I should therefore be glad to hear from any members of the profession who would be interested in medically oriented language courses. Alternatively, members

may prefer to write direct to the Secretary of the Medical Education Panel at B.M.A. House with suggestions for the organization of such courses.—I am, etc.,

JAMES D. E. KNOX

Chairman,
Medical Education Panel,
Board of Science and Education, B.M.A.
B.M.A. House,
London W.C.1

Certification of Competence

SIR,—I am still able to recall the thrill of hearing my name announced as a successful candidate in the Conjoint examination. I was at last a "proper doctor" and after a further period of training was entrusted to embark upon my chosen career as a general practitioner. After 15 years I have learned much, and in the next 15 I shall learn much more. I have endeavoured to keep myself abreast of medical knowledge and I hope and believe that I have provided a competent service for my patients. I am naive enough to believe that qualification laid such a trust upon me.

But current rumours suggest that this is not to be enough. It is proposed in certain quarters that I shall need to obtain a "certificate of competence" in order to be financially rewarded for practising one aspect of my art.

My question to the members of my profession (and to our Association) is—why only for family planning? Is it not logical also to propose similar certificates for competence in the treatment of eye diseases, interpretation of E.C.G.s, or in the use of potentially dangerous psychotropic drugs? Let it be quite clear that I am a committed disciple of continuing postgraduate education, but my plea to our Association is that we might have full debate on the principle of recognition of competence before we are presented with another fait accompli. (The precedent is already with us in the form of the obstetric list.) For once, let us know the mind of the profession before we step further along the road which leads to two grades of general practitioner—those who are certifiable and those who are not!—I am, etc.,

GEOFFREY R. HORTON

Newark, Notts

Points from Letters

Breast Feeding

Dr. MARY BARTLETT (Wiggington, near Banbury) writes: Bearing in mind the ethologists' concept of "imprinting," education of mothers in the advantages of breast feeding will be useless unless midwives are also educated not to give the baby anything through a rubber teat in the first few days of life. This used to be the rule when I was first in practice, and breast feeding was respectable and practised.

Radioactivity and Lymphocytic Chromosomes

Dr. N. W. RAMSEY, Ph.D. (Charing Cross Hospital Fulham) writes: In your leading article (14 April p.68) reference is made to the dose to the synovial tissues after a typical intra-articular treatment of the knee joint by the injection of either radioactive

Earnings of G.P.s and Hospital Doctors

SIR,—For the second time in a month I find myself in conflict with views expressed by an adviser to the B.M.A. I refer to the letter from Mr. C. I. Brown (25 August, p. 458) on the subject of the remuneration of consultants and general practitioners. In this letter he states that "a significant number of [consultants] receive additions of 20% or more to their basic remuneration in the form of distinction awards." I suggest that it is this destructive argument that is responsible for the present low level of whole-time consultants' salaries. It is surely time that it is finally rejected. It is the level of basic remuneration that has fallen so far behind and is causing dissatisfaction. Distinction awards are supposedly given for especially meritorious service. They are not received by all consultants and should not be confused with basic remuneration.

I find it depressing that an adviser to the B.M.A. should not be able to reach this clear conclusion for himself and should reproduce this well-worn fallacy about distinction awards in the *B.M.J.*. When a letter such as this appears it is difficult for whole-time consultants to believe that the B.M.A. has any real sympathy with their predicament or indeed any real understanding of it. The fact is that many competent and hard-working whole-time consultants are entirely dependent on their basic remuneration. Perhaps it is significant that the general practitioners have rejected distinction awards altogether.—I am, etc.,

R. D. H. RYALL

Liverpool Regional Radiotherapy Centre,
Clatterbridge Hospital West,
Bebington, Cheshire

Widow's Pensions

SIR,—May I strongly urge that it is vital that doctors' widows' pensions should be raised at least another 16½% to make them 50% of their husband's pension; or in the case of those who forwent the retirement grant the figure should be raised from 50% to 66½%.

This should clearly be a first charge on any surplus in the pensions fund.—I am, etc.,

T. S. GOODWIN

Wannock,
near Polegate, Sussex

gold or yttrium. I think it should be emphasized that the figures quoted are approximate figures based on a mean dose, assuming that the administered radioactivity is dispersed through a volume of tissue of weight 300 g. These figures therefore provide a rough guide to the radiation level, but it is important to bear in mind that the dose to the synovium is 6,000-7,000 rads¹ the depth of tissue, receiving a dose at this level or higher being 1-2 mm.

Impatient Patients

Dr. J. MAHMOOD (London E.11) writes: Am I unique in having met during the past few days a mini-epidemic of patients—both of long-standing registration and newcomers—asking to be referred for investigation before I have even been asked for my advice, diagnosis, or treatment? My reaction is to give them their hearts' desire and then ask the executive council to remove them from my list. Am I oversensitive?