CORRESPONDENCE

Correspondents are asked to be brief

Doctors in South Africa

I. N. Bernadt, F.R.C.S.	632
Female Sterility Produced by Investigation	
F. W. Wright, F.F.R., and Sir John Stallworthy, F.R.C.O.G.	632
Measles	
S. R. Saxena, M.D.	632
Analgesics and the Kidney	
I. C. Calder, PH.D., and others	633
Irradiation Treatment of Rheumatoid Arthritis	
J. M. Gumpel, M.R.C.P.	633
Survival in Coalworkers' Pneumoconiosis	
B. Curry, M.Sc.	633
Edrophonium in Diagnosis of Cholinergic Crisis	
D. L. McLellan, M.R.C.P.	634
Sterility after Vasectomy	
I. L. Craft, F.R.C.S., and others; G. J. Fellows, F.R.C.S.	634
Anaesthetic Safety Devices	
R. Parfitt, F.F.R.	635
Nephrotoxic Drugs	
B. M. Tune, M.D., and R. L. Kempson, M.D.	635
Hypotension in Tetanus	
B. Fistein, M.R.C.P.ED., D.T.M.&H.	635

Rubella Surveillance: Recent Data from Scotland

N. R. Grist, F.R.C.P.ED., F.R.C.PATH.	636
Treatment of Diabetic Retinopathy	
W. G. Oakley, F.R.C.P.; E. N. Wardle, M.R.C.P.	636
Neonatal Jaundice and Maternal Oxytocin Infusion	
A. Ghosh, M.R.C.P., and F. P. Hudson, F.R.C.P.ED.; W. G. Mills, F.R.C.O.G.; M. C. R. Fisher, S.C.M.	636
Apnoea after Sacral Extradural Analgesia	
J. A. Lee, F.F.A.R.C.S.	637
Delayed Attacks of Malaria in Visitors to the Tropics	
T. Wilson, M.D.; L. J. Bruce-Chwatt, M.D.	637
Specific Immunoglobulin for Prevention of Serum Hepatitis	
D. N. S. Kerr, F.R.C.P.	637
Hazards of Laparoscopy	
A. D. Noble, F.R.C.S.ED.	638
Depression and Organic Disorder	
G. Dodds, M.B.	638
Protection of Cytomegalovirus-infected Cells by IgG Antibody	
P. Diosi, M.D., and Georgeta Herzog	638

Endotoxic Shock Complicating Trans-hepatic Cholangiography

B. B. Hickey, F.R.C.S.	638
Antibiotic-induced Meningitis	
W. I. H. Shedden, M.D.	638
Hypomagnesaemia after Parathyroidectomy	
R. D. M. Scott, M.R.C.P., and J. A. Strong, F.R.C.P.	639
Incidence and Prevalence	
R. Ulman, M.R.C.P.	639
Impatient Patients	
D. H. Ryde, M.R.C.G.P.	639
Aetiology of Reye's Syndrome	
P. G. B. Johnston, M.R.C.P.; A. Elkeles, M.D.	640
Erudition and Aphorisms	
J. Apley, F.R.C.P.	640
Problem Oriented Record Card	
P. R. V. Tomson, M.R.C.G.P.	640
Diuretics and Renal Pain	
N. G. Kounis, M.D.	641
Conference of Royal Colleges and Faculties in Scotland	
J. H. Croom, F.R.C.P.ED., and others	641

Doctors in South Africa

SIR,—The list of salary scales, set out in the Table below, is taken from a recent advertisement in the *South African Medical Journal* for doctors in one of the provincial administrations. You will notice the differential salary scale, in which coloured and Asian medical officers and registrars receive from 76% to 81%, and African doctors 61% to 74%, of the white doctors' salaries. Similar salary scales exist in all state hospitals, in all provincial hospitals and in most municipal hospitals. Nurses' salary scales are equally discriminatory.

TABLE—Minimum and Maximum Commencing Annual Salaries (in Rands) Granted in Respect of Previous Appropriate Experience

I. Medical: White		(Rands p.a.)	
Principal specialist		12,600	(fixed)
Senior specialist		12,000	(fixed)
N.B. To the ranks mentioned hereunder a pensionable allowance of 15% in respect of Whites and 17% in respect of non-Whites are payable on the basic salary of full-time personnel with effect from 1 April 1973.			
Specialist		8,400	9,300
Senior superintendent			8,400
Senior medical officer			8,400
Medical officer		5,700	8,100
Registrar		5,700	8,100
(Must be qualified for at least two years)			
Clinical assistant		5,700	8,100
(Must be qualified for at least two years)			
II. Medical: Coloured/Asian			
Registrar		4,350	6,600
Medical officer		4,350	6,600
III. Medical: Non-White			
Registrar		3,480	6,000
Medical officer		3,480	6,000

This must be a unique situation in the world.

Though no law prevents coloured doctors from issuing instructions to white nurses, this is not allowed in provincial and state hospitals. Coloured doctors are not allowed to visit white wards. Any coloured colleague from Britain would not be able to attend ward rounds in the white wards of South African teaching hospitals.

The South African Medical Association has frequently condemned these discriminatory practices, but has apparently not done much more about it.

The South African College of Medicine has not commented on this situation, as far as I know, but coloured and African candidates for the college examinations are never allowed to examine white patients in their practicals.

The comments of the B.M.A., the royal colleges, and your readers would be most interesting. It is only by exposing this remarkable situation that the combined weight of informed opinion may be mobilized to exert pressure to effect change.—I am, etc.,

IAN BERNADT

Cape Town, South Africa

*The B.M.J. has now stopped accepting advertisements from South Africa unless assured that no discriminatory salary attaches to the post advertised.—Ed., B.M.J.

Female Sterility Produced by Investigation

SIR,—Some doctors still use iodized oil as the contrast medium for hysterosalpingography. It has serious disadvantages in that oil embolism is a danger and deaths have been reported, and pelvic adhesions may

form following peritoneal reaction to the iodized oil. This can be clinically silent and detected later by repeat hysterosalpingography, laparoscopy, or at laparotomy. On the other hand, there can be an acute reaction with a clinical picture of pelvic peritonitis. When unsuspected pelvic tuberculosis is present we have seen violent reactions with pelvic abscess formation.

We have investigated several patients with bilateral tubal occlusion which appeared to develop at the fimbrial end of the tubes following hysterosalpingography using iodized oil. Normal tubal patency had been confirmed at the initial examination.

We consider oil contrast media for hysterosalpingography to be both dangerous and unnecessary. It is tragic, and there could be medicolegal consequences, when a technique demonstrating normal tubal patency causes subsequent sterility. In reminding doctors of the possible complications of injecting iodized oil into the peritoneal cavity we recommend the use of water soluble contrast media, which do not have any of these disadvantages and which allow a quicker, more certain, and less dangerous demonstration of tubal anatomy and patency.—We are, etc.,

F. W. WRIGHT
JOHN STALLWORTHY

Churchill Hospital,
Oxford

Measles

SIR,—I am glad that the question of pathogenesis of measles has been raised in your leading article (28 July, p. 187). Despite the advent of measles vaccine this disease remains a challenging problem for the treating doctor, especially in India.

Br Med J: first published as 10.1136/bmj.3.5881.632-b on 22 September 1973. Downloaded from <http://www.bmj.com/> on 26 April 2023 by guest. Protected by copyright.