SIR,—While the report by Dr. D. P. Davies and others (1 September, p. 476) shows a clear association between an increased incidence of neonatal jaundice and induced labour there can be little to commend the title of their communication. They discuss in some detail the mechanism by which hepatic enzyme immaturity might occur in the offspring of women who do not spontaneously enter labour. They quote the "gestational age" as being around 39 weeks in the group of women whose labour was induced without reference to the well recognized inaccuracy of calculating the anticipated date of confinement from a patient's recollection of her last period.

There is a strange uniformity of the 39-week figure in the three groups. I submit that those who started spontaneously in labour might well have been at term, and those who were induced (presumably because they failed to start in labour at "39 weeks") might have been significantly immature. It is seldom that the precise date of conception can be ascertained and, though postmaturity is well recognized, the majority of induced labours will probably be preterm.

The title of the article could only be justified if the authors had compared their results with a control series in which labour was induced without oxytocin. It should surely have read "Neonatal Jaundice and Induced Labour."—I am, etc.,

WILFRID G. MILLS

Birmingham Maternity Hospital

SIR,—Dr. D. P. Davies and others (1 September, p. 476) wonder if the infants in Group C (oxytocin induced) are particularly susceptible to hyperbilirubinaemia. Could not the explanation be that a fair proportion of these babies are immature?

Estimation of the gestational period is nowadays notoriously difficult. Could not this immaturity be the reason why the mothers had not gone spontaneously into labour?—I am, etc.,

M. C. R. FISHER

Oxford

Apnoea after Sacral Extradural Analgesia

SIR,—Dr. S. V. Humphries's interesting "Personal View" (25 August, p. 452) points out a danger of sacral extradural analgesia and how he dealt with it successfully. This is an excellent and safe form of pain relief increasingly being used in both our labour wards and operating theatres.

The risk of puncture of the spinal dural sac, a structure which usually but not always ends at the level of the second sacral vertebra, can be minimized by using a needle no more than 4 cm and by inserting it no further once the sacrococcygeal membrane has been pierced. In spite of care and considerable experience in the use of the technique, I have pierced the dura in this situation twice in one morning. The injection of the local analgesic solution should not be made until the anaesthetist is certain that neither blood nor cerebrospinal fluid can be aspirated. Should cerebrospinal fluid appear, while experienced workers may decide to carry on with the form of analgesia, the occasional anaesthetist would be well advised to abandon it to avoid the possibility

of apnoea, so dramatically described by Dr. Humphries.

I am writing this letter to defend an excellent and safe form of pain relief, which, however, may like other techniques result in avoidable complications. It is extremely rare for harm to come to patients who develop apnoea after extradural block, either lumbar or sacral, if they are adequately treated by intermittent positive pressure ventilation, plasma volume expanders, elevation of the legs, and very occasionally the injection of pressor amines.—I am, etc.,

J. ALFRED LEE

Southend General Hospital

Delayed Attacks of Malaria in Visitors to the Tropics

SIR,—I feel sure that Dr. Paul Horstmann (25 August, p. 440) would have changed the wording—and perhaps the title—of his article if he had been able to consult some of the papers on this subject published in the early 1950s. The development of a primary attack of vivax malaria 6-14 months after leaving the malarious area was commonplace in servicemen returning from the war in Korea. This long latent period was regarded as typical of temperate zone strains of *Plasmodium vivax* and contrasted with the short latent periods found in a number of tropical zone strains, notably those from New Guinea. From the details given, Dr. Horstmann's patients might have acquired their vivax infections almost anywhere along the route to India.

I live in retirement and can quote only three references,¹⁻³ but more should be available to anyone within reach of a library dealing with tropical medicine. Dr. Horstmann's article may well have served a useful purpose by reminding today's doctors that these long latent periods do occur in some of the vivax malaria infections that might develop in returned travellers.—I am, etc.,

T. WILSON

Portstewart,
N. Ireland

¹ Covell, G., *Journal of Tropical Medicine and Hygiene*, 1954, 57, 151.

² Field, J. W., and Shute, P. G., *Studies from the Institute of Medical Research, Malaya*, No. 24, 1956.

³ Hankey, D. D., et al., *American Journal of Tropical Medicine and Hygiene*, 1953, 2, 958.

SIR,—Dr. Paul Horstmann's interesting paper (25 August, p. 440) is a salutary reminder of the well-known fact that chemoprophylaxis with chloroquine must not be mistaken for causal prophylaxis (especially when it comes to *Plasmodium vivax*). The latter results from the action of drugs on the pre-erythrocytic stages of plasmodia and cannot normally be obtained by 4-aminoquinolines. On the other hand, antifolic drugs, such as pyrimethamine or proguanil, have causal, prophylactic action, especially on *P. falciparum*.^{1,2}

Nevertheless, 4-aminoquinolines exert a high degree of suppressive, antiparasmodial effect, which is greater for *P. falciparum* than for *P. vivax*. This explains why persons exposed to malaria and given insufficient dosage of 4-aminoquinolines, during or shortly after their exposure, may have delayed attacks of *P. falciparum* but particularly of *P. vivax* and other relapsing infec-

tions. In Dr. Horstmann's study of six persons who had a delayed attack of vivax malaria the once-weekly dosage of chloroquine (500 mg of chloroquine phosphate, equivalent to 300 mg of base) was manifestly insufficient, especially as the drug regimen was limited to two weeks after the group's return to Denmark. The recommendations adopted by the World Health Organization,^{3,4} and generally accepted, call for at least four weeks of chloroquine prophylaxis at the above dosage after returning to a non-malarious area. Even then many *P. vivax* infections cannot be suppressed and the delayed primary attack or the first clinical relapse may occur several months later.

This drug-induced delay of the primary attack must not be confused with the infection by some *P. vivax* strains having a naturally long incubation period. It is regrettable that in the discussion of Dr. Horstmann's paper this essential difference has not been sufficiently stressed or fully explained. On the other hand, Dr. Horstmann's remarks concerning the practical diagnostic importance of recognizing the possibility of malaria in persons who were exposed to the infection several months before are pertinent and timely.—I am, etc.,

L. J. BRUCE-CHWATT

Ross Institute of Tropical Hygiene,
London W.C.1

¹ Peters, W., *British Medical Journal*, 1971, 2, 95.

² Bruce-Chwatt, L. J., *British Medical Journal*, 1971, 2, 91.

³ The Ross Institute Information and Advisory Service, Bulletin No. 2, *Antimalarial Drugs*, London, Ross Institute, 1972.

⁴ World Health Organization, *Chemotherapy of Malaria and Resistance to Antimalarials*, Technical Report Series No. 529, Geneva, W.H.O., 1973.

Specific Immunoglobulin for Prevention of Serum Hepatitis

SIR,—A limited supply of immunoglobulin containing a high titre of antibodies to hepatitis B-associated antigen (HB-Ag) has been prepared at the Lister Institute and will be made available through the Medical Research Council for a series of controlled clinical trials.

It has been decided, however, that there is one situation in which the uncontrolled use of this material is justified. There is already considerable evidence for the value of this¹ and similar^{2,3} material in preventing serum hepatitis after inoculation of material containing HB-Ag. Consequently a limited supply has been made available for distribution through the Public Health Laboratory Service for the treatment of patients, staff, and other persons who are accidentally inoculated with infective material such as blood products, and laboratory reagents which are known to contain the antigen. The recipient and the physician administering it will be asked to participate in a follow-up study which is being conducted by the Medical Research Council to establish its effects and to find the incidence of adverse reactions, if any occur.

In the event of a suspected accident involving inoculation with infective material or splashing of such material into the mouth or conjunctival sac, physicians should contact the nearest Hepatitis Reference Laboratory of the Public Health Laboratory Service, or in cases of doubt, should call Dr. Sheila Polakoff or Dr. Yvonne Cossart at the Cen-