to disagree with this term, but it seems appropriate to us and its meaning has been clearly defined.—We are, etc.,

R. HALL
DAVID EVERED

Department of Medicine,
University of Newcastle upon Tyne

- 1 Masi, A. T., *American Journal of Public Health*, 1965, **55**, 658.
- 2 Snyder, P. J., and Utiger, R. D., *Journal of Clinical Investigation*, 1972, **51**, 2077.
- 3 Ormston, B. J., Alexander, L., Evered, D. C., Clark, F., Bird, T., Appleton, D., and Hall, R., *Clinical Endocrinology*, in press.

I.U.D. and Hydrorrhoea

SIR,—Hydrorrhoea gravidarum is a rare syndrome. I should like to report a case in which the contributory factor was an intra-uterine contraceptive device, an association which has not previously been reported.

The patient conceived in spite of a Grafenburg ring which had been inserted two years previously. When some eight weeks pregnant she developed vague lower abdominal pain which resolved slowly over the next three weeks, uterine growth being satisfactory. An ultrasound examination at this time demonstrated an intrauterine pregnancy but failed to show the ring.

At a gestation of 22 weeks she developed a profuse watery vaginal discharge. The size of the uterus was compatible with her dates and the cervical os was tightly closed. The fetal heart was clearly audible and regular. From then on a watery fluid drained intermittently and in great quantity. It was alkaline and glucose free, but contained substantial quantities of protein. No fetal cells could be seen. An abdominal x-ray showed a normal fetus with a cephalic presentation. The ring was lying below the fetal head and above the internal os. She remained well until 32 weeks and fetal growth was good, but she then developed some pink staining of her discharge which was shown to be blood. A placental scan showed a grade I anterior placenta praevia.

Uterine growth continued and serial oestriol estimations were satisfactory until 35 weeks gestation when she developed some fresh painless vaginal bleeding, despite which the uterus remained lax and the cervix closed. The fetal heart was audible and regular and this was confirmed by a fetal electrocardiograph which showed no evidence of hypoxia. Bleeding continued throughout the day and so caesarean section was done. On incision of the lower segment a blood clot was seen from which protruded the broken Grafenburg ring. This was lying at the lower edge of the placenta, which had partially separated. The membranes were intact. A live female infant weighing 2.31 kg was delivered. The pelvis was normal apart from some fibrinous adhesions surrounding the right fallopian tube and ovary. Subsequent progress of mother and child was uneventful and there was no undue vaginal discharge.

Moir suggests¹ the hydrorrhoea gravidarum represents a failure of fusion of the decidua vera and the decidua capsularis. One patient is mentioned who had this complaint in three successive term pregnancies. Paalman and Veer² reported a series of 41 cases in which 11 (28.6%) had an associated placenta circumvallata and nine had a history of bleeding in early pregnancy. They felt that

the fluid probably resulted from partial stripping of the membranes.

These reports and the present case suggest that hydrorrhoea gravidarum is caused by some irritant between the membranes and the decidua, probably blood but possibly a foreign body. In the present case it may be that the collection of fluid was lying just below the placental margin and thus caused slight placental separation—yet one more hazard of the intrauterine contraceptive device.—I am, etc.,

J. T. WRIGHT

Royal Free Hospital,
Liverpool Road Branch,
London N.1

- 1 Kerr, J. Munro, *Operative Obstetrics*, p. 252 8th edn. by J. Chassar Moir and P. Myerscough. London, Bailliere Tindall and Cassell, 1971.
- 2 Paalman, R. J., and Veer, C. G. V., *American Journal of Obstetrics and Gynecology*, 1953, **65**, 491.

Propellant a Factor in Asthma Deaths?

SIR,—In suggesting causes of the recent epidemic of deaths among asthmatics both your leading article (25 November 1972, p. 443) and Herxheimer¹ have overlooked the strong possibility that the propellant used in aerosol cans for dispensing medication may have been a contributor. Some of these aerosol propellants have been implicated in sudden deaths due to "aerosol sniffing,"^{2,3} and one aerosol product (Pertussin) has been banned in the United States because of associated sudden deaths. Various halogenated hydrocarbons (haloalkanes), isobutane, propane, vinyl chloride, and other agents have been used as propellants in aerosol cans.^{2,4} Probably not all, but several of them have been used in preparations for inhalation by asthmatics. Some of these agents, notably some of the haloalkanes, have been shown to be associated with cardiac arrhythmia and ventricular fibrillation as well as other effects in animals.⁴⁻¹² In one human experiment, premature atrial and ventricular contractions were noted after haloalkane and isoproterenol inhalation.¹³ In some of these experiments, it was thought that the agent simply sensitized the heart to anoxia or to exogenous adrenergic agents,^{4,12} but subsequent experiments have shown cardiac effects without either anoxia or exogenous adrenergic agents.^{6,7,10,11} In addition, "aerosol sniffing" deaths have occurred from aerosols containing no adrenergic agents.² Both anoxia and adrenergic agents, however, enhance the cardiotoxicity of the haloalkanes.^{4,9,11,12} Stolley focused attention on the high concentration of isoprenaline in some of the preparations for asthmatics, and showed that excess asthma deaths occurred in those countries that had used the "high concentration" preparations.¹⁴

Herxheimer pointed out that isoprenaline is not cardiotoxic, and pointed out some discrepancies in the distribution of excess asthma deaths which Stolley's hypothesis failed to explain.¹ He went on to attribute the excess deaths to inadequate medical management as others had done.¹⁵ Unfortunately, this hypothesis is even less adequate than Stolley's in explaining the distribution of excess asthma deaths. It utterly fails to account for the rise in asthma deaths during the time period which accompanied the introduction of the pressurized aerosol asthma preparations in England,^{14,15} and the fall in those deaths after an official warning about use of pressurized medication by asthmatics was issued.¹⁷

Although several investigators have argued that

asthmatic aerosol preparations are safe when properly used,^{9,15,18,20} it appears likely that in sensitive individuals, a combination of overuse, hypoxia, adrenergic agents and certain aerosol propellants could produce cardiac arrhythmias and ventricular fibrillation. Increases in either hypoxia or concentration of adrenergic agents would probably increase the likelihood of ventricular fibrillation from these preparations.

The associations noted by Stolley¹⁴ may have been with a preparation which accidentally combined high concentrations of adrenergic agent with the most cardiotoxic of the propellants. Other combinations of aerosol propellant and adrenergic agent could readily account for the spotty distribution of excess deaths noted by Herxheimer.—I am, etc.,

VICTOR E. ARCHER

National Institute for Occupational Safety and Health,
U.S. Department of Health, Education and Welfare,
Salt Lake City,
Utah, U.S.A.

- 1 Herxheimer, H., *British Medical Journal*, 1972, **4**, 795.
- 2 Bass, M., *Journal of the American Medical Association*, 1970, **212**, 2075.
- 3 Baselt, R. C., and Cravey, R. H., *Journal of Forensic Sciences*, 1968, **13**, 407.
- 4 Reinhardt, C. F., Azar, A., Maxfield, M. E., Smith, P. E., and Mullin, L. S., *Archives of Environmental Health*, 1971, **22**, 265.
- 5 Kilen, S. M., and Harris, W. S., *Journal of Pharmacology and Experimental Therapeutics*, 1972, **183**, 245.
- 6 Taylor, G. J., Harris, W. S., and Bogdonoff, M. D., *Journal of Clinical Investigation*, 1971, **50**, 1546.
- 7 Taylor, G. J., and Harris, W. S., *Journal of Laboratory and Clinical Medicine*, 1970, **76**, 857.
- 8 Taylor, G. J., and Harris, W. S., *Journal of the American Medical Association*, 1970, **214**, 81.
- 9 Clark, D. G., and Tinston, D. J., *Annals of Allergy*, 1972, **30**, 536.
- 10 Flowers, N. C., and Horan, L. G., *Journal of the American Medical Association*, 1972, **219**, 33.
- 11 Mullin, L. S., Azar, A., Reinhardt, C. F., Smith, P. E., and Fabryka, E. F., *American Industrial Hygiene Association Journal*, 1972, **33**, 389.
- 12 Azar, A., Zapp, J. A., Reinhardt, C. F., and Stopps, G. J., *Journal of American Medical Association*, 1971, **215**, 1501.
- 13 Brooks, S. M., Mintz, S., and Weiss, E., *American Review of Respiratory Disease*, 1972, **105**, 640.
- 14 Stolley, P. D., *American Review of Respiratory Disease*, 1972, **105**, 883.
- 15 Harris, M. C., *Annals of Allergy*, 1971, **29**, 250.
- 16 Smith, J. M., *Lancet*, 1966, **1**, 1042.
- 17 Inman, W. H. W., and Adelstein, A. M., *Lancet*, 1969, **2**, 279.
- 18 Azar, A., Reinhardt, C. F., Maxfield, M. E., Smith, P. E., and Mullin, L. S., *American Industrial Hygiene Association Journal*, 1972, **33**, 207.
- 19 McClure, D. A., *Toxicology and Applied Pharmacology*, 1972, **22**, 221.
- 20 Silverglade, A., *Journal of the American Medical Association*, 1971, **215**, 118.

Hypergastrinaemia in Rheumatoid Arthritis

SIR,—While one may excuse Dr. P. J. Rooney and his nine colleagues (30 June, p. 752) for not knowing of early studies suggesting an increased incidence of achlorhydria in rheumatoid arthritis^{1,3} it is more surprising that they fail to note the report of Buchanan *et al.*⁴ concerning Sjogren's syndrome. These authors demonstrated achlorhydria in four of six patients with this condition. Since 20% of the patients reported by Dr. Rooney and colleagues have keratoconjunctivitis sicca this is surely of importance in discussing the significance of the demonstrated hypergastrinaemia. As theirs was a preliminary report one wonders whether a further paper is already in preparation describing the results of gastrin assays in patients with Sjogren's syndrome? Should this be the case the authors may care to note that among the many prescient "objectives for further