CORRESPONDENCE

Correspondents are asked to be brief

Health on the Cheap

P. J. E. Wilson, F.R.C.S. 112

Fatal Insect Stings

C. W. Thomson, F.F.A.R.C.S.; K. K. Nayak, M.D. 113

Lawn Mower Injuries

J. R. Hulme, B.M., and A. R. Askew, F.R.C.S.ED. 113

Septic Abortion and the Dalkon Shield

D. F. Hawkins, F.R.C.O.G. 113

Sensitivity to Rifampicin

D. J. Girling, M.R.C.P., and others. 114

Influenza in Families

W. O. Williams, F.R.C.G.P. 114

Computer Assisted Tomography

G. A. S. Lloyd, F.F.R., and J. E. Wright, F.R.C.S. 114

Nursing Staff and Work Load

R. Bennett, S.R.N., and others. 115

Accidental Injection of Potassium

A. G. L. Whitelaw, M.R.C.P. 115

Oral Contraceptives and Health

Ellen C. G. Grant, M.B. 115

Achalasia of the Cardia

F. R. Berridge, F.R.C.P., F.F.R. 116

Salaries in Occupational Medicine

R. E. W. Fisher, M.F.C.M. 116

Clinical Public Health Doctors and Community Medicine

P. O. Nicholas, M.F.C.M. 116

Review Body Report

N. Gordon, M.R.C.G.P., and others; D. K. Sambrook, F.R.C.S.; June Phillips, M.B., M.F.C.M.; M. H. Hughes, F.R.C.PATH. 117

G.P.s and Family Planning

D. D. Cowen, M.R.C.G.P. 117

Reward for Effort

I. C. Gilchrist, M.B.; W. B. Jamison, M.R.C.G.P. 117

Working Hours of Junior Staff

D. G. H. Stone, F.R.C.P. 118

An Overpaid Profession?

Anne H. Garner, M.B.; E. F. Brooks, M.B.; M. I. Davies, B.M.; N. H. Barley, M.R.C.G.P.; J. W. Baker, B.M.; R. R. Watkin, F.F.A.R.C.S.; A. R. Rogers, M.B.; C. W. D. Phillips, M.B. 118

Private Beds and the N.H.S.

S. O. Aylett, F.R.C.S. 119

Health on the Cheap

SIR,—Quite soon after the system of extra duty payments for hospital junior staff came into operation consultants were informed of a rising unease in administrative quarters that the sums being paid were too large. It was implied that some doctors were getting too much, that others might be cheating, and that consultants should therefore verify each and every claim before countersigning their juniors' claim forms. Consultant reaction to these overtures of bureaucratic distress was that the extra duty scheme, though ill-structured and inherently susceptible to anomaly and abuse, had been freely entered into by the Department of Health and Social Security and that it was no part of a consultant's duty to scrutinize the fiscal bona fides of his junior colleagues. This certainly was and is my own view.

After 1 July 1974, when the contractual 100-hour week for junior hospital doctors became an 80-hour week, it is obvious that the bill for extra duty payments will rise sharply, but no extra revenue is forthcoming to pay this. Instead, the alienated mandarins at the Elephant and Castle have issued suggestions

(or are they directives?) whereby the extra duty bill may be minimized. These include interspecialty cross-cover, reduction of cover at night, compression of clinical work into "office" hours, and the fatuous idea that consultants shall do their own juniors' extra duties for them for no extra payment.

Now if the bill for extra duty payments were in truth outrageous or exorbitant, consultants might feel constrained at least to examine the feasibility of some or all of the above proposals. I believe that we must ignore these proposals totally and resolutely. My belief is strengthened by a few simple calculations derived from an official statement of extra duty payments actually made to hospital doctors in this city during 1973-4. Badly presented, the grand aggregate (£32,000) might perhaps, as it was meant to do, cause an uncritical eye to bulge a little. But the heart of the matter is this. What has been the cost of guaranteeing to the public a professional service by trained specialists and their teams for every hour of every day of every year? And has this cost been unreasonable or excessive when

compared with similar services provided by other trades and professions?

The accompanying table embodies some of the major district specialties and "minor" but relatively expensive subregional specialties in one 470-bed hospital, which must differ little from many others in the country.

It will be seen that, on the basis of a 100-hour week for juniors and a 168-hour week for consultants ("the whole or substantially the whole of their time"), 40% of the hours worked are reimbursed by 10% of the moneys paid. Had an 80-hour week obtained, no less than 52% of the hours worked would have been paid for by extra duty allowances. Only if the total sum of such payments quadrupled would there begin to be an equitable relation between the cover provided and the remuneration given. Consultants' extra duties remain unpaid, it must be emphasized.

This ailing nation gets its emergency hospital services very cheaply. Taking Swansea as an exemplar, the cost of extra duty payments per bed per day in 1973-4, giving guaranteed multispecialty cover, was 11p in one of its two major hospitals and 8p in the other. The new scales of payment

Cost of Medical Staffing for some General and Special Acute Services in a 470-bed District Hospital, 1973-4

Specialty	Consultant	Senior Registrar	Registrar	S.H.O.	H.O.	Extra Duty Payment per annum	Extra Duty Payment per diem	Total per annum	Total per diem	Extra Duty Payment as % of Total
General medicine	£14,916 (2)	—	£3,192	£2,850	£2,061	£1,697	£4.65	£24,716	£67.71	6.8%
General surgery	£18,603 (2)	—	£6,384 (2)	—	£4,142 (2)	£1,109	£3.04	£30,238	£82.84	3.7%
Obstetrics and Gynaecology	£12,402 (2)	£4,179	£3,192	£2,664	—	£2,777	£7.61	£25,214	£69.00	11.0%
Paediatrics	£14,358 (2)	—	£3,192	£7,992 (3)	—	£2,653	£7.27	£28,195	£77.25	9.4%
Neurosurgery	£12,751 (2)	£4,179	£3,192	£5,700 (2)	—	£2,219	£6.08	£28,041	£76.82	7.9%
Neurology	£6,201	—	£3,363	£2,664	—	£2,390	£6.55	£14,618	£40.05	16.4%
Thoracic surgery	£7,599	—	£3,192	—	—	£2,394	£6.56	£13,185	£36.12	18.2%

Figures in parentheses denote number of posts in that grade. Part-time consultant salaries are reckoned as at the mid-point of the incremental scale, as are full-time salaries in all grades except where specifically known to be otherwise. Where extra duty payments were not made over a full year, notional daily average payments were calculated from actual total payments.

recommended by the Halsbury Committee would inflate these mind-boggling sums to 12p and 9p respectively.

This, surely is the form in which doctors and the public at large should have N.H.S. expenditure presented to them. It would then be plain to all that for the medical profession to help the D.H.S.S. to wallow a little less deeply in the financial mire in which it now flounders is not only super-erogatory but irrelevant. More important, this way of looking at the costs of the care of acute illness also offers a way of meeting those costs. If every patient in hospital, except perhaps the very young and the very old, contributed 10p a day to that hospital he could rest secure in the knowledge that this trivial premium—the cost of five cigarettes, a can of coke, a bar of chocolate, or 1½ pints of petrol—would ensure that whatever illness he had, whatever urgent service he might need, whatever unforeseen complication or reverse he might suffer, cover would be provided at any hour of the day or night. Doubling the levy to a mere 20p a day would, moreover, underwrite the desperately needed increase of salaries for nurses and ancillaries.

Politicians of all colours have long lacked the courage to tell their electors that this kind of service must be paid for. Perhaps, after all, the D.H.S.S. and the public do not want the kind of service we now struggle to provide. If so, let them say so. But neither public, politicians, nor placemen should expect the medical profession to compound their parsimonious folly.—I am, etc.,

P. J. E. WILSON

Neurosurgical Unit,
Morriston Hospital,
Swansea

Fatal Insect Stings

SIR,—In your leading article on this subject (18 May, p. 345) it was pointed out that respiratory tract obstructions—oedema of the glottis or bronchi—accounted for 70% of the deaths and that 58% of the deaths occurred within the first hour. It was also suggested that antihistamines may be injected intramuscularly or intravenously to combat the respiratory oedema; and by way of example promethazine hydrochloride 50 mg was quoted. It should be remembered that promethazine is a sedative and if injected intravenously can produce such a degree of sedation that the patient may become unconscious, a dangerous state to be in if the airway is partly blocked.

I recently saw a patient who had developed a minor degree of upper respiratory obstruction, having inadvertently eaten some pork, to which she was sensitive. Her breathing was laboured and she indicated that she could feel a sensation of swelling at the back of the tongue. I treated her with hydrocortisone intravenously to a total dose of 700 mg and at one stage I injected 5 mg of chlorpheniramine slowly into a vein. Immediately after this she became extremely sleepy and her breathing, which had until then been laboured but adequate, became slow and shallow and plainly inadequate. I found I could maintain her respiration by loud vocal encouragement and continuous tactile stimulation—in short, by shouting

and prodding—till she woke up 20 minutes later.

The fact that this was only a mild case of respiratory obstruction which ended harmlessly should not obscure the fact that the intravenous antihistamine, by depressing her consciousness, made the situation worse. Anaesthetists are trained to be aware of the danger of giving soporific drugs to patients with respiratory obstruction and I would urge that antihistamines be included in this group and should be used only with extreme caution.—I am, etc.,

CHARLES THOMSON

Department of Anaesthetics,
Newcastle General Hospital,
Newcastle upon Tyne

SIR,—I wish to draw attention to the potential danger of intramuscular or intravenous antihistamines, as recommended in your leading article (18 May, p. 345), in the treatment of acute anaphylactic shock caused by insect stings, particularly where bronchoconstriction is in evidence. Antihistamines may result in further bronchoconstriction and in any case they are probably useless in the treatment of acute anaphylaxis. The physiological antihistamine adrenaline together with intravenous corticosteroids and, of course, other supportive therapy would be the logical approach. I am rather surprised that you do not mention intravenous corticosteroids in your article.—I am, etc.,

K. K. NAYAK

Department of Clinical Research,
Philips-Duphar B.V.,
Weesp, Holland

** The sedative effects of an antihistamine may vary from individual to individual but it would be unlikely that the recommended dosage would produce serious respiratory depression in most patients. The patient quoted by Dr. Thomson was still rousable, and was fully awake in 20 minutes—a remarkably short period if the antihistamine and not the anaphylactic shock was chiefly to blame.

There is no evidence that antihistamines, in therapeutic dosage, can cause bronchoconstriction though they may not relieve it. Antihistamines should still be considered as important in the treatment of acute anaphylaxis, second only to the all-important adrenaline.

Corticosteroids are of little or no value in acute anaphylactic emergencies. We agree that once the emergency has passed they should be used to prevent recurrent or delayed reactions.—Ed., *B.M.J.*

Lawn Mower Injuries

SIR,—These summer months will produce their share of lawn mower injuries. In the United States in 1970 it was calculated¹ that there were 27m. mowers in use, resulting in 140,000 injuries during that year. (Figures for Britain are not available.) Most of these appear to be caused by rotary mowers, whose blades rotate horizontally to the ground at speeds of up to 3,000 r.p.m., the blade tips reaching a speed of 232 miles (373 km) per hour.

Lawn mower injuries are of two types:

(1) Direct injuries. These are caused by the blades of the mower striking either the operator or someone else, usually a child who has fallen into the path of the mower.

Injuries to adults usually involve the forefoot, causing at the least lacerations of the toes and, more seriously, complete or partial amputation of one or more toes, the wounds being contaminated with grass and soil. We have found² that these forefoot injuries tend to occur when the operator is using the mower going down a wet or dew-damp slope, when his feet slip from under him and slide under the protective canopy of the mower into the blades. Lasting sequelae of such injuries are disturbance of balance, limping, and diminished walking distance due to pain, especially if the hallux and/or first metatarsal are involved.

(2) Missile injuries. When small, hard objects such as stones are struck by the rotating blades they may be hurled from under the mower for distances of 50 yards (46 m) or more. Apart from causing structural damage to windows etc., they may cause serious injuries to the eye, even blindness.³ A survey in New Zealand⁴ reported that such missiles were the most common single cause of traumatic hyphema.

The following points should help prevent these accidents: (1) remove stones etc. from the lawn before mowing; (2) keep children and pets away from mowers in use; (3) remember that slopes, especially if wet, are potentially hazardous; and (4) wear robust footwear capable of secure purchase on the ground—golf shoes or ordinary shoes with spikes inserted will prevent slipping.—We are, etc.,

J. R. HULME

Basingstoke District Hospital,
Basingstoke, Hants

A. R. ASKEW

Radcliffe Infirmary,
Oxford

¹ DeMuth, W. E., et al., *Journal of the American Medical Association*, 1973, 225, 355.

² Hulme, J. R., and Askew, A. R., *Injury*, 1974, 5, 217.

³ Barsky, D., *Archives of Ophthalmology*, 1960, 64, 385.

⁴ Ferguson, R. H., and Poole, L. W., *Transactions of the Ophthalmological Society of New Zealand*, 1968, 20, Suppl. 10, p. 54.

Septic Abortion and the Dalkon Shield

SIR,—To remove an intrauterine contraceptive device (I.U.C.D.) in early pregnancy by traction on the strings, as advised by Dr. J. S. Templeton (15 June, p. 612), is very likely to cause abortion. Legal considerations apart, the risks of incomplete abortion, haemorrhage, and sepsis inherent in such a procedure must be far greater than the remote chance of intrauterine sepsis if the device remains in situ.

It seems likely that cases of septic abortion occurring in patients with I.U.C.D.s who become pregnant are due to attempted interference. If it is true that the infection can occur spontaneously, then an appropriate course of action would be to culture a high vaginal (not intracervical) swab and give a short course of antibiotics followed by a maintenance dose until the risk is considered to have passed.

The advice in Dr. Templeton's last paragraph could easily be misinterpreted by the patient as a promise of a therapeutic abortion if she became pregnant while wearing a Dalkon Shield. Such an undertaking would not come within the provisions of the Abortion Act, which, at least by implication, requires the opinions of two practitioners based on circumstances which present only