instances. There was no evidence of side effects.—We are, etc.,

ELIZABETH GREEN

Cell Barnes Hospital,
St. Albans

D. M. LYNCH

Bridge Hospital,
Witham

JAMES A. MCFADZEAN

I. M. PUGH

Chemotherapeutic Research Group,
May and Baker Ltd.,
Dagenham

- ¹ Allen, A. V. H., and Ridley, D. S., *Journal of Clinical Pathology*, 1970, 23, 545.
- ² Catzel, P., *Paediatric Prescriber*, 3rd edn. Oxford, Blackwell, 1966.

Malaria in the United Kingdom

SIR,—Professor B. G. Maegraith's (27 July, p. 252) merits in stressing the dangers of imported tropical disease are too well known to need emphasizing and all of us owe him a debt of gratitude for his pioneering work in this and many other fields. It is therefore all the more regrettable to find ourselves so misunderstood and taken to task for a non-existing misdeed (29 June, p. 707).

Professor Maegraith appears to disagree with one point of our paper—namely, that the protection of the health of each individual is, in the final account, his own responsibility and that the medical profession or any other bodies can only inform, advise, and guide. No one denies that the airlines or travel agents have shown little interest in the problem of imported malaria in spite of the fact that this has been frequently stressed by the daily press, many professional journals, and the B.B.C. It is also true that the excellent "Notice to Travellers" produced by the Department of Health and Social Security should be given to any traveller when he buys his ticket. Better distribution of this leaflet is now being arranged by the D.H.S.S. But the only way to improve the situation is to stress again and again the potential seriousness of the imported tropical disease and to obtain the willing co-operation of travel agents and airline managers. Sanctions or legalistic measures relevant to malaria are unlikely to be acceptable at a time when the general trend of the new international health regulations leans towards some relaxation of vexing sanity policy, even with regard to such diseases as cholera. The World Health Organization gave much thought to the problem of introduction of malaria and the relevant conclusions of a special meeting¹ were as follows:

"International travellers (tourists) should not be subjected to any compulsory antimalaria measures. Information on preventive measures should be given to them for their own protection when they enter or leave malarious areas together with warning cards instructing them how and where to report should they develop fever."

Such warning cards are used by the military² and have been adopted by the Council of Europe for civilian use, though the results of the latter method have not been spectacular. Obviously, more could be done by the medical profession, by the information media, and by the travel industry to make the public aware of the importance of diseases contracted abroad.

Neither the Liverpool nor the London School of Tropical Medicine has been idle in this respect. But once again, in the final account it is the informed individual who must assume the responsibility for his or her

own protection from malaria, either by reducing the degree of exposure, by taking prophylactic drugs, or by consulting his doctor in case of sickness after travelling overseas. No one could disagree with this and we are not aware of any serious difference of opinion between Professor Maegraith and ourselves.—We are, etc.,

L. J. BRUCE-CHWATT

B. A. SOUTHGATE

C. C. DRAPER

Ross Institute of Tropical Hygiene,
London School of Hygiene and Tropical Medicine,
London W.C.1

- ¹ World Health Organization, *Prevention of the Re-introduction of Malaria*, Technical Report Series No. 374, Geneva, W.H.O., 1967.
- ² McKelvey, T. P. H., et al., *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 1971, 65, 286.

Aspirin and What Else?

SIR,—Your leading article (6 July, p. 5) draws attention to the widespread use of aspirin in Britain and discusses the possible haemostatic defects that may arise. Until recently in our transfusion centre little attention was paid to aspirin ingestion in blood donors, despite the fact that platelet-rich plasma or platelet concentrate would be prepared from some of the donations. A survey in the U.S.A. showed that 42 (36%) of 118 donors gave a history of having taken a drug known to affect platelet function and the most common of these was aspirin: 36 (88%) of the 42 showed abnormal platelet aggregation.¹ In addition to the abnormal platelet aggregation in vitro aspirin also inhibits platelet function in vivo, since the bleeding time of normal subjects is increased after aspirin ingestion.² In view of these observations and the increasing use of platelet preparations, we decided to ascertain the frequency at which local blood donors had taken aspirin-containing compounds during the seven days before blood donation and to investigate platelet aggregation using adrenaline and collagen.

Donors were asked whether they had ingested any aspirin-containing substances within the previous seven days (as an aide memoire they were shown a list containing the names of 102 such substances). At the end of the normal blood donation in these donors a small sample of blood was taken and platelet aggregation (in response to collagen and adrenaline) was measured by a turbidometric method.³ Platelet counts were performed and plasma salicylate levels also measured. The findings of the medical questionnaire were broken down into the following groups. Group I, donors who had taken no aspirin-containing compounds or any other drug for seven days before donation. Group II, donors who had taken aspirin-containing compounds during the period of seven days before donation. Group III, donors who had taken other drugs during the period of seven days before donation but excluding those in group II. The donors in group III had taken either paracetamol, sleeping tablets, anti-histamines, or antidepressants. Altogether 111 donors were studied in this survey.

Our survey showed that 33% of donors had taken aspirin during the seven days before donation and a further 14% had taken other medicines within this period. There was a predominance of females in the aspirin-ingestion group. Of the donors with a history of taking aspirin (group II) 24% did not respond to collagen-induced aggregation compared with 8% of donors who had not taken any medication (group I). With adrenaline 70% of group II donors showed abnormal platelet aggregation compared with 29% of donors in group I who could not recall taking drugs. In group III the response to collagen and adrenaline was similar to group I. Positive plasma salicylate levels (>5 mg/100 ml) in 13% of group I donors showed that some may have unknowingly ingested aspirin during the seven days before donation.

The question now posed is: "Does abnormal platelet aggregation in aspirin-ingesting donors mean that these platelets are haemostatically ineffective? As little as 150 mg of aspirin has been found to suppress the second wave of adrenaline-induced platelet aggregation for as long as seven days but the in-vitro addition of only 10% normal platelets corrected the abnormality.¹ Stuart *et al.*⁵ showed that if the donor had not taken aspirin within 36 hours the platelets should prove haemostatically effective, but if within this time the donation should be pooled with those of other donors who have not taken aspirin. In view of this,, for regular platelet concentrate production of small pools (that is, for paediatric transfusions) aspirin-ingesting donors should be avoided but for adults, when large amounts are required, aspirin ingestion should not pose an unduly difficult problem.—We are, etc.,

R. M. BARNES

P. R. EVANS

D. S. SMITH

Wessex Transfusion Centre,
Southampton

- ¹ Schwartz, A. D., *Transfusion*, 1972, 12, 344.
- ² Quick, A. J., *American Journal of the Medical Sciences*, 1966, 252, 265.
- ³ O'Brien, J. R., et al., *Thrombosis et Diathesis Haemorrhagica*, 1966, 16, 752.
- ⁴ O'Brien, J. R., *Lancet*, 1968, 1, 894.
- ⁵ Stuart, M. J., et al., *New England Journal of Medicine*, 1972, 287, 1105.

SIR,—Aspirin in normal doses (650-1,300 mg) slightly prolongs the bleeding time in normal people. This is attributed to its effect in preventing the release of adenosine diphosphate (ADP) from platelets and thereby interfering with platelet stickiness. Your leading article (6 July, p. 5), suggested that profuse gastric haemorrhage after aspirin ingestion was due to an underlying abnormal haemostatic mechanism being potentiated by this effect.

Small doses of aspirin (10-100 mg) do not produce significant changes in bleeding times in normal individuals. However, in patients with the syndrome of nasal polyposis and asthma associated with aspirin idiosyncrasy Fisherman and Cohen¹ have shown greatly prolonged bleeding times after such minute doses. Using Macfarlane's method for measuring bleeding times I have been able to confirm that the bleeding times were prolonged to double or more in 12 patients with this syndrome, whereas in matched controls the bleeding times were unchanged or even shortened.

No patient in this series had a history of haematemesis, and only one patient of a larger series (45) with this syndrome had suffered a haematemesis. This would tend to support the hypothesis made in the leading article that an additional haemostatic defect needs to be present before aspirin will result in a profuse gastric haemorrhage.—I am, etc.,

J. C. DELANEY

Walton Hospital,
Liverpool

- ¹ Fisherman, E. W., and Cohen, G. N., *Annals of Allergy*, 1973, 31, 182.

Metaclopramide and Prolactin

SIR,—Dr. H. W. C. Ward (20 July, p. 169) draws attention to the possible dangers of administration of metaclopramide to breast