CORRESPONDENCE

Correspondents are asked to be brief

Repeated Acupuncture and Serum Hepatitis	
P. Alexander, D.Sc., and others	466
Smoking and Deep Vein Thrombosis	
Sir Richard Doll, F.R.C.P., F.R.S.	466
Endometriosis and Pseudopregnancy	
J. A. Chalmers, F.R.C.O.G.	466
Pitocin Buccal in Late Pregnancy	
T. M. Coltart, F.R.C.S.ED., and T. G. Nash, F.R.C.S.	467
Eugynon 30 and Dermatitis	
Delia F. Morris, M.B., M.F.C.M.	467
Tetracycline-resistant Group A Streptococci	
J. A. N. Emslie, M.D.	467
Antiemetics, Prolactin, and Breast Cancer	
M. O. Thorner, M.R.C.P., and others	467
Blood Donors and the Transfusion Service	
J. Wallace, F.R.C.P.GLAS.; M. H. Cooper, B.A.	467
Hypothermia, Thrombosis, and Acute Pancreatitis	
E. P. Savides, L.M.S.S.A.	468
Swimmers' Ears	
A. M. Birt, M.R.C.S.; C. R. Lucas, M.B.; T. Smyth; J. A. Lack, F.F.A.R.C.S.	468
Respiratory Complications of Obesity	
M. J. B. Farebrother, M.R.C.P., and G. J. R. McHardy, F.R.C.P.	469

Maternal Deaths in Scotland	
Louise F. Giles, M.A.	469
Death in Hospital	
C. J. Bennett, M.B.	469
Cleft Lip and Palate	
N. R. Dennis, M.R.C.P.	469
Removal of Tattoos	
G. T. Watts, F.R.C.S.	469
Allergy to Aspirin	
F. D. Hart, F.R.C.P., and E. H. M. Foxen, F.R.C.S.	470
Age Limit for Contraceptives	
Elizabeth H. Gregson, M.B.	470
Diabetic Autonomic Neuropathy	
G. E. Heard, F.R.C.S.	470
Measurement of Side Effects of Drugs	
A. Herxheimer, M.R.C.P.	470
Kidneys for Transplantation	
Elizabeth D. Ward, J.P.	470
Effects of Posture on Limb Blood Flow in Late Pregnancy	
J. W. Downing, F.F.A.R.C.S.; A. Singer, F.R.C.S.	470
Medicine in Belgium	
R. De Smet, M.D.	471
Non-comatose "Diabetic Coma"	
J. W. Todd, F.R.C.P.	471

Gut Motility Changes Causing Drug Malabsorption	
M. J. Kendall, M.D., and Linda Beeley, M.R.C.P.	471
Venoarterial Difference in α_1-Antitrypsin Levels	
J. S. Milledge, F.R.C.P.ED.	471
Perinatal Metabolism of Diazepam	
R. Erkkola, M.D., and others	472
Was it a Drug?—Forceval Protein	
A. C. A. Daniel, M.B.	472
Antibiotics in <i>Bacteroides fragilis</i> Infections	
D. T. L. Turner, F.R.C.S.	472
Cloxacillin Levels in Synovial Fluid	
J. H. Newman, F.R.C.S.	472
Medical Synovectomy	
J. M. Gumpel, M.R.C.P.	472
Aceiform Lesions in Combined Rifampicin Treatment in Africans	
U. Nwokolo, F.R.C.S.ED.	473
Treatment of Genital Herpes	
Paula J. H. Gosling, M.B.	473
Alternative Health Service	
P. Joan Bishop, M.B.	473
N.H.S. Contraceptive Services	
D. R. Cargill, B.M.	473
Saudi Arabian Medical School	
Susan Barlow, PH.D.	473

Repeated Acupuncture and Serum Hepatitis

SIR,—The letter by Dr. K. K. Hussain (6 July, p. 41) suggesting a possible link between repeated acupuncture and serum hepatitis is of considerable interest in view of the situation in China. On a recent visit to the People's Republic of China, under the auspices of the British Council, we saw the widespread use of acupuncture as a therapeutic measure, particularly in the commune hospitals. We even saw it being practised by children, on each other, in a class where acupuncture was being taught. The needles were placed in a sterilizing fluid but there was no evidence that any other form of sterilization was used.

We also noted the high incidence of hepatocellular carcinoma, and were told that one-third of these patients had a classical history of previous hepatitis and that the

liver usually showed cirrhosis at the time of diagnosis of the hepatic carcinoma. A sequence of events is suggested: virus hepatitis is extremely common, some patients proceed to develop cirrhosis, and some of these go on to develop primary liver tumours. The interest lies in how the virus is disseminated, and superficially it seemed to us that acupuncture could easily be responsible for this.—We are, etc.,

P. ALEXANDER

Chester Beatty Research Institute,
Belmont,
Sutton, Surrey

G. HAMILTON FAIRLEY
D. W. SMITHERS

Imperial Cancer Research Fund,
Department of Medical Oncology,
St. Bartholomew's Hospital,
London E.C.1

Smoking and Deep Vein Thrombosis

SIR,—The observation that venous thromboembolism is more likely to occur in hospital patients admitted for the treatment of myocardial infarction if they were non-smokers on admission than if they were smokers who then stopped smoking, which was reported by Dr. A. J. Handley and Mr. D. Teather and by Drs. P. Marks and P. A. Emerson (27 July, p. 230 and 232 respectively), is of considerable interest. At first sight the difference appears surprising, but it accords with a number of similar observations in other comparable circumstances. Avery Jones, Pygott, and I,¹ for example, found that gastric ulcers healed less well in non-smokers than in smokers who stopped smoking (61% average healing in four weeks against 83%), and Badaracco and Vessey² found that women admitted to hospital with "idiopathic" venous thromboembolism who had not been taking oral contraceptives were more likely to relapse than women who had been taking them but subsequently stopped

(36% within three years compared with 8%). Finally, we may note that smoking does not appear to be a cause of venous thrombosis.³

These results agree, therefore, in indicating that patients who have a probable cause for their disease fare better, once that cause has been removed, than patients whose condition has no such removable cause. Dr. Handley and his colleagues suggest that among those who develop myocardial infarction non-smokers tend to be intrinsically more susceptible to both arterial and venous thrombosis. In the light of the above observations their suggestion seems attractive.—I am, etc.,

RICHARD DOLL

Radcliffe Infirmary,
Oxford

¹ Doll, R., Jones, F. A., and Pygott, F., *Lancet*, 1958, **1**, 657.

² Badaracco, M. A., and Vessey, M. P., *British Medical Journal*, 1974, **1**, 215.

³ Vessey, M. P., and Doll, R., *British Medical Journal*, 1969, **2**, 651.

Endometriosis and Pseudopregnancy

SIR,—I think your leading article, "Endometriosis and Pseudopregnancy" (29 June, p. 682) underestimated the usefulness of hormone therapy. Andrews and Larsen,¹ whom you quote, treated only one-seventh of their patients by pseudopregnancy alone. Though in the much larger group treated by surgery alone results in terms of subsequent pregnancy were better than in the small group treated only with hormones they do not clearly indicate the reasons for their choice of treatment, and this selection pattern may have had a considerable effect on the results.

In a small personal series of 77 cases reported in 1969² there were 20 women who desired pregnancy: 14 of them became pregnant, some of them twice, after infertility which in three cases had been present for eight, nine, and 11 years respectively. The percentage success in this admittedly small group was materially higher than that quoted by Andrews and Larsen for their patients treated surgically. Subsequent surgery has so far been necessary in only two of the 77 cases in our series. One of these was a woman aged 23 with an extensive endometriosis of the uterosacral area causing dyspareunia so severe that she had to abandon intercourse. After treatment with norethisterone acetate and ethinyl oestradiol for nine months the mass in the pouch of Douglas was greatly diminished and her pain disappeared. She became pregnant and was delivered of a healthy child. A year later her pelvic mass and her dyspareunia recurred. Once again pseudopregnancy for 12 months was followed by such improvement that a second pregnancy occurred and was successfully concluded. She remained symptom free for a further year, but required hysterectomy at the age of 30 because severe symptoms reappeared. I am confident that conservative surgery in this patient would have been valueless and that a primary surgical approach would have deprived her of all hope of having children. We have not shared Page's³ experience that conservative therapy for endometriosis may be followed by prolonged anovulation. All the pregnancies in our series occurred within six months of the completion of treatment.

Even if, as Andrews and Larsen found, the need for subsequent operation is not