who later required oxytocin to accelerate progress did not differ significantly from bilirubin levels in those infants whose mothers entered labour spontaneously and did not subsequently need oxytocin. The findings of Dr. Gould and colleagues only confirm these observations. What we did record, however, were raised serum bilirubin levels in the infants of mothers whose labours had been artificially induced by amniotomy and followed immediately by oxytocin infusion. Because these mothers received higher doses of oxytocin for more prolonged periods than mothers in whom oxytocin was used only to accelerate the progress of labour, it was felt that this possible effect of oxytocin on neonatal jaundice was dose-dependent. However, an alternative explanation was also suggested—namely, that this association between higher bilirubin levels and maternally administered oxytocin was a spurious one and that instead the infants of mothers who had not entered labour spontaneously were perhaps peculiarly susceptible in some other way to neonatal jaundice. The effects of other drugs administered to these mothers before and during labour and differences in the corticosteroid status of their infants² were suggested as perhaps having a role.

It would be interesting if Dr. Gould and his colleagues could provide data relating neonatal jaundice to oxytocin infusion when this has been used to induce rather than to expedite labour.—We are, etc.,

D. P. DAVIES

Department of Child Health,
Welsh National School of Medicine,
University Hospital of Wales,
Cardiff

R. GOMERSALL

Department of Obstetrics and Gynaecology,
Princess Mary Maternity Hospital,
Newcastle upon Tyne

- ¹ Davies, D. P., et al., *British Medical Journal*, 1973, 3, 476.
- ² Murphy, B. E. P., *American Journal of Obstetrics and Gynecology*, 1973, 115, 521.

Mycoplasma pneumoniae Infection

SIR,—Dr. N. D. Noah, in his article on *Mycoplasma pneumoniae* infection in the United Kingdom in 1967-73 (8 June, p. 544), states that "the full epidemiological pattern of infection in larger communities, as, for example, in the population of a country, is not known." May I draw your attention to a study on the incidence of *M. pneumoniae* infection in Denmark from 1958 to 1969?¹ This study encompasses the population (about five million) of a whole country. During the 12-year period three epidemics of *M. pneumoniae* infection occurred at intervals of four and a half years, the third one culminating in the winter 1967-8. In a follow-up study² it was shown that the next epidemic in Denmark occurred from the winter of 1971-2 to that of 1972-3. Again, the middle of the epidemic was four and a half years after the previous one.

The curve in the figure in Dr. Noah's article depicts the reported cases in the U.K. The top of the curve in the winter of 1967-8 might well be somewhat delayed owing to an underrepresentation of cases, especially for 1967 when the reports, starting in May, were presumably not complete. Unfortunately, the total number of four-weekly reports of respiratory viruses and related organisms are not given. Thus the possi-

bility cannot be excluded that the interval between the two epidemics described for the U.K. might be four and a half years. The seasonal variation in *M. pneumoniae* infection in the U.K. found by Dr. Noah is in accordance with that described for Denmark.^{1,3}

The previous report¹ and the present information from Denmark, Finland, and Norway of *M. pneumoniae* epidemics in 1972² indicate that these epidemics run a concomitant course in the Scandinavian countries. Comparison of data from more European countries might add to our knowledge of the epidemiology and ecology of *M. pneumoniae* infection.—I am, etc.,

K. LIND

The Mycoplasma Laboratory,
Statens Seruminstitut,
Copenhagen, Denmark

- ¹ Lind, K., *Acta Pathologica et Microbiologica Scandinavica*, 1971, B79, 239.
- ² Lind, K., *Mycoplasma pneumoniae Infection. Serological Aetiological and Epidemiological Studies*, Chap. vi, 1973. Copenhagen, F.A.D.L.'s Forlag.
- ³ Lind, K. To be published.

Death during Dentistry

SIR,—The report by your legal correspondent (10 August, p. 419) contains a statement by Mr. Bryan Thomas that during his hospital training there had been no tuition in the use of intravenous anaesthesia. This oversight should be a matter of grave concern for all responsible for the training of dental students.

Ever since the days of Riggs and Wells dentists have been giving anaesthetics. Many of those when practising under the Dentists' Act 1921 displayed in their surgeries a diploma from the Society of Dental Anaesthetists stating that the holder was a competent anaesthetist for dental work. Dental surgeons with a qualifying diploma granted by the Royal College of Surgeons also frequently administered nitrous oxide and oxygen to their patients as a sedation during painful procedures. Even the introduction of local analgesia had many opponents; but the public voted with their feet in no uncertain terms for both these advances in pain control.

The dental surgeon, unlike his medical colleagues, is not compelled to do a pre-registration hospital course. During his course the medical student has as much clinical work as he wants in any branch that interests him. Not so for the dental student. All the more important therefore to ensure that during the undergraduate days the groundwork be as thorough and comprehensive as possible. The control of the airway and the importance of oxygen can never be overstressed. The basic principles of anaesthesia do not change just because a drug is administered intravenously instead of by inhalation, though the various stages are speeded up. All this should be made crystal clear during the most formative years. Post-graduate tuition later, if the dental surgeon is still that interested, will have a firm foundation. No dental surgeon should ever be able to reproach his Alma Mater for being inadequately prepared.—I am, etc.,

M. W. P. HUDSON

University College Hospital Dental School,
London W.C.1

Catecholamine Excretion Levels in Neuroblastoma

SIR,—The recent report of a retrospective study of 487 children with neuroblastoma (3 August, p. 301) draws attention to some of the prognostic factors in this disease, in particular age at diagnosis, clinical stage of the tumour, and histological grade. However, no mention is made of catecholamine levels. It may be that in the period 1962-7 routine estimation of urinary levels of vanillyl-mandelic acid (V.M.A.) were not being carried out.

In the series of 50 patients with neuroblastoma who were seen at the Great Ormond Street Hospital for Sick Children during the period 1970-3 the initial V.M.A. level was raised in 89% of all patients in whom it was measured (41/46). The actual level of V.M.A. excreted in 24 hours varied enormously in different patients and was not correlated to survival. However, in 91% (11/12) children who have survived disease-free the V.M.A. returned to normal following treatment and has not been raised again.

Gitlow *et al.*¹ recently drew attention to the fact that a good prognosis is associated with a rapid return of catecholamine metabolism to normal following therapy. Serial determinations are useful both in assessing response to treatment and when considering the need for further radiotherapy or chemotherapy.² While most children who die from neuroblastoma will do so within two years of diagnosis there are occasional late relapses,^{3,4} and we have recently seen two patients in whom a relapse has been associated with a persistently raised V.M.A. level for up to three years after initial therapy in the absence of any other clinical evidence of active disease.—I am, etc.,

JANE V. BOND

Hospital for Sick Children,
Great Ormond Street,
London W.C.1

- ¹ Gitlow, S. E., et al., *Cancer*, 1973, 32, 898.
- ² Liebner, E. J., and Rosenthal, I. M., *Cancer*, 1973, 32, 623.
- ³ Konrad, P. N., Singher, L. J., and Neerhout, R. C., *Journal of Pediatrics*, 1973, 82, 80.
- ⁴ Jaffe, N., *Journal of Pediatrics*, 1973, 82, 1094.

Age Limit for Contraceptives

SIR,—Dr. Elizabeth H. Gregson (17 August, p. 470) may not be aware that when an invitation to attend the F.P.A. Clinic Doctors' National Council was given to me in November 1972 by an F.P.A. administrative official I asked for a "doctors only" meeting. This was not acceptable to the administration and therefore I declined the invitation.

In September 1970, as Dr. Gregson will doubtless recall, an administrative document suddenly appeared from the F.P.A.'s National Executive. This included an "instruction" that any F.P.A. doctors' meeting must have an F.P.A. administrator in attendance. This attempt by the F.P.A. administration to deny clinic doctors' groups free discussion (which could include criticism of F.P.A. policy) and the opportunity to put independent resolutions forward based on such criticism was not conducive to giving doctors confidence in the F.P.A.'s democracy. There was an element of "Big Brother is watching you" in all this, which was unpleasant.

It is rather surprising that concern for the