of glucagon in 10 ml of diluent. In one patient rapid injection was followed by a severe syncopal attack with nausea and vomiting, but no ill effects followed the slow administration of glucagon to this or any other patient. There was no clinically detectable difference between the results of intermittent injection at four-hourly intervals and continuous infusion. Infusion, however, was less disturbing to the patient and easier for the nursing staff. Patients who continued to take bran, as advised, had no recurrence of symptoms of diverticular disease.

CONVENTIONAL TREATMENT

Response to conventional treatment though ultimately satisfactory is slow. The early days, during which response to treatment is barely perceptible, are trying to the patients and a cause of anxiety to their attendants. The principal symptoms—namely, pain, disturbance of bowel function, and abdominal distension—persisted in the patients selected for comparison for periods of 1.5–6 days (mean four days) after beginning hospital treatment, during which analgesics and sedatives were required. The characteristic tender colonic mass gradually diminished over an average period of five days.

Discussion

The relief afforded by glucagon to 20 consecutive patients suffering from severe symptoms of acute diverticulitis encourages us in our belief that a new therapeutic role for this hormone may have been discovered. This belief is strengthened by the fact that glucagon acted so quickly on symptoms which had often persisted over long periods and resisted other treatment. The rapid resolution of abdominal guarding or rigidity, pyrexia, and leucocytosis—even when these were severe enough to raise the suspicion of pericolic abscess—tends to underline the importance of colospasm in the aetiology of acute diverticulitis, a

theory propounded on histological grounds by Morson (1963). During the study glucagon was given to every patient in whom the diagnosis of acute diverticulitis, or colospasm, seemed to be a reasonable possibility. As other conditions may cause similar symptoms or occur in conjunction with diverticulitis, not surprisingly in four patients the original clinical impression proved erroneous. The fact that lasting beneficial effects occurred only in patients suffering from uncomplicated acute diverticulitis suggests that glucagon may have a specific action in this condition by its inhibitory action on intestinal motility. If this suggestion is confirmed the hormone may prove to be of diagnostic as well as therapeutic value.

Comparison with a group of patients treated conventionally but selected retrospectively may be open to criticism. We therefore plan to conduct a double-blind comparative study between glucagon and a placebo, which we consider justified on the basis of our data.

We thank the research committee of the Welsh Hospital Board for generous support and Dr. G. L. Smith, of Eli Lilly and Company Limited, Basingstoke, Hants for valuable advice and help. Mr. Rushton of the Dylade Company Ltd., Runcorn, Cheshire kindly loaned a Sage pump, model 240, for the investigation. We thank Mr. R. J. Marshall, director of the department of medical illustration, United Cardiff Hospitals, for the illustrations; Mr. Philip Wade, librarian of the Royal Society of Medicine for valuable help with the references, and Miss Iona Williams for secretarial help.

References

- Dotevall, G., and Kock, N. G. (1963). *Gastroenterology*, **45**, 364.
- Knight, M. J., Condon, J. R., and Smith, R. (1971). *British Medical Journal*, **2**, 440.
- Lefebvre, P. J., and Unger, R. H. (1972). *Glucagon*. Oxford, Pergamon Press.
- Morson, B. D. (1963). *British Journal of Radiology*, **36**, 385.
- Necheles, H., Sporn, J., and Walker, L. (1966). *American Journal of Gastroenterology*, **45**, 34.
- Painter, N. S. (1962). M.S. Thesis, London University.

MEDICAL MEMORANDA

Case of Post-primary Tuberculosis and Massive Haematemesis

B. W. HANCOCK, D. B. BARNETT

British Medical Journal, 1974, **3**, 722–723

Tuberculous aortitis is extremely rare and the prognosis is grave, most cases progressing to aneurysm formation and rupture (Volini *et al.*, 1962). The case reported here is unique in that the patient had post-primary cryptic disseminated tuberculosis (Proudfoot *et al.*, 1969) presenting with massive haematemesis from a tuberculous aorto-oesophageal fistula without aneurysm formation.

Case History

The patient, a man aged 52, was admitted to this hospital as an emergency case. He had vomited fresh blood of unknown quantity

passed 2 pints (1,140 ml) of fresh melaena. As he was confused a history was obtained from the relatives. He had been unwell for 12 weeks with anorexia, weight loss, and upper abdominal pain. Other than a daily intake of 6 pints (3.4 l.) of beer there was nothing else relevant in the history.

On admission he was found to be shocked and there was the appearance of recent weight loss. The abdomen was soft with some tenderness in the epigastrium. No masses were palpable and rectal examination showed fresh melaena. The preliminary differential diagnosis included bleeding from peptic ulcer, gastric neoplasm, and oesophageal varices. Haemoglobin was 73% and he was transfused with 4 units of blood. Next morning he remained confused though the bleeding had stopped. Endoscopy showed some changed blood adherent to the mucosa near the cardia of the stomach. The oesophagus, stomach, and proximal duodenum appeared normal.

The following investigations gave normal results: blood sugar, bleeding and clotting values, serum vitamin B₁₂ and folic acid, serum proteins and electrophoresis, and barium swallow, meal, and follow-through examination. A portable chest x-ray film on admission showed miliary nodularity thought to be industrial in origin. This was not seen in later films. Liver function tests showed slightly raised transaminases and the platelet count was 70,000/mm³. In view of the patient's continued confusion, previous alcohol intake, and abnormal liver function liver disease was considered despite the absence of demonstrable varices. He improved over the next few days though the fever remained. The platelet count returned to normal and it was intended to perform liver biopsy, but the patient collapsed again 12 days after admission with further haematemesis and melaena. He was rapidly transfused with 2 units of plasma and 4 units of blood. His condition worsened and one

Section of Therapeutics, Academic Division of Medicine, Royal Infirmary, Sheffield S6 3DA

B. W. HANCOCK, M.B., M.R.C.P., Registrar
D. B. BARNETT, M.B., M.R.C.P., Lecturer

hour later there was a cardiac and respiratory arrest due to massive haematemesis. Resuscitation was unsuccessful.

NECROPSY

In the mediastinum a dense white mass 2 by 1 cm was found through which a fistulous communication existed between the aorta opposite the origin of the left subclavian artery and the oesophagus (fig. 1). There was fresh thrombus in the fistula. The trachea and main bronchi were normal but there were small pleural effusions and some apical fibrosis bilaterally. The stomach was dilated and full of blood. A 6-cm tear was present at the cardia and smaller

FIG. 1—Necropsy specimen showing aorto-oesophageal fistula.

FIG. 2—Necropsy specimen showing gastric tears.

tears in the mucosa were noted along the lesser curve (fig. 2). No gastric or duodenal ulceration was present.

Histologically the white mass in the mediastinum was a lymph node largely replaced by caseous tissue with tuberculous follicles. Scattered caseating tuberculous lesions were present in the lungs and one of the apical foci of scarring showed evidence of reactivation. Acid-fast bacilli were identified in another mediastinal node. Tuberculous granulomata were also found in the liver.

In summary the findings were of an aorto-oesophageal fistula produced by erosion from caseous tuberculous mediastinal lymph nodes, active pulmonary and miliary tuberculosis, and probable terminal Mallary-Weiss's syndrome.

Comment

Only four cases of tuberculous false aneurysm of the thoracic aorta with rupture into the oesophagus have been recorded (Stiefel, 1958). We have been able to find only two reports of tuberculous aorto-oesophageal rupture without aneurysm formation (Ferrabouc and Fauvert, 1931; Hong and Sugai, 1965) and we believe that the present case is the first to be recorded of post-primary cryptic disseminated tuberculosis presenting with haematemesis from a tuberculous aorto-oesophageal fistula without aneurysm.

In retrospect the patient's history contained certain features which might have allowed a diagnosis to have been made in life. He had been unwell and losing weight for 12 weeks before admission. After his initial haematemesis he remained ill and feverish despite transfusion. Hepatic disease was suspected from the abnormal liver enzyme values and initial thrombocytopenia. Liver biopsy was considered but delayed because of the low platelet count. Had this investigation been performed the diagnosis might have been made in life, as 75% of patients with miliary tuberculosis have positive liver biopsy findings (Brunner and Haemmerli, 1964). The initial overpenetrated portable chest x-ray film showed miliary nodularity thought to be industrial in origin. This nodularity was not noted on a later film but the initial findings were probably significant in view of the scattered pulmonary tuberculous deposits found at necropsy.

Tuberculous aortitis, especially with perforation, carries a grave prognosis. Few instances of successful surgery in such cases have been recorded (see, for example, Rob and Eastcott, 1955) and chemotherapy has little effect (Volini *et al.*, 1962). Haematemesis in a patient with tuberculosis should suggest aorto-intestinal rupture even though peptic ulceration is commoner in these cases (Citron, 1973). Prompt surgical reaction of the lesion offers the only possible chance of survival.

We are grateful to Professor R. Kilpatrick for permission to report details of this patient, and to Dr. K. Scott, department of pathology, for the necropsy report and photographs.

References

- Brunner, K., and Haemmerli, U. P. (1964). *German Medical Monthly*, 9, 372.
- Citron, K. M. (1973). *Medicine*, 14, 911.
- Ferrabouc, L., and Fauvert, R. (1931). *Annales d'Anatomie Pathologique*, 8, 764.
- Hong, L. C., and Sugai, K. (1965). *Singapore Medical Journal*, 6, 164.
- Proudfoot, A. T., *et al.* (1969). *British Medical Journal*, 2, 273.
- Rob, C. G., and Eastcott, H. H. G. (1955). *British Medical Journal*, 1, 378.
- Stiefel, J. W. (1958). *Archives of Pathology*, 65, 506.
- Volini, F. I., *et al.* (1962). *Journal of the American Medical Association*, 181, 78.