notes the substantial benefit many children with moderate to severe chronic asthma derive from the use of steroid aerosols. However, it fails to emphasize the important place that regular use of bronchodilators has in management.

It is inferred that if adequate control of asthma is not achieved with regular prophylactic disodium cromoglycate and bronchodilator therapy given early in the course of any acute wheezing attack steroid therapy in one form or another should be used. As about 0.5% of children have chronic airways obstruction as a result of asthma^{1,2} and this is often inadequately controlled with regular disodium cromoglycate and intermittent bronchodilators, steroids appear to be recommended for a large number of children. This is despite the well-known complications of oral corticosteroids and the lack of information of the long-term effects of topical steroids on the human lung.

In fact many children with moderate chronic asthma whose airways obstruction persists despite regular disodium cromoglycate and intermittent bronchodilator therapy will be adequately controlled with regular bronchodilator therapy. The effectiveness of regular oral theophylline in full therapeutic dosage is now well documented.³ While the value of combining one of the newer β -adrenergic stimulants either orally or by inhalation with regular theophylline is not proved, we have used this combination for many years and have achieved very adequate control in a substantial number of children with moderate to severe chronic asthma without any apparent side effects.⁴ Orciprenaline or salbutamol by inhalation using a hand nebulizer driven by small air compressor pump as described in detail elsewhere⁵ is particularly valuable both for the control of persisting airways obstruction and for the rapid relief of acute exacerbations.

Certainly no child should be started on corticosteroids, either by inhalation or orally, until it has been clearly demonstrated that the combination of regular disodium cromoglycate, oral theophylline, and orciprenaline or salbutamol by inhalation has failed to give adequate control. The criteria for control outlined in your article are adequate, but in older children objective evidence from pulmonary function testing is also of considerable importance. Some parents of children with chronic asthma fail to appreciate the severity of the condition and may report that their child is normal whereas he has substantial persisting airways obstruction. Conversely there are a small number of children, usually from disturbed backgrounds, who on reporting by parents seem to be having a great deal of trouble with their asthma, but this is not supported by objective data. There is a substantial risk of undertreating the former group and overtreating the latter if history alone is the guide to therapy.—I am, etc.,

PETER D. PHELAN

Department of Thoracic Medicine,
Royal Children's Hospital,
Melbourne

Motor Insurance and Ischaemic Heart Disease

SIR,—Some interesting points have been raised by Dr. G. J. Rockley (22 March, p. 679) concerning the underwriting of motor insurance where the medical condition of the policy-holder is relevant.

Motor insurance contracts are governed by the principle of utmost good faith—that is to say that there is a requirement on the part of the policy-holder to notify his insurer of any fact which would materially alter the risk. This would include any change in health affecting the ability to drive advised to the patient by his doctor. Insurers would always expect a doctor to advise a patient on the extent and nature of his future activities relative to his medical condition, whether his advice is sought specifically or not. It remains as a matter for the patient's own conscience and common sense as to whether he accepts or rejects the doctor's advice. In the case of illness or some form of disability insurers must defer to the opinion of a registered medical practitioner on the question of the policy-holder's ability and suitability to drive.

In these circumstances it is vital for a patient to understand and appreciate the fact that if he fails to advise his insurers of a change in his medical condition then he would certainly run the risk of having a claim for damage to property (including damage to his own car) repudiated. In so far as personal injury claims are concerned the Road Traffic Act deals with this situation—particularly section 148.

When an insurer requires a report from a doctor concerning fitness to drive, it is normally expected that the report will come from the patient's general practitioner, and it would be assumed that the G.P. would be in possession of all facts concerning the case. In the case of ischaemic heart disease, where fitness to drive may improve over the months after clinical recovery, a certificate of fitness to drive would be required by the insurers from the patient's doctor stating that the condition had improved to the extent where driving a vehicle would not be abnormally hazardous. Equally, the insurers would be expected to be notified of any decline in the patient's condition which would affect fitness to drive. Insurers require only an affirmative opinion from a medical practitioner regarding the patient's ability to drive. The extent to which a doctor feels it necessary to examine a patient and to consult with specialists, etc., is entirely a matter of opinion for the doctor in individual cases. It follows from this that the fee which a doctor charges a patient for such a professional opinion would reflect the amount of work, time, and consultation necessary for him to produce such an opinion.

Because they have a duty placed upon them to provide insurance cover for the motoring public insurers feel that they must satisfy themselves at all times regarding the level of fitness of a driver, and it is to this end that it is necessary for them to require their policy-holders to seek medical advice. In many cases, while insurers are prepared to provide insurance cover they may wish—and indeed they have the right—to impose special terms and to charge higher than normal premiums.

With regard to the certificate of ability to drive mentioned in Dr. Rockley's letter, may I say that this certificate, which is couched

in the affirmative form, has been the subject of discussion and agreement between insurers and the B.M.A.—I am, etc.,

R. C. W. BARDELL
Secretary General,
British Insurance Association

London E.C.4

Prophylaxis of Postoperative Deep Vein Thrombosis

SIR,—We are obliged to Mr. F. S. A. Doran (24 May, p. 442). He has done as much as anyone else to popularize electrical calf muscle stimulation for the prevention of postoperative deep vein thrombosis. We also thank Mr. J. A. Lewis (5 July, p. 41) for pointing out the superiority of random number tables (which we now use). The method of randomization in the published trial (22 March, p. 649) based on the month of birth of the patient produced an unfortunate excess of patients in the control group. This made the figures clumsily uneven but it can surely not be claimed that it was anything but strictly random or that the method of prophylaxis used was determined by our preference and not purely by chance.

Mr. Doran's labour in calculating χ^2 of the stimulator group against the heparin group has been largely wasted. He has added together laparotomies for benign disease, laparotomies for malignant disease, and operations on the bladder and prostate and has produced a χ^2 of 4.924—favouring heparin with a probability of over 95%. If he had taken laparotomies for benign disease alone and compared the two groups he would have found a χ^2 of 0.875 (no significant advantage of heparin over the stimulator). In the laparotomies for malignant disease on the other hand χ^2 is 13.298 and the chance that heparin is no better than electrical stimulation is less than one in 1000.

He makes another statistical mistake twice. First, in comparing major deep vein thrombosis (D.V.T.) between the two groups. The total number of cases of major D.V.T. in the stimulator group was five and in the heparin group one. Add these together and divide by two and the total expected number is three. No statistician is prepared to apply the χ^2 test (or indeed any other test) when the expected number is less than four. Similarly, four haemorrhages in the heparin group against none in the stimulator group gives an expected number of two and, again, cannot be analysed statistically. To put it another way, if you toss a penny four times there is a strong possibility that it will come down heads on each occasion.

Twenty-two patients were withdrawn from the trial. There were four deaths, two in the stimulator group, one in the control group, and one in the heparin group. Their details are as follows: (1) Stimulator. A man of 80 died two days after laparotomy, suture of a caecal perforation, and transverse colectomy for an obstructing carcinoma of sigmoid colon. Overwhelming sepsis. No necropsy. (2) Stimulator. A man of 76 died two days after gastrectomy, splenectomy, and transverse colectomy for advanced carcinoma of the stomach. Superior mesenteric artery occlusion. No necropsy. (3) Control. A woman of 71 died three days after gastrectomy and transverse colectomy for gastric reticulosis. Cardiac and respiratory failure. No necropsy. (4) Heparin. A man of 74 died four days after right hemicolectomy for a perforated carcinoma of the caecum. Uraemia and bronchopneumonia. Necropsy showed no pul-

¹ McNicol, K. N., and Williams, H. E., *British Medical Journal*, 1973, 4, 7.

² Hill, D. J., et al., *Archives of Disease in Childhood*, 1972, 47, 874.

³ Weinberger, M. M., and Bronsky, E. A., *Journal of Pediatrics*, 1974, 84, 421.

⁴ Williams, H. E., and Phelan, P. D., *Respiratory Illness in Children*, p. 116. Oxford, Blackwell Scientific, 1975.

⁵ Phelan, P. D., and Stocks, J. G., *Archives of Disease in Childhood*, 1974, 49, 143.

monary embolus and no more bleeding at the operation site than would have been expected.

Of the 14 patients withdrawn because of failure to observe the protocol, one was in the heparin group (the preoperative dose was forgotten) and the remaining 13 in the stimulator group: in seven of these the surgeon omitted to use the machine and in six the machine failed to function.—We are, etc.,

I. LAWRENCE ROSENBERG

St. James's Hospital,
Leeds

MARY EVANS
A. V. POLLOCK

Scarborough Hospital,
Scarborough, N. Yorks

Survival after Postoperative Avascular Necrosis of Lesser Curve of Stomach

SIR,—I read with interest the paper by Dr. J. F. Halvorsen and others (14 June, p. 590) reporting the third death due to avascular necrosis of the lesser curve of the stomach following highly selective vagotomy (H.S.V.) for duodenal ulcer. I should like to describe a similar case in which the patient survived.

An otherwise healthy male patient aged 46 with a long history of chronic duodenal ulceration underwent H.S.V. on 28 November 1974. His condition remained satisfactory until the third postoperative day, when he developed left lower chest pain associated with fever, mild hypotension, and a pleural friction rub. It was assumed that he had developed a chest infection and he was treated accordingly with physiotherapy, systemic antibiotics, etc. Several hours later his condition deteriorated: his blood pressure fell, abdomen became distended, and bowel sounds were absent. An abdominal tap was performed and the fluid obtained was mistakenly thought to be blood. Intraperitoneal haemorrhage was therefore diagnosed and appropriate resuscitation was carried out. Once a suitable blood pressure was attained laparotomy was carried out, when a large defect was found in the lesser curve and about 2½ l of blood-stained fluid, mostly gastric contents, in the peritoneal cavity. The defect was sutured and peritoneal toilet carried out. The patient's postoperative condition was satisfactory but he subsequently developed a left subphrenic abscess which required treatment with drainage and antibiotics. He was discharged on 10 January 1975 and when last seen at the clinic in May 1975 he was very well, had no recurrence of dyspepsia, and was prepared to go back to work.

This case also demonstrates the difficulty in diagnosis of this rare condition, which nonetheless should be thought of if deterioration occurs after H.S.V.

I should like to thank Mr. G. E. Schofield for permission to report this case which was under his care.

—I am, etc.,

STUART W. BRODIE

Law Hospital,
Carlisle, Lanarkshire

G.M.C. and Indian Qualifications

SIR,—Mr. A. K. Varshneya (5 July, p. 43) writes that there is an increasing number of teaching schools in India with poorly equipped departments and that the standard of medical education has gone down in recent years. I spent nearly 13 years in the Indian Medical Service, so that I know the background, and I have been back twice, in 1969-70 and 1972, touring the Indian states on behalf of the World Health Organization. During these tours I visited a large number of primary health centres and district and

general hospitals and some teaching hospitals. Though I saw in the primary health centres many examples of Indian doctors supplying a cheerful and unselfish service to their countrymen, the lack of even the simplest equipment and the elementary nature of the procedures in these centres was striking. In the district and general hospitals laboratory equipment was of the scantiest and in the teaching hospitals there was a conspicuous lack of the sort of modern equipment one would have expected to find there.

In the hospitals the few laboratory tests attempted were carried out usually in one room and there was a grave lack of adequately trained technical staff. I noticed that doctors generally had a knowledge of only the simplest techniques such as blood cell counting, examining malaria slides, or examining faecal specimens for helminth ova, and while one could argue that medicine could possibly be practised in rural India without a knowledge of laboratory procedures, such inadequately trained doctors would not be suitable for participating in a highly technical hospital service such as Britain's.

No doubt some graduates of the good Indian medical schools display a wider knowledge of medical practice, but those of us who know India well are aware that there are a great number of doctors coming to this country who are simply not well enough trained. It is absolutely imperative that there be a sorting-out examination, and no matter how disappointing its results may be for some candidates an exacting standard must be maintained.

It would be futile to run a three-month course (as has been suggested by an Indian doctor on the radio from Birmingham) to prepare doctors who have arrived in the U.K. to take the G.M.C. examination. How could a three-month habilitation course, especially for someone whose English is weak, make up for the deficiencies of several years in an inferior medical school?—I am, etc.,

R. J. HENDERSON

Public Health Laboratory,
Royal Infirmary,
Worcester

Sodium Nitroprusside in Anaesthesia

SIR,—We should like to amplify your brief statement (7 June, p. 524) on the metabolism of sodium nitroprusside (SNP).¹

In patients infused with SNP we have found that there was an increase in blood cyanide (HCN) levels, even with small doses of the drug. Of this HCN, 98% was present in the red cells. Though β -mercaptopyruvate sulphur transferase occurs in the red cells of the rat,² it has not yet been shown to play any significant role in the direct detoxication of HCN in human erythrocytes. The blood HCN is slowly detoxicated to thiocyanate (SCN) by tissue rhodanese so that plasma SCN levels showed only a gradual rise. The ratio SCN:HCN in body fluids is normally around 1000:1 and so at the relatively low levels of SCN attained during hypotensive anaesthesia using SNP there would be insignificant formation of HCN from SCN by the ill-defined thiocyanate oxidase system.

You suggest a maximum SNP dose rate of 3 mg kg⁻¹ h⁻¹, but this may be too high. Death due to cyanide poisoning is attri-

buted to inhibition of cytochrome oxidase. In vitro 50% inhibition occurs at a concentration of 1.5 μ mol HCN/l.³ In vivo there will no doubt be a plasma/tissue concentration gradient and so plasma levels at which inhibition would occur would be somewhat higher. In dogs given a single intravenous dose of 1 mg SNP/kg, resulting in peak plasma levels of 3 μ mol HCN/l, we have found evidence for histotoxic hypoxia.

Our studies in man have shown that plasma HCN levels immediately following short-term infusions (over a period of two hours or less) show a linear relationship to the total dose of SNP. A plasma level of 3 μ mol HCN/l would be reached by the infusion of about 1.5 mg SNP/kg.

Deaths have been reported⁴ following SNP infusions in which doses of approximately 4 mg/kg per two-hour period have been given. It therefore seems reasonable to suggest that plasma HCN levels should not be allowed to rise above 3 μ mol/l and the maximum total dose of SNP be set at 1.5 mg/kg for hypotensive anaesthesia of relatively short duration. In the event of untoward symptoms attributable to SNP we would certainly encourage the use of hydroxocobalamin and sodium bicarbonate as you suggest, and in addition sodium thiosulphate.—We are, etc.,

CYRIL J. VESEY
PETER COLE
PETER SIMPSON

Department of Anaesthesia,
St. Bartholomew's Hospital,
London E.C.1

1 Vesey, C. J., et al., *British Medical Journal*, 1974, 2, 140.

2 Sorbo, B., in *Proceedings of the First International Pharmacological Meeting*, ed. B. B. Brodie, et al., vol. 6, p. 21. London, Pergamon, 1962.

3 Schubert, J., and Brill, W. A., *Journal of Pharmacology and Experimental Therapeutics*, 1969, 162, 352.

4 Merrifield, A. J., and Blundell, M. D., *British Journal of Anaesthesia*, 1974, 46, 324.

Emigration of Doctors

SIR,—Dr. M. P. White (7 June, p. 561) writes from his experience as a district medical officer, but many of us who have served from the grass roots up to Ministry levels in many developing countries are unable to agree with his statements. There is no such thing as tropical medicine; it is the medical care of poverty, national and individual; and the paramedical and auxiliary cadres, urban and rural health centres, and a reconsideration of medical student training are all appearing in the so-called developed countries as they face up to the economic problems of medical care.

Dr. White specifically raises four points as some of the "certain facts . . . always ignored" by the protagonists of community health care.

I agree that in the developing countries there is a decline of morale, discipline, and output of the health services paralleling those of the U.K. But the root causes are very different. In the developing countries the indigenous are now taught by indigenous physicians and paramedicals who in turn were taught by expatriates. Thus the indigenous students continue to be taught an inappropriate medical care system. Indigenous training in many areas is still not orientated to a country's individual epidemiological and social needs and priorities. Doctors and paramedicals soon discover this