Aspects of Sexual Medicine

Surgery of Male Sexual Disorders

JOHN P. PRYOR

British Medical Journal, 1975, 3, 585-587

Surgery has a small but important role in correcting sexual disorders of men. Some of the symptoms requiring surgical correction are well recognized, but others are less well known and are often ignored. This article will concentrate on those symptoms which may require urological investigation or treatment.

Painful Coitus

The tight foreskin which causes a painful erection is readily recognizable and easily cured by circumcision. But some patients requesting circumcision, or other simple procedures such as vasectomy, are really seeking advice for an underlying sexual dysfunction, and this should always be suspected when the prepuce retracts easily and does not appear to be tight. Less commonly a tight fraenum may cause the pain, and simple division of the band is all that is required. Paraphimosis can usually be reduced by manipulation, and circumcision may be performed later as an elective procedure. Patients with a "zip fastener" injury to the penis sometimes require surgical intervention to release the prepuce, and circumcision may be required to give a satisfactory cosmetic appearance.

Hyperflexion of the erect penis may rupture the tunica albuginea. It is usually accompanied by a searing pain and occasionally by an audible crack. The pain is followed by detumescence, and the penis becomes oedematous and bruised. Operation is indicated in the acute stage to repair the defect in the tunica albuginea and to prevent subsequent fibrosis and deformity.

Pain in the penile shaft during erection is usually due to Peyronie's disease, and this will be discussed more fully later. It is sufficient to note that the pain associated with Peyronie's disease always improves spontaneously, though the deformity may persist for many years.

A perplexing symptom is one of pain at the tip of the penis at the time of ejaculation. The origin of this pain is difficult to determine, though some patients with it suffer from prostatitis. This condition may also give rise to perineal or testicular discomfort during coitus.

Penile Deformity

Ventral curvature of the penis is a feature of hypospadias. It is usually corrected in childhood, and persisting chordee may

be of little concern until erections occur. In such patients surgical excision of the chordee is indicated. Similarly, epispadias is accompanied by dorsal chordee, and this may require surgical excision. Ventral curvature of the penis may also be due to chordee without hypospadias, a condition termed "congenital short urethra." In some of these patients there is fibrosis of the corpus spongiosus. Surgical treatment lies between lengthening the urethra with an inlay of skin and excising a segment of the tunica albuginea on the dorsal surface of the penis to counteract the relative excess length of the corpora cavernosa.

Dorsal curvature of the erect penis is a feature of Peyronie's disease and the angulation is often severe enough to prevent penetration of the vagina. The aetiology of Peyronie's disease is unknown, but it is characterized by fibrosis in the tunica albuginea, particularly in the septum between the corpora cavernosa. Occasionally calcification and even ossification occur. Patients with Peyronie's disease present with penile deformity, painful erection, and sometimes a lump, which may mistakenly be diagnosed as a neoplasm. Such an error should not occur, as the lump is characteristically within the substance of the penis on its dorsal aspect and is not related to an epithelial surface. Over the course of many months there is a tendency for the deformity to diminish, and various drugs are used in an attempt to speed its resolution. At present a clinical trial is being conducted to assess the relative efficacy of vitamin E and procarbazine. Persistence of the deformity in Peyronie's disease is often accompanied by impotence, a combination particularly difficult to treat. When prolonged medical treatment fails to improve the deformity, surgical excision of the fibrous tissue is indicated. The defect in the tunica albuginea is replaced with a dermal graft taken from the anterior abdominal wall.

Localized scarring of the cavernous tissue, either congenital or the result of trauma, may produce a deformity with angulation of the penis during erection. If the deformity is sufficient to prevent coitus, a surgical attempt should be made to achieve an erect organ by excising an appropriate segment of the tunica albuginea.

Priapism

The initial satisfaction produced by a long-sustained erection soon passes to embarrassment and finally fear. Consequently medical advice is rarely sought during the first 12 hours, and conservative management is indicated only while the patient is being prepared for operation. During that time sickle cell disease and leukaemia should be diagnosed and the additional treatment for these conditions instituted. Surgical decompression of the engorged corpora cavernosa can be achieved by shunting the blood away by anastomosing the corpora cavernosa to the corpus spongiosus or the long saphenous vein. Priapism lasting more than 24 hours is usually followed by impotence. Those patients who were treated with a long-saphenous-vein shunt may be cured by dividing the vein if it is still patent. In other patients a penile implant proves necessary, as the corpora cavernosa have become fibrosed.

Impotence

Impotence is usually psychogenic and its treatment requires counselling of both partners. In some patients reassurance and judicious medication may be sufficient to restore normal potency. The hypothalamic gonadotrophin releasing hormone may promote erectile activity, and its use in impotence is under review.

Organic causes for impotence (table) should be readily diagnosed, and in some of these patients a penile implant may be considered. One method of treatment was to insert a Silastic rod beneath the tunical albuginea, but this was prone to fracture or displacement. It has recently been replaced by a softer model where a separate implant is passed through a perineal incision into each corpus cavernosus.

Ejaculatory Disorders

Normal erection and ejaculation are controlled by both parasympathetic and sympathetic pathways. Their exact relationship to the various phases of coitus is not fully understood, and all aspects are subject to psychogenic influence. Primary ejaculatory failure and most instances of secondary ejaculatory failure are due to psychogenic factors. Failure to ejaculate may be due to sympathetic nervous blockade by drugs or operations such as abdomino-perineal resection of the rectum, lumbar sympathectomy, retroperitoneal node dissection, or vascular operations on the aorta and iliac vessels. These operations produce variable effects. Erections normally persist, but ejaculation is diminished and consists of seminal fluid oozing from the penis rather than the usual forceful expulsion. It should be remembered that the force and efficiency of ejaculation decrease with advancing age, which is why older men take longer to obtain negative sperm counts after vasectomy.

Failure to ejaculate may be due to retrograde ejaculation, and this is common after prostatectomy. It also occurs in other conditions where there is incompetence of the bladder neck. Potency after prostatectomy is usually maintained, and the chances of impotence are little more than those associated with any surgical procedure. Retrograde ejaculation after prostatectomy occurs in almost half the patients and is seldom of concern to the patients once its significance is explained. A urethral stricture may prevent normal ejaculation, but in these patients the disturbance of micturition will be readily apparent and of greater concern.

Finally, in those patients with congenital absence or inflammation of the seminal vesicles the volume of seminal fluid is much reduced. Normally 80% of the ejaculate is produced by the seminal vesicle, and its lack is readily appreciated.

Ejaculatory failure does not require surgical treatment. If infertility is due to retrograde ejaculation, it is necessary to retrieve the spermatozoa from the bladder and use these for artificial insemination of the wife.

Haemospermia

This distressing symptom is rarely due to any serious lesion, and a solitary episode in a young man may be safely ignored. Repeated haemospermia should be investigated by urine analysis, intravenous urography, cystourethroscopy, and possibly vesiculography. Inflammatory and neoplastic diseases of the prostate or seminal vesicles occasionally present with haemospermia, and so may, on even rarer occasions, a bladder neoplasm.

Fertility Problems

Anxiety about unwanted pregnancies or failure to achieve a wanted pregnancy often cause sexual dysfunction. The initial zest of a couple trying to conceive disappears as the years roll by and the duration of infertility increases. It is important to

take an accurate history of coitus in these patients, as occasionally infertility is due to a failure to ejaculate intravaginally.

A clear understanding of the pathophysiology of male infertility has lagged behind the study of infertility in the female or in animal husbandry. In an attempt to improve this a Fertility Investigation Group (F.I.G.) based at the Institute of Urology has been formed. In the past too much importance has been attributed to the sperm count, and it is now realized that infertility is due to lack of sperms only when the count is less than 5-10 million per millilitre of seminal fluid. A full discussion of infertility is not relevant to the present subject, but surgical treatment is required in those patients who have oligospermia associated with varicocele, or in those with obstructive azoospermia. In the latter group of patients the success of operation (epididymovastostomy) depends on the amount of normal epididymis present. Testicular biopsy is recommended for patients with azoospermia or severe oligospermia so that an accurate diagnosis can be made and rational treatment offered.

Male sterilization by vasectomy has proved to be an acceptable method of limiting family size and is usually performed when a couple have decided they do not want more children. In these circumstances the only exception to male sterilization should be when the need for it stems from severe mental disorder in the wife. Bilateral vasectomy can readily be performed under local anaesthesia. The essential functions of the operation are to divide the vas deferens and separate the ends. It is not necessary to excise any of the vas, but it is essential to confirm subsequently on two occasions the absence of spermatozoa from the ejaculate. The formation of a sperm granuloma may give rise to the persistence of spermatozoa in the ejaculate in about 0.5% of patients. Most instances of pregnancies following bilateral vasectomy are not accompanied by a positive sperm count in the husband, particularly when two negative counts have previously been obtained.

Men undergoing vasectomy should always consider it irreversible, but if any unexpected accidents occur advice is sought about reversal. Recent experience suggests that, provided spermatozoa were present at the time of reanastomosis, more than 80% of patients should have spermatozoa in the ejaculate after vasovasostomy, though in only about half of these will a pregnancy ensue. In closing this discussion of vasectomy one should mention the psychological aspects of the procedure. It is inevitable that the operation should be accompanied by psychological changes, but in most people these are minor. The likelihood of notable psychological disturbance varies with the individual patient and his counsellor and the incidence in published reports has varied also with the nature of the survey.

Miscellaneous Procedures

Occasionally young men are troubled by congenital absence or removal of one or both testicles. It is then justifiable to insert a testicular prosthesis. Various Silastic models are available in a wide range of sizes, and they may be inserted into the scrotum.

Penile lengthening is occasionally necessary in patients with severe epispadias or after injuries to the penis. Various techniques are described, but basically they rely on mobilizing the corpora cavernosa from the pubic rami. These procedures are considered only when very little penile shaft remains, as satisfactory intercourse can occur with a short stump. Many men are able to have intercourse after a partial amputation of the penis for tumour, but in others pain is troublesome.

Conclusion

"The pen is mightier than the sword," and this philosophy is certainly true in the management of sexual dysfunction. In the course of this paper I hope that while stressing the importance of the psyche I have been able to draw attention to the relatively

few patients who might benefit from surgical procedures to improve their well being.

Bibliography

- Campbell, M. F., and Harrison, J. H. (eds.), *Urology*, 3rd edn. Philadelphia, W. B. Saunders, 1971.
Horton, C. E. (ed.), *Plastic and Reconstructive Surgery of the Genital Area*. Boston, Little, Brown, 1973.

Organic Causes of Impotence

- Congenital: absence or fibrosis of corpora cavernosa
Traumatic: spinal cord injury, fractured pelvis, surgical (abdominal perineal resection, sympathectomy, cystectomy)
Vascular: priapism, arteriosclerosis (Leriche syndrome)
Neurological: spinal cord lesions, peripheral neuropathy
Endocrine: diabetes, hypogonadism
Pharmacological: parasympathetic blockade, sympathetic blockade

Letter from . . . South Australia

A Dean Down Under

PHILIP RHODES

British Medical Journal, 1975, 3, 587-588

South Australia is a progressive state. It has a Premier who seems to be an intelligent intellectual. He heads a Labour government, which therefore has some accord with the central Commonwealth Government of Mr. Gough Whitlam. Our premier, Mr. Don Dunstan, known as "Donnie," is a lawyer who was able to devote an hour's television programme to poetry and his delight in it, and he recited it too. It was a great success. Recently he intervened in the Maslins Beach episode.

For many years Maslins Beach has been the haunt of illegal nude sun and sea bathing. The police had in general turned a blind eye or at most made an arrest when they had time from their other more arduous duties. The beach is about 20 miles (32 km) south of Adelaide and, like the whole long coastline, is relatively devoid of people. Then during a lull in the news the local paper started a story on Maslins. The police had to take some action and arrested a few offenders. Then a stipendiary magistrate, taking his duties seriously, announced that he would punish to the maximum the law allowed and gaol for three months those guilty of this heinous offence. Our Donnie stepped in and the law was changed almost overnight. Nude disporting at Maslins became legal, and this was the first beach in Australia to be so licensed. Now there is also another.

After the brouhaha in the paper, cars blocked the road on the cliff-top overlooking Maslins, apparently filled with people sporting binoculars and telescopes, and light aeroplanes, of which there are many, used the beach as a reporting-in station. The controllers of Adelaide airport are reputed to have played the game by allowing these daring pilots to come in to land at 500 feet (152 m). It was all less than a nine days wonder, though householders in the area became anxious that the value of their properties might fall. They obviously had little confidence that there might be enough voyeurs in the country to raise the asking price.

Community Medicine

Community medicine and community practice are all the rage in Australia. The Commonwealth Government is convinced of their value and has given generously of taxpayers' money to set

up chairs of community medicine and health centres. Nevertheless, the Government, and many people, are confused about the difference between community medicine and practice. They make few distinctions. What they usually mean, when they appear to think about the problem at all, is family practice, or the delivery of primary medical care.

There is little understanding of the population as a patient, using the disciplines of epidemiology, statistics, and sociology. Perhaps community medicine is thought to be done by government teams involved in health care resources and is not seen as an academic discipline. This could well lead to confusion on the international scene since the holders of chairs of community medicine could well be general practitioners, so causing some difficulties in communication. But there are signs that the problem is being recognized and embarrassment may be avoided. Both disciplines are obviously needed but their aims are different, so it is well to keep some lines of distinction between them.

The teaching health centres in Adelaide are just getting under way, with Government support. They are superb. Purpose built and lavishly equipped, they are sited in shopping centres, which are the hub of life for small communities. The suburbs are spread out and nearly every house is on its own plot of land, so that the populace is relatively thinly spread. But nearly all have cars and the women in particular whizz in and out of the shopping centres, which cater for most of their needs. A health centre at St. Agnes is staffed by general practitioners whose consulting rooms are plush by U.K. standards. Wood panelling and carpets abound and the medical equipment leaves little to be desired. There is a room for a physiotherapist, a treatment room, and diagnostic radiological apparatus. Close by are dental services and a chiropodist. Behind the general practice suite is the consultants' suite, where visiting specialists come regularly. The pathology service is excellent and run centrally by the Institute of Medical and Veterinary Science associated with the Royal Adelaide Hospital. (I will say more of these on some future occasion.) The Mini vans of the I.M.V.S. are to be seen all about the city, labelled "Pathology Collection Service," bringing specimens to the institute.

Health centres are staffed, apart from the doctors, by nurses and secretaries, social workers, clinical psychologists, counsellors, and, indeed, all who nowadays are essential for the full delivery of health care. Visiting patients in their homes is being built up, and all who work in these centres are greatly enthusiastic, and this alone will ensure their success. The patients' reaction is harder to assess but seems very favourable. At the least they may be saved a long journey to a centrally situated hospital. The newer immigrants from Europe, however, are used to going to hospital for their primary care, for a variety of reasons, and