addition to calcium sensitising properties. EMD 53998 seems to be the first compound in which calcium sensitising properties predominate.^{40 41}

Of these compounds, pimobendan has currently progressed the furthest in clinical trials. Because it has both calcium sensitising and phosphodiesterase inhibiting activities it increases systolic calcium concentrations, though the increase is much less than occurs, for example, with $\bar{6}$ stimulation.⁴² Comparisons of energy requirements with pimobendan and with $\bar{6}$ stimulation indicate that pimobendan performs more favourably, causing little or no increase in energy utilisation.¹⁷ It also has beneficial inotropic effects in patients with cardiac failure, including an improvement in New York Heart Association functional class, increased cardiac and stroke volume indices, increased exercise capacity, and decreased right atrial pressure and catecholamine concentrations, and all this with few side effects.⁴³ Furthermore, these results were maintained after one month of treatment.⁴³ Left ventricular relaxation rate was actually improved by the drug.⁴⁴

In conclusion, then, calcium sensitivity is now known to be an important factor in modulating the contractile force of the heart. Sensitivity might be expected to be depressed in the diseased myocardium, and this makes it likely that calcium sensitisers will prove to be beneficial inotropic agents. Compared with conventional agents calcium sensitisers would be expected to cause fewer arrhythmias and increase energy requirements less for a given inotropic effect. In the next few years such drugs seem likely to become of clinical importance as a new class of positive inotropic agents.

JOHN A LEE

Department of Pathology,
Royal Victoria Infirmary,
Newcastle upon Tyne NE1 4LP

DAVID G ALLEN

Professor of Physiology,
University of Sydney,
New South Wales 2006,
Australia

- 1 Allen DG, Kurihara S. Calcium transients in mammalian ventricular muscle. *Eur Heart J* 1980;1(suppl A):5-15.
- 2 Morgan JP, Blinks JR. Intracellular Ca transients in the cat papillary muscle. *Can J Physiol Pharmacol* 1982;60:520-8.
- 3 Wier WG, Hess P. Excitation-contraction coupling in cardiac Purkinje's fibres. Effects of cardiotonic steroids on the intracellular Ca²⁺ transient, membrane potential and contraction. *J Gen Physiol* 1984;83:395-415.
- 4 Kurihara S, Konishi M. Effects of β -adrenoceptor stimulation on intracellular Ca transients and tension in rat ventricular muscle. *Pflugers Arch* 1987;409:427-37.
- 5 Katz AM. Cyclic adenosine monophosphate effects on the myocardium. *J Am Coll Cardiol* 1983;2:143-9.
- 6 Ebashi S, Endo M. Calcium ion and muscle contraction. *Prog Biophys Mol Biol* 1968;18:123-83.
- 7 Allen DG, Blinks JR. Calcium transients in aequorin-injected frog cardiac muscle. *Nature* 1978;273:509-13.
- 8 Smith TW, Antman EM, Friedmann PL, Blatt CM, Marsh JD. Digitalis glycosides: mechanisms and manifestations of toxicity. Part I. *Prog Cardiovasc Dis* 1984;26:413-58.
- 9 Smith TW, Antman EM, Friedmann PL, Blatt CM, Marsh JD. Digitalis glycosides: mechanisms and manifestations of toxicity. Part II. *Prog Cardiovasc Dis* 1984;26:495-540.
- 10 Smith TW, Antman EM, Friedmann PL, Blatt CM, Marsh JD. Digitalis glycosides: mechanisms and manifestations of toxicity. Part III. *Prog Cardiovasc Dis* 1985;27:21-56.
- 11 Farah AE, Alousi AA, Schwartz RP. Positive inotropic agents. *Annu Rev Pharmacol Toxicol* 1984;24:275-328.
- 12 Orchard CH, Elsner DA, Allen DG. Oscillations of intracellular Ca²⁺ in mammalian cardiac muscle. *Nature* 1983;304:735-8.
- 13 Wier WG, Kort AA, Stern MD, Lakatta EG, Marban E. Cellular calcium fluctuations in mammalian heart: direct evidence from noise analysis of aequorin signals in Purkinje fibers. *Proc Natl Acad Sci USA* 1983;80:7367-71.
- 14 Lederer WJ, Tsien RW. Transient inward current underlying arrhythmogenic effects of cardiotonic steroids in Purkinje fibres. *J Physiol (Lond)* 1976;263:73-100.
- 15 Kass RS, Lederer WJ, Tsien RW, Weingart E. Role of calcium ions in transient inward currents and aftercontractions induced by strophanthidin in cardiac Purkinje fibres. *J Physiol (Lond)* 1978;281:187-208.
- 16 Katz AM. Potential deleterious effects of inotropic agents in the therapy of chronic heart failure. *Circulation* 1986;73(suppl 3):184-90.
- 17 Hasenfuss G, Holubarsch C, Just H, Blanchard E, Mulieri LA, Alpert NR. Energetic aspects of inotropic interventions in rat myocardium. *Basic Res Cardiol* 1987;82(suppl 2):251-9.
- 18 Blinks JR, Endoh M. Modification of myofibrillar responsiveness to Ca²⁺ as an inotropic mechanism. *Circulation* 1986;73(suppl 3):85-98.
- 19 Ruegg JC. Effects of new inotropic agents on Ca²⁺ sensitivity of contractile proteins. *Circulation* 1986;73(suppl 3):78-84.
- 20 Fabiato A, Fabiato F. Effects of pH on the myofilaments and the sarcoplasmic reticulum of skinned cells from cardiac and skeletal muscles. *J Physiol (Lond)* 1978;276:233-55.
- 21 Kentish JC. The effects of inorganic phosphate and creatine phosphate on force production in skinned muscles from rat ventricle. *J Physiol (Lond)* 1986;370:585-604.
- 22 Solaro RJ, Moir AGJ, Perry SV. Phosphorylation of troponin I and the inotropic effect of adrenalin in the perfused rabbit heart. *Nature* 1976;262:615-6.
- 23 Bailey IA, Williams SR, Radda GK, Gadian DG. Activity of phosphorylase in total global ischaemia in the rat heart. *Biochem J* 1981;196:171-8.

- 24 Rovetto MJ, Whitmer JT, Neely JR. Comparison of the effects of anoxia and whole heart ischemia on carbohydrate utilisation in isolated working rat hearts. *Circ Res* 1973;32:681-99.
- 25 Matthews PM, Bland JL, Gadian DG, Radda GK. A ³¹P-NMR study of metabolism in the hypoxic perfused rat heart. *Biochem Soc Trans* 1981;9:236-7.
- 26 Allen DG, Orchard CH. Intracellular calcium concentration during hypoxia and metabolic inhibition in mammalian ventricular muscle. *J Physiol* 1983;339:107-22.
- 27 Lee JA, Allen DG. The effects of repeated exposures to anoxia on intracellular calcium, glycogen and lactate in isolated ferret heart muscle. *Pflugers Arch* 1988;413:83-9.
- 28 Allen DG, Lee JA, Smith GL. The consequences of simulated ischaemia on intracellular Ca²⁺ and tension in isolated ferret ventricular muscle. *J Physiol* 1989;410:297-323.
- 29 Jewell BR. A reexamination of the influence of muscle length on myocardial performance. *Circ Res* 1977;40:221-30.
- 30 Allen DG, Kentish JC. The cellular basis of the length-tension relation in cardiac muscle. *J Mol Cell Cardiol* 1985;17:821-40.
- 31 Endoh M, Blinks JR. Actions of sympathomimetic amines on the Ca²⁺ transients and contractions of rabbit myocardium: reciprocal changes in myofibrillar responsiveness to Ca²⁺ mediated through α and β adrenoceptors. *Circ Res* 1988;62:247-65.
- 32 Morano I, Hofmann F, Zimmer M, Ruegg JC. The influence of P-light chain phosphorylation by myosin light chain kinase on the calcium sensitivity of chemically skinned heart fibres. *FEBS Lett* 1985;189:221-4.
- 33 O'Dowd JJ, Robins DJ, Miller DJ. Detection, characterisation and quantification of carnosine and other histidyl derivatives in cardiac and skeletal muscle. *Biochim Biophys Acta* 1988;967:241-9.
- 34 Wendt JR, Stephenson DG. Effect of caffeine on Ca-activated force production in skinned cardiac and skeletal muscle fibres of the rat. *Pflugers Arch* 1983;398:210-6.
- 35 Solaro RJ, Ruegg JC. Stimulation of Ca²⁺ binding and ATPase activity of dog cardiac myofibrils by AR-L 115BS, a novel cardiotonic agent. *Circ Res* 1982;51:290-4.
- 36 Van Meel JCA. Effects of some cardiotonic agents on contractility of skinned fibres from mammalian heart. *Drug Res* 1987;37:679-82.
- 37 Herzig JW, Quast U. Increase in calcium sensitivity of myocardial contractile structures by DPI-106. *J Mol Cell Cardiol* 1984;16(suppl 3):6P.
- 38 Lues I, Siegel R, Harting J. Effect of isomazole on the responsiveness to calcium of the contractile elements in skinned cardiac muscle fibres of various species. *Eur J Pharmacol* 1988;146:145-53.
- 39 Müller-Beckmann B, Fruend P, Honerjager P, Kling L, Ruegg JC. In vitro investigations on a new positive inotropic and vasodilating agent (BM 14.478) that increases myocardial cyclic AMP content and myofibrillar calcium sensitivity. *J Cardiovasc Pharmacol* 1988;11:8-16.
- 40 Allen DG, Lee JA. EMD 53998 increases tension with little effect on the amplitude of calcium transients in isolated ferret ventricular muscle. *J Physiol (Lond)* 1989;416:43P.
- 41 Ferroni C, Spurgeon HA, Klockow M, Lakatta EG, Capogrossi MC. Contractile potentiation without increasing cytosolic calcium in single rat ventricular myocytes. *FASEB J* 1989;3:A1039.
- 42 Lee JA, Ruegg JC, Allen DG. Effects of pimobendan, a novel inotropic agent, on intracellular calcium and tension in isolated ferret ventricular muscle. *Clin Sci* 1989;76:609-18.
- 43 Renard M, Walter M, Liebens I, Dresse A, Bernard R. Pimobendan in chronic congestive heart failure. *Chest* 1988;93:1159-64.
- 44 Rousseau SS, Hanet C, Etienne J, Leonard J, Menegeot P, Pouleur H. Effects of UDCG 115 on inotropic state and left ventricular diastolic function in patients with heart failure. *Acta Pharmacol Toxicol* 1986;59:182.

Surgery and salmonella

Complications require prompt diagnosis and treatment

Salmonella infections have become all too familiar to the general public in recent years, but they are not perceived as illnesses that may require surgery. Nevertheless, surgical complications do occur—though fortunately they are rare. Human infection from different species of salmonella is usually seen in one of three forms: typhoid fever; enteritis—the most common form of salmonella disease; and extra-intestinal focal infections such as cholecystitis.

Typically, typhoid fever presents with a stepladder rise of fever; bacteraemia is followed finally by localisation of *Salmonella typhi* in Peyer's patches, which become ulcerated in the long axis of the bowel. Two main complications—haemorrhage and perforation—may be encountered in the third week of the illness. Frank haemorrhage calls for blood transfusion and heralds the possibility of perforation. Tiny perforations¹ may occur in the antimesenteric border of the terminal ileum,^{2,3} though in a 10th of cases the perforations occur in the caecum.

Typhoid perforation is a grave complication requiring prompt diagnosis, early resuscitation, and operative management—but the choice of surgical management is controversial.⁴ The bowel close to the perforation will hold sutures,⁵ and simple closure^{6,7} or excision of the Peyer's patch and closures have both been advocated.¹ The defect should be closed in a transverse direction to avoid narrowing of the bowel. Nevertheless, the rate of reoperation for leaks has been reported as a worrying 15%,¹ and resection of the bowel may be preferable, especially in the presence of multiple perforations.⁴ Right hemicolectomy is justified in cases of caecal

perforation. Any ulcers visible from the serosal surface but unperforated should be oversewn.

Gastroenteritis resulting from the various serotypes of *S enteritidis* is becoming increasingly common and may be extremely severe. The presentation varies but usually includes abdominal cramps, pain, and diarrhoea that may be bloody. This picture may be confused with that of ulcerative colitis.⁸ Diagnosis may sometimes be difficult: salmonella enteritis and ulcerative colitis are both common and their sigmoidoscopic appearances are similar. If the colitis is treated with corticosteroids then simultaneous systemic antibiotic cover is essential.⁸

The micro-organisms invade focal areas of the small and large bowel,⁹ and occasionally the colitis progresses to perforation. This complication has been reported after infection with many different species, including *S hadar*,¹⁰ *S typhimurium*,¹¹ and *S newport*.¹⁷ We recently encountered perforation and faecal peritonitis in a patient infected in the recent outbreak of *S kegougou*. A high index of suspicion is necessary during an outbreak of salmonella enteritis; if perforation of the colon does occur prompt colectomy is essential for a favourable outcome.¹²

Extraintestinal focal lesions with salmonella are often associated with other chronic disease. The focal lesions may manifest themselves long after an episode of enteritis or the original bowel infection may have been silent. *S enteritidis* has a predilection for blood vessels. Possibly, intravascular salmonellas localise at sites of atheromatous ulceration and may cause thrombosis of major vessels or may colonise vascular grafts.¹³ In addition they may damage the venous system, causing septic thrombophlebitis,¹⁴ iliac vein thrombosis, or pulmonary embolism. These focal lesions have a tendency to chronicity and may mimic tuberculosis, particularly in osteomyelitis of a vertebra or paravertebral abscess. Circulating *S typhi* is trapped by the liver and excreted in the bile. Acute cholecystitis as a complication of typhoid has become rare since the advent of antibiotics, but it may develop in a normal gall bladder and may proceed to perforation and biliary peritonitis.

Finally, the general surgeon may be asking for help in the management of patients who are chronic typhoid carriers. The typhoid carrier passes organisms in the faeces, presumably derived from a focus of infection in the gall bladder or biliary tract. Chronic typhoid cholecystitis is symptomless. The most notorious carrier was Mary Mallon, who was the source of an outbreak affecting 1300 people in New York in 1903. Although prolonged administration of ampicillin in high dosage has been successful in eliminating carriage of *S typhi* even in patients with gall stones and non-functioning gall bladders,¹⁵ cholecystectomy is regarded as the most effective way of permanently curing the carriers—provided there is no associated infection in the biliary or urinary tract.

J G MOSLEY
Consultant General and Vascular Surgeon

A K CHAUDHURI
Consultant Microbiologist

Leigh Infirmary,
Leigh,
Lancashire WN7 1HS

- 9 Mandel BK, Mani V. Colonic involvement in salmonellosis. *Lancet* 1976;i:887-8.
- 10 Pascall CR, Stearn EJ, Mosley JG. Salmonella hadar peritonitis. *Br Med J* 1980;280:26.
- 11 Tauner WA, Collins PG, Fahy AM. Spontaneous perforation of the colon in salmonella enterocolitis. *Br J Med Sci* 1980;149:37.
- 12 Gill KP, Feeley TM, Keane FBV. Toxic megacolon and perforation caused by salmonella. *Br J Surg* 1989;76:796.
- 13 Boyd JF. Colonic involvement in salmonellosis. *Lancet* 1976;i:1415.
- 14 Kurty JB. Leg abscesses caused by Salmonella heidelberg. *Lancet* 1976;i:200-1.
- 15 Christie AB. Treatment of typhoid carriers with ampicillin. *Br Med J* 1964;i:1609.

Raynaud's syndrome

Thymoxamine, iloprost, and ACE inhibitors are among the effective treatments now available

Raynaud's phenomenon was described over 100 years ago, but we still do not understand the pathogenesis of the abnormally prolonged (sometimes painful) episodic peripheral vasospasm in response to cold or emotional stimuli. The controversy has yet to be settled between the hyperactivity of the sympathetic nervous system described by Raynaud¹ and Lewis's local fault.²

The clinical classification into primary and secondary Raynaud's syndrome and phenomenon is also unsatisfactory. Patients who have Raynaud's phenomenon but are otherwise fit may commonly be shown to have one or more autoantibodies present in low titre. Some of these patients will later develop one of the connective tissue diseases.^{3,4}

So what do we know? Patients with Raynaud's syndrome may be shown to have increased sensitivity or density of the peripheral α adrenoceptors, or both, in both the vessel walls⁵ and the platelet membranes.⁶ Other factors implicated in the disease are abnormal platelet adhesiveness,⁷ increased whole blood and plasma viscosity,⁸⁻¹⁰ reduced red cell deformability,¹¹ reduced activity of the fibrinolytic system,^{12,13} an imbalance in the cyclo-oxygenase products of arachidonic acid—thromboxane (TXA₂) and prostacyclin (PGI₂)—and hypersensitivity of serotonin (5-HT₂) receptors.¹⁴ Whether any or all of these are crucial in the pathogenesis or are simply epiphenomena remains to be determined. Attempts to correct these abnormalities have been reflected in numerous therapeutic approaches to the disease. Despite the high proportion of women affected the contribution of oestrogen and other female sex hormones remains vague and poorly defined.¹⁵ A possible exception may be those patients whose symptoms begin at or near the menopause, in whom hormone replacement treatment may be beneficial.¹⁵

In the clinical setting primary Raynaud's syndrome is common.¹⁶ The average general practitioner may expect to see one or two new cases a year. Fortunately it is also usually benign and compatible with an entirely normal lifespan (unpublished data). The prognosis of secondary Raynaud's syndrome is that of the underlying disease and depends on the severity with which target organs are affected.^{3,17}

As the common stimulus is cold, either local or general, advice not to smoke and to keep warm remains the cornerstone of management. Smoking a cigarette may produce a fall in temperature of 2°C or 3°C in the fingertips of normal people in a comfortable ambient temperature. Patients may be helped to keep warm with appliances ranging from chemically activated handwarmers to electrically heated gloves, foot warmers, and hand and ear muffs. Further information concerning these may be obtained from the Raynaud's Association (112 Crewe Road, Alsagar, Cheshire ST7 2JA; tel 0270 872 776).

About one third of patients also have attacks in response to

- 1 Keenan JP, Hadley SP. The surgical management of thyroid perforations in children. *Br J Surg* 1984;71:928-9.
- 2 Dunkerley GE. Perforation of the ileum in enteric fever. *Br Med J* 1946;iii:454-7.
- 3 Dickson JAS, Cole GJ. Perforation of the terminal ileum. *Br J Surg* 1964;51:893-7.
- 4 Gibney EJ. Typhoid perforation. *Br J Surg* 1989;76:887-9.
- 5 Angorn IB, Pillary SP, Hegarty MM, Baker LW. Typhoid perforation of the ileum. *S Afr Med J* 1975;49:781-5.
- 6 Chouhan MK, Pandu SK. Typhoid enteric perforation. *Br J Surg* 1982;69:173-5.
- 7 Archampong EO. Operative treatment of typhoid perforation of the bowel. *Br Med J* 1969;iii:273-6.
- 8 Dronfield MW, Fletcher J, Langman MJS. Coincident salmonella infections and ulcerative colitis. *Br Med J* 1974;i:99-100.