abuse by women, and research must be undertaken to identify similarities and differences between men and women offenders—especially with regard to the motivation to abuse, the selection of victims, the efficacy of treatment, and the propensity to abuse again. Despite the feminist tenet “No penis, no harm,” clearly sexual abuse can no longer be considered the exclusive preserve of men.¹⁸

ROBERT WILKINS

Consultant Child and Adolescent Psychiatrist

Paxton House,
Reading RG3 2BA

- 1 Mrazek PB. Annotation: sexual abuse of children. *J Child Psychol Psychiatry* 1980;21:91-5.
- 2 American Humane Association. *National study on child neglect and abuse*. Denver: American Humane Association, 1981.
- 3 McCarty LM. Mother-child incest: characteristics of the offender. *Child Welfare* 1986;65:447-59.
- 4 Kendall-Tackett KA, Simon AF. Perpetrators and their acts: data from 365 adults molested as children. *Child Abuse Negl* 1987;11:237-45.

- 5 Mrazek P, Lynch M, Bentovim A. Recognition of sexual abuse in the United Kingdom. In: Mrazek P, Kempe CH, eds. *Sexually abused children and their families*. Oxford: Pergamon Press, 1987:35-50.
- 6 Mathis J. *Clear thinking about sexual deviations: a new look at old problems*. Chicago: Nelson-Holt, 1972.
- 7 Finkelhor D. Recognition of sexual abuse in children. *Pediatrics in Review* 1986;8:25-39.
- 8 Finkelhor D, Russell D. Women as perpetrators. In: Finkelhor D, ed. *Child sexual abuse: new theory and research*. New York: Free Press, 1984:171-87.
- 9 Margolis M. A case of mother-adolescent son incest: a follow-up study. *Psychoanal Q* 1984;53:355-85.
- 10 Faller KC. Women who sexually abuse children. *Violence and Victims* 1987;2:263-76.
- 11 Margolin L. The effects of mother-son incest. *Lifestyles: A Journal of Changing Patterns* 1986;5:104-14.
- 12 Krug RS. Adult male report of childhood sexual abuse by mothers: case descriptions, motivations and long-term consequences. *Child Abuse Negl* 1989;13:111-9.
- 13 Groth AN. The incest offender. In: Sgroi SM, ed. *Handbook of clinical intervention in sexual abuse*. Lexington: Heath, 1982:215-39.
- 14 Parsons T. The incest taboo in relation to social structure and the socialization of the child. *Br J Sociol* 1954;5:101-17.
- 15 Nasjleti M. Suffering in silence: the male incest victim. *Child Welfare* 1980;59:269-75.
- 16 Groth AN. Treatment of the sexual offender in a correctional institution. In: Greer J, Stuart I, eds. *The sexual aggressor: current perspectives on treatment*. New York: Van Nostrand Reinhold, 1983:160-76.
- 17 Groth AN. *Men who rape*. London: Plenum, 1979.
- 18 Banning A. Mother-son incest: confronting a prejudice. *Child Abuse Negl* 1989;13:563-71.

Unwanted pregnancies

High abortion rates indicate failure of family planning services

The vote last week in the House of Commons reducing the time limit for legal termination of pregnancy from 28 to 24 weeks' gestation (with certain exceptions) has been accepted by all concerned as marking the end of attempts by this parliament to amend the Abortion Act 1967. Public attention should now switch to the inconsistencies and defects in the way the act has been operating and in the provision of family planning services. Whenever parliament has debated the time limit for termination the medical profession has tried to explain that the way to reduce the numbers of late abortions is to make early abortion easier. One of the main reasons for late abortions remains the delay between a woman going to her doctor and the procedure being carried out: a survey by the Royal College of Obstetricians and Gynaecologists found that 45% of women operated on between 13 and 16 weeks' gestation and one fifth of those operated on after 20 weeks had seen their doctors before 13 weeks.¹ Much of this delay is linked with the variation throughout Britain in the way the act is interpreted. Half of the British women who have terminations pay for them—but the proportion varies from 82% in the west midlands to 15% in the Northern region.² Far too many women seeking termination still meet covert opposition and obstruction from doctors who dislike the concept of abortion on demand—yet in most of the country that is the way the act is interpreted.

Other countries—France, Italy, and The Netherlands—have recently given women a right to choose an abortion in the early months of pregnancy. Opinion polls in Britain show that around 80% of people think that women should have this right²—and if they did we believe that the numbers of late abortions would fall substantially. Meanwhile the Departments of Health have a duty to ensure that all women—wherever they live—have equal access to NHS clinics providing a fast, caring, and efficient service for early terminations.

Late or early, however, the total number of abortions is far too high. In Britain in the 1980s one in every five pregnancies was terminated; and almost all abortions are done essentially because the pregnancy was unwanted (less than 2% are for fetal abnormalities).³ Since the 1970s four out of every 10 conceptions outside marriage have ended in termination of the pregnancy. These figures horrify the campaigners against

the Abortion Act and they should horrify us all. Termination of a pregnancy must always be a miserable event for the woman, her partner, her family, and the doctors and nurses concerned.

The most effective way to reduce the numbers of abortions is not to make them illegal or more difficult but to reduce the numbers of unwanted pregnancies. Research studies have consistently shown that the way to achieve that goal is by making sex education and contraception readily and freely available to the whole population—but especially to the young and single. The rate of abortion per 1000 women aged 15-44 in England and Wales is more than twice that in The Netherlands—clear evidence that our services need improvement.²

Yet throughout Britain family planning clinics are being shut and services reduced as health authorities look for ways of making financial savings. Last week the Labour Party's shadow health minister, Harriet Harman, quoted cuts of 20-50% in every NHS region, affecting particularly rural and peripheral clinics and clinics for teenagers. Many general practitioners who offer contraceptive advice do not fit intra-uterine devices or caps, nor do general practitioners provide condoms. Alternatives to oral contraceptives are needed, however, at a time when many women have been influenced by reports linking the contraceptive pill with cancer and when the use of barrier contraceptives is being encouraged to reduce the risks of acquiring sexually transmitted diseases.

High priority should, we believe, be given to the maintenance and improvement of family planning services, and that will require specific government action. Most health authorities face such financial problems that they cannot take the long view. We are told that the NHS must become more efficient. Preventing unwanted pregnancy must be one of its most cost effective activities—and one that also helps reduce the total sum of human unhappiness.

TONY SMITH

Deputy editor, *BMJ*

- 1 Royal College of Obstetricians and Gynaecologists. *Report on fetal viability and clinical practice*. London: Royal College of Obstetricians and Gynaecologists, 1985.
- 2 Munday D, Francombe C, Savage W. Twenty one years of legal abortion. *Br Med J* 1989;298:1231-4.
- 3 Office of Population Censuses and Surveys. *1988 Abortion statistics. England and Wales*. London: HMSO, 1989. (Series AB, No 15.)