screening by assay of immunoreactive trypsin in dried blood spots,¹¹ but the general tendency seems to have been to await DNA based diagnostic methods, which should give more clear cut results. Now that the cystic fibrosis gene has been identified,⁹ the commonest mutation is easy to detect,^{12,13} and when the polymerase chain reaction is used dried Guthrie blood spots seem to be an adequate source of DNA for diagnosis.¹⁰ It would be unwise, however, to rush into large scale neonatal screening for cystic fibrosis. Precise genetic diagnosis of a recessively inherited condition raises many new and unsolved problems, as shown by experience with neonatal screening for sickle cell disease.

Abnormal haemoglobins can be detected cheaply and reliably by electrophoresis, and neonatal screening with follow up both reduces early mortality in sickle cell disease and also allows parents to be counselled about later pregnancies.¹⁴ Neonatal screening identifies about 40 heterozygotes for every homozygote detected (with cystic fibrosis the ratio would be closer to 100:1). What should be done for these babies? Clearly they should also be followed up because their parents will include numerous couples at risk who could be counselled before they have affected children. But there are no special counselling resources for families of carriers; the neonatal period is a sensitive time; most people are unfamiliar with recessive inheritance; there are pitfalls such as accidental revelation of non-paternity; and there is no mechanism to ensure that a genetic diagnosis remains attached to a person's medical records for life. Current thinking holds that it is better to leave families in ignorance than to inform them without counselling, so unless a trained haemoglobinopathy counsellor is available many families with asymptomatic (heterozygous) infants are not informed at all. Adequate resources for counselling for common genetic traits will be available only when it is perceived as a task for primary care workers and they have been given training, and that will take time.

Furthermore, neonatal screening is an inefficient method for detecting couples at risk of having children with recessively inherited disorders. Only half of the couples at risk can be found by following up the parents of heterozygous infants

as one quarter will already have had an affected child and the quarter who have had a normal child will escape detection. This leads to the consideration that most really precise genetic methods could be used just as well antenatally—to detect adult carriers of a disorder—as postnatally, when the result tells them that their child is already affected. Antenatal screening or screening before pregnancy allows the choice of prenatal diagnosis, and genetic counselling—however imperfect—is now becoming a familiar concept in antenatal clinics.¹⁵

As diagnostic methods become more precise the possibilities for neonatal diagnosis will certainly increase; but when the prime objective is reproductive counselling screening is likely to be pushed back to the antenatal period and earlier. Ultimately, this could actually reduce some indications for neonatal screening.

BERNADETTE MODELL

Consultant in Perinatal Medicine,
University College and Middlesex School of Medicine,
University College London, London WC1E 6HX

- 1 Cuckle HS, Wald NJ. Principles of screening. In: Wald NJ, ed. *Antenatal and neonatal screening*. Oxford: Oxford University Press. 1984:1-24.
- 2 World Health Organisation Working Group. Glucose-6-phosphate dehydrogenase deficiency. *Bull WHO* 1989;67:601-11.
- 3 McNeil TF, Sveger T, Thelin T. Psychological aspects of screening for somatic risk: the Swedish α_1 -antitrypsin experience. *Thorax* 1988;43:505-7.
- 4 Udall JN, Dixon M, Newman AP, Wright JA, James B, Bloch KJ. Liver disease in alpha-1-antitrypsin deficiency. A retrospective analysis of the influence of early breast- vs bottle-feeding. *JAMA* 1985;253:2679-82.
- 5 Dellamonica Ch, Robert JM, Cotte J, Collombel C, Dorche C. Systematic neonatal screening for Duchenne muscular dystrophy. *Lancet* 1978;ii:1100.
- 6 Gardner-Medwin D. Recognising and preventing Duchenne muscular dystrophy. *Br Med J* 1983;287:1083-4.
- 7 Smith RA, Rogers M, Bradley DM, Sibert JR, Harper PS. Screening for Duchenne muscular dystrophy. *Arch Dis Child* 1989;64:1017-21.
- 8 Hoffman EP, Brown RH, Kunkel LM. Dystrophin: the protein product of the Duchenne muscular dystrophy locus. *Cell* 1987;51:919-28.
- 9 Riordan J, Rommens JM, Kerem B-S, et al. Identification of the cystic fibrosis gene: cloning and characterisation of complementary DNA. *Science* 1989;245:1066-71.
- 10 Williams C, Weber L, Williamson R, Hjelm M. Guthrie spots for DNA-based carrier testing in cystic fibrosis. *Lancet* 1988;ii:639.
- 11 Wilken B, Chalmers C. Reduced morbidity in patients with cystic fibrosis detected by neonatal screening. *Lancet* 1985;ii:1319-21.
- 12 Newton CR, Heppinstall LE, Summers C, et al. Amplification refractory mutation system for prenatal diagnosis and carrier assessment in cystic fibrosis. *Lancet* 1989;ii:1481-3.
- 13 Ballabio A, Gibbs RA, Caskey CT. PCR test for cystic fibrosis deletion. *Nature* 1990;343:220.
- 14 Grover R. Program effects on decreasing morbidity and mortality. Newborn screening in New York City. *Pediatrics* 1989;83(suppl):819-22.
- 15 Royal College of Physicians of London. *Prenatal diagnosis and genetic screening; community and service implications. Report*. London: Royal College of Physicians of London, 1989.

Thiazides in the 1990s

The risk:benefit ratio still favours the drug

Though thiazide diuretics have been in regular use for over 30 years, debate about their role in therapeutics seems to increase as the years go by. Thiazides have been a mainstay in treating hypertension, in which (together with β adrenoceptor antagonists) they have been the drugs of first choice. They also have useful diuretic actions, particularly in conditions such as congestive heart failure. Thiazides work in hypertension in the long term by lowering peripheral resistance rather than by their diuretic effect,¹ and they are usually regarded as superior to loop diuretics in this condition.² The dose commonly used in hypertension, however, has probably been unnecessarily high; studies with cyclopenthiiazide have shown that the fall in blood pressure is as good with 125 μ g (a dose lower than currently marketed) as it is with 500 μ g.^{3,4}

Thiazide diuretics have always been known to be able to produce adverse effects, though when set against their volume of use their safety record is good.^{5,6} Some controversy has arisen in the context of long term treatment of hypertension when thiazide diuretics have been alleged to have long term

harmful effects. In the multiple risk factor intervention trial thiazide diuretics were suggested as the cause of an increase in the number of cardiac deaths in a subgroup of patients with minor electrocardiographic abnormalities.⁷ This was, however, a retrospective conclusion, and data from similar studies such as the hypertension detection and follow up program and the Medical Research Council hypertension trial failed to confirm the observation.^{8,9} Thiazide diuretics are well known to cause hypokalaemia; in Britain some 76 000 patients having long term treatment with these drugs may have a serum potassium concentration below 3.0 mmol/l.¹⁰ Hypokalaemia (by causing cardiac arrhythmias) has been suggested as the cause of the postulated increase in the number of deaths. Nevertheless, the serum potassium concentrations do not correlate well with intracellular concentrations of potassium, and while the issue remains controversial,^{10,11} the evidence does not suggest that hypokalaemia induced by thiazides is an important cause of cardiac arrhythmias.¹²⁻¹⁴ Thiazides have also been alleged to be harmful to patients in the long term

because of an increase in the serum cholesterol concentration — particularly in that of the low density lipoprotein fraction.¹⁵ Most of the evidence suggests that this increase in low density lipoprotein cholesterol concentration is short term, and the cholesterol concentration falls to baseline concentrations or below in the long term.^{14 16 17}

Thiazides cause other adverse effects, and a recent observation was that of impotence in men taking thiazides in the Medical Research Council trial.⁹ Perhaps as a result the quality of life in patients taking thiazide diuretics may be inferior to that of patients taking propranolol.¹⁸ Sodium depletion is rarely a problem in patients taking thiazide diuretics alone but may be a problem in patients, particularly in elderly patients, if given together with amiloride — usually as Moduretic.^{5 6}

A diabetogenic effect of thiazides was suspected soon after their first clinical use.¹⁹ Worsening of glucose tolerance may occur with long term use of thiazides²⁰ and seems to be greater with the longer acting members of the class — for example, clopamide.²¹ Diabetes may also occur in patients taking loop diuretics, but clinical problems are unusual with both groups of diuretics. Gout is, however, a common adverse effect of thiazides and other diuretics; but though half of all patients taking thiazides have hyperuricaemia, less than 2% of patients suffer from clinical gout.²² Rashes are occasionally a problem with thiazide diuretics, but only loop diuretics seem to cause ototoxicity.^{5 6} Clinical problems may arise when diuretics are given with non-steroidal anti-inflammatory drugs as these drugs blunt the hypotensive effects of thiazides.²³

This liturgy of side effects may persuade some clinicians to stop prescribing thiazides for patients with hypertension, but there are benefits besides the obvious diuretic or hypotensive effect. Thiazides reduce the urinary excretion of calcium (unlike loop diuretics, which increase it),^{5 6} and there are suggestions that thiazides reduce the rate of recurrence of urinary calculi.²⁴ Nevertheless, further trials are needed. Two recent studies, however, have shown a reduced incidence of hip fracture in elderly patients receiving long term treatment with thiazides.^{25 26}

Overall, the risk:benefit ratio for thiazides still favours the drugs. We should aim at using the lowest possible dose, as there is a flat dose response curve.²⁷ It should rarely be necessary to use more than 2.5 mg of bendrofluzide a day. We should be alert to the possibility of hypokalaemia and take

avoiding action either through the diet or the use of potassium sparing diuretics. Combined preparations of thiazide with potassium rarely help.^{5 6} Finally, we should remember that in large parts of the developing world thiazide diuretics offer the only cost effective way of treating hypertension.

M ORME

Professor of Clinical Pharmacology,
University of Liverpool,
Liverpool L69 3BX

- 1 Van Brummelen P, Man In't Veld AJ, Schalekamp MADH. Hemodynamic changes during long term thiazide treatment of essential hypertension in responders and non-responders. *Clin Pharmacol Ther* 1980;27:328-36.
- 2 Anderson J, Godfrey BE, Hill DM, Munro-Faure AD, Sheldon J. A comparison of the effects of hydrochlorothiazide and of frusemide in the treatment of hypertensive patients. *Q J Med* 1971;40:541-60.
- 3 McVeigh G, Galloway D, Johnston D. The case for low dose diuretics in hypertension: comparison of low and conventional doses of cyclopentiazide. *Br Med J* 1988;297:95-8.
- 4 Carlsen JE, Køber L, Torp-Pedersen C, Johansen P. Relation between dose of bendrofluzide, antihypertensive effect, and adverse biochemical effects. *Br Med J* 1990;300:975-8.
- 5 Lant AF. Diuretics. A review. *Drugs* 1985;29:57-87.
- 6 Lant AF. Diuretics. A review. *Drugs* 1985;29:162-88.
- 7 Multiple Risk Factor Intervention Trial Research Group. Baseline rest electrocardiographic abnormalities, antihypertensive treatment and mortality in the multiple risk factor intervention trial. *Am J Cardiol* 1985;55:1-5.
- 8 Hypertension Detection and Follow-up Program Cooperative Research Group. The effect of antihypertensive drug treatment on mortality in the presence of resting electrocardiographic abnormalities at baseline: the HDFP experience. *Circulation* 1984;70:996-1003.
- 9 Medical Research Council Working Party. MRC trial of treatment of mild hypertension: principal results. *Br Med J* 1985;291:97-105.
- 10 Poole-Wilson PA. Diuretics, hypokalaemia and arrhythmias in hypertensive patients: still an unresolved problem. *J Hypertens [Suppl]* 1987;5:51-5.
- 11 Hollifield JW. Electrolyte disarray and cardiovascular disease. *Am J Cardiol* 1989;63:21B-6B.
- 12 Khair GZ, Kochar MS. Mild hypertension, diuretics and cardiac arrhythmias: consensus amid controversy. *Am Heart J* 1988;116:216-21.
- 13 Langford HG. The effect of potassium depletion by thiazides. *Arch Intern Med* 1988;148:1265-6.
- 14 Freis ED, Papademetriou V. How dangerous are diuretics? *Drugs* 1985;30:469-74.
- 15 Grimm RH Jr, Leon AS, Hunninghake DB, Lenz K, Hannan P, Blackburn H. Effect of thiazide diuretics on plasma lipids and lipoprotein in mildly hypertensive patients. A double blind controlled trial. *Ann Intern Med* 1981;94:7-11.
- 16 Ames RP. The effects of antihypertensive drugs on serum lipids and lipoproteins. 1. Diuretics. *Drugs* 1986;32:260-78.
- 17 Burriss JF, Freis ED. Thiazides do not cause long term increases in serum lipid concentrations. *Arch Intern Med* 1985;145:2264-5.
- 18 Williams GH, Croog SH, Levine S, Testa MA, Sudilovsky A. Impact of antihypertensive therapy on quality of life: effect of hydrochlorothiazide. *J Hypertens [Suppl]* 1987;5:29-35.
- 19 Goldner MG, Zarowitz H, Akgun S. Hyperglycaemia and glycosuria due to thiazide derivatives administered in diabetes mellitus. *N Engl J Med* 1960;262:403-5.
- 20 Murphy MB, Lewis PJ, Kohner E, Schumer B, Dollery CT. Glucose intolerance in hypertensive patients treated with diuretics: a fourteen year follow up. *Lancet* 1982;ii:1293-5.
- 21 Hicks BH, Ward JD, Jarrett RJ, Keen H, Wise P. A controlled study of clopamide and clorexolone and hydrochlorothiazide in diabetes. *Metabolism* 1973;22:101-9.
- 22 Beevers DG, Hamilton M, Harpur JE. The long term treatment of hypertension with thiazide diuretics. *Postgrad Med J* 1971;47:639-43.
- 23 Favre L, Glasson PH, Riondel A, Vallotton MB. Interaction of diuretics and non-steroidal anti-inflammatory drugs in man. *Clin Sci* 1983;64:407-15.
- 24 Churchill DN, Taylor DW. Thiazides for patients with recurrent calcium stones: still an open question. *J Urol* 1985;133:749-51.
- 25 Roy WA, Giffin MR, Downey W, Melton WJ III. Long term use of thiazide diuretics and risk of hip fracture. *Lancet* 1989;i:687-90.
- 26 La Croix AZ, Wienpahl J, White LR, et al. Thiazide diuretic agents and the incidence of hip fracture. *N Engl J Med* 1990;322:286-90.
- 27 Cranston WI, Juel-Jensen BE, Semmence AM, Handfield-Jones RPL, Forbes JA, Mutch LMM. Effects of oral diuretics on raised arterial pressure. *Lancet* 1963;ii:966-70.

The mental health of Asians in Britain

Little disease or underreporting?

Any general statements on the mental health of British Asians are hard to sustain, given the cultural heterogeneity, range of length of settlement in Britain, and variation in eagerness towards (and success in) assimilating into Western culture. Nevertheless, some 50 papers and several reviews have discussed the mental health of people in Britain whose origins lie in the Indian subcontinent, and these suggest that less mental illness is found among most groups of British Asians than might be expected.^{1 3} The competing explanations are that British Asians genuinely experience less mental illness, that general practitioners are less good at identifying or treating mental illness in their Asian patients, and that Asians are reluctant to present with such illnesses.

There is evidence to support the suggestion that British

Asians have less mental illness than the indigenous population. Despite evidence from small studies to the contrary^{4 5} Asians seem to consult their general practitioners more frequently than do native Britons,^{6 9} but these high consultation rates do not apply when mental illness alone is considered.⁹ Referrals to one east London child psychiatry service, for example, included remarkably few Bengalis.¹⁰ Moreover, a considerable body of evidence shows that British Asians have low levels of neurotic traits or psychiatric symptoms.^{5 11-15}

Possibly, however, general practitioners are simply not recognising mental illness in their Asian patients. Such a failure may be part of a wider failure to communicate between British doctors and Asian patients.^{16 17} One study discovered that general practitioners find it harder to recognise psychi-