central laboratories, because of its reliability, safety, and economy.⁴¹ Although it is only one factor in the trend towards more testing nearer the patient and population screening, dry chemistry is undoubtedly the leader and the one with the greatest potential.

VINCENT MARKS

Professor of Clinical Biochemistry,
University of Surrey,
Guildford GU2 5XH

- 1 Marks V. The clinical laboratory: nearer the patient. *Advances in Pathology* 1988;1:1-30.
- 2 Sandler V. Immediacy: speed of response from the point of view of the physician. In: Marks V, Alberti KGMM, eds. *Clinical biochemistry nearer the patient*. Edinburgh: Churchill Livingstone, 1985:38-51.
- 3 Hilton S. Near patient testing in general practice, a review. *British Journal of General Practice* 1990;40:31-6.
- 4 Belsey R, Baer D, Sewell D. Laboratory test analysis near the patient. *JAMA* 1986;255:775-86.
- 5 Iwamoto I, Yamazaki H, Kimura A, Ochiai K, Tomioka H, Yoshida S. Comparison of a multi-allergen dipstick IgE assay to skin-prick test and RAST. *Clin Exp Allergy* 1990;20:175-9.
- 6 Imam AA, Novey HS, Orgel HA, Klein GL, Meltzer EO, Ziering RW. A simplified screening test for the diagnosis of allergy. *West J Med* 1990;162:313-6.
- 7 Twigg JT, Gray RL, Pichler K, Weiner R. Evaluation of multi-allergen dipstick screening test. *Ann Allergy* 1989;63:225-8.
- 8 Newall RG, Howell R, eds. *Clinical urinalysis: the principles and practice of urine testing in the hospital and community*. Stoke Poges: Ames Division, Miles Limited, 1990.
- 9 Anonymous. Is routine urinalysis worthwhile? [Editorial]. *Lancet* 1988;i:747.
- 10 Tuel SM, Meythaler JM, Cross LL, McLaughlin S. Cost effective screening by nursing staff for urinary tract infection in the spinal cord injured patient. *Am J Phys Med Rehabil* 1990;69:128-31.
- 11 Flanagan PG, Rooney PG, Davies EA, Stout RW. Evaluation of four screening tests for bacteriuria in elderly people. *Lancet* 1989;i:117-9.
- 12 Pels RJ, Bor DH, Woolhandler S, Himmelstein DU, Lawrence RS. Dipstick urinalysis screening of asymptomatic adults for urinary tract disorders. 2. Bacteriuria. *JAMA* 1989;262:1221-4.
- 13 Doran HM, Kensit JG. Screening for bacteriuria with Clinitec-200. *J Clin Pathol* 1988;41:1127-9.
- 14 Richardson H. Urine testing in schools: clinical considerations. In: Newall RG, Howell R, eds. *Clinical urinalysis: the principles and practice of urine testing in the hospital and community*. Stoke Poges: Ames Division, Miles Limited, 1990:86-92.
- 15 Propp DA, Weber D, Ciesla ML. Reliability of a urine dipstick in emergency department patients. *Ann Emerg Med* 1989;18:560-3.
- 16 Britton JP, Dowell AC, Whelan P. Dipstick haematuria and bladder cancer in men over 60: results of a community study. *BMJ* 1989;299:1010-2.
- 17 Marks V, Dawson A. Rapid stick method for determining blood glucose concentration. *BMJ* 1985;290:293-4.
- 18 Stahler F. The technology of dry versus wet chemistry. In: Marks V, Alberti KGMM, eds. *Clinical biochemistry nearer the patient* II. Eastbourne: Baillière Tindall, 1986:10-20.
- 19 Price CP, Koller PU. A multicentre study of the new Reflotron system for the measurement of urea, glucose, triacylglycerols, cholesterol, γ -glutamyltransferase and haemoglobin. *J Clin Chem Clin Biochem* 1988;26:233-50.
- 20 Maguire TA, McElroy JC. Performance of "bench top" cholesterol analysers. *Pharmaceutical Journal* 1990 May 12:566-8.
- 21 Okuda K. A compact, automated, dry-reagent chemistry analyzer. *Journal of the International Federation of Clinical Chemistry* 1990;2:39-44.
- 22 Zuk RF, Ginsberg VK, Houts T, et al. Enzyme immunochemistry—a quantitative immunoassay requiring no instrumentation. *Clin Chem* 1985;31:1144-50.
- 23 Cuss FM, Palmer JB, Barnes PJ. Rapid estimation of plasma theophylline concentrations. *BMJ* 1985;291:384.
- 24 Mould G, Marks V. The use of solid-phase chemistry in therapeutic drug monitoring. *Clin Pharmacokinet* 1988;14:65-70.
- 25 Chandler MHH, Clifton GD, Louis BA, et al. Home monitoring of theophylline levels: a novel therapeutic approach. *Pharmacotherapy* 1990;10:294-300.
- 26 Anonymous. TDM for community pharmacists? [Editorial]. *Pharmaceutical Journal* 1990 Sept 15:357.
- 27 Paz G, Yogev L, Gottreich A, et al. Determination of urinary luteinizing hormone for prediction of ovulation. *Gynecol Obstet Invest* 1990;29:207-10.
- 28 Marks V. The economics of near-patient testing. In: Kerkhof PLM, van Dieijen-Visser MP, eds. *Laboratory data and patient care*. New York: Plenum, 1988:17-23.
- 29 Harter G, Klimm HD, König B, Koberstein R. Evaluation of dry chemistry laboratory methods for step diagnosis in general medicine. *Fortschr Med* 1990;108:61-6.
- 30 Shephard MDS, Penberthy OA, Fraser CG. Urinalysis in an Australian teaching hospital. *Med J Aust* 1982;i:300-1.
- 31 Drucker RF, Williams DRR, Price CP. Quality assessment of blood glucose monitors in use outside the hospital laboratory. *J Clin Pathol* 1983;36:948-53.
- 32 Rady AA, Jones KW, Halloran SP, Marks V. Bedside blood glucose measurement: quality control and assessment. *Ann Clin Biochem* 1987;24(suppl 2):170-1.
- 33 Smith LA, McNaught J, Hutchinson AS. How well do nurses perform blood glucose analyses at the diabetic clinic? An assessment using the Reflotron analyser. *Ann Clin Biochem* 1990;27:156-7.
- 34 Blackburn J. Curbing the cost of consumables. *Doctor* 1990 April:72.
- 35 Venables MGM. Cutting the cost of urine tests. *Practice Update* 1990 June:1206.
- 36 Gallagher EJ, Schwartz E, Weinstein RS. Performance characteristics of urine dipsticks stored in open containers. *Am J Emerg Med* 1990;8:121-3.
- 37 Fraser CG, Geary TD, McLellan GH. Guidelines for the performance of clinical biochemistry tests outside laboratories. *Clinical Biochemistry Reviews* 1981;2:24-5.
- 38 Anonymous. Cholesterol testing: council guidelines [Editorial]. *Pharmaceutical Journal* 1988 Dec 17:786.
- 39 Marks V. Essential considerations in the provision of near-patient testing facilities. *Ann Clin Biochem* 1988;25:220-5.
- 40 Department of Health and Social Security. *Blood glucose meters*. London: DHSS, 1987. (Hazard notice HN (87)13.)
- 41 Ashby JP, Blundell G. Dry slide clinical chemistry. *Lancet* 1990;336:886.

Bovine somatotrophin and public health

Need for further independent and more extensive investigation

Last December the European Council of Agriculture Ministers agreed to extend for a year the ban on the commercial use of recombinant bovine somatotrophin to boost the milk yield of dairy cows, "pending the results of relevant scientific experiments."¹ For the past four years it has been used only on a few commercial farms (of undisclosed identity) for purposes of evaluation, although milk from treated cows has been mixed with the normal supply to consumers. Concerns have been expressed that treatment may jeopardise animal health, but official pronouncements have so far denied that milk from treated animals poses any threat to human health.² Yet preserving consumer confidence, strained by recent food scares, may demand more than ex cathedra statements.

Potential risks to human health of bovine somatotrophin excreted in milk have been much debated. Several reasons have been advanced for discounting them. Only a very small amount of bovine somatotrophin finds its way into milk and what does is biologically inactive in humans, is destroyed by pasteurisation, and is likely to suffer extensive proteolysis on ingestion.² Critics claim that as most recombinant somatotrophins are xenobiotics (differing from pituitary somatotrophin by from one to nine amino acid residues) they might induce immunogenic or allergenic responses at the gut wall if not systemically.³ Insulin-like growth factor 1 is potentially a more serious risk as its concentration in milk increases in parallel with bovine somatotrophin's galactopoietic effect. Bovine insulin-like growth factor 1 is identical to human insulin-like growth factor 1: it is a potent mitogen and is not destroyed by pasteurisation.² Moreover, in colostrum insulin-

like growth factor 1 exists in a truncated form, up to 10 times more potent than insulin-like growth factor 1 itself.⁴

Assurances of the safety of milk from cows treated with bovine somatotrophin are therefore based on the presumption that increased concentrations of insulin-like growth factor 1 in milk, even if statistically significant, are not biologically important. Evidence used to support this view includes the observations that concentrations of insulin-like growth factor 1 remain in the "physiological range" and that when given orally to rats no systemic effects occurred.² Such assertions, however, need qualifying. Firstly, including the concentration of insulin-like growth factor 1 in colostrum in the "physiological range," though scientifically valid, is misleading. Colostrum, whose concentration of insulin-like growth factor 1 falls from greater than 150 μ g/l at calving to about 25 μ g/l within four days,⁵ is not marketed.

Moreover, reports of the increase in insulin-like growth factor 1 concentration vary greatly:² one described a 360% increase after just seven daily injections,⁶ whereas it is proposed that in commercial practice "14 days' worth" would be injected once every fortnight. Even if all concentrations remained in the physiological range the mean values would increase. Secondly, some significant responses to oral insulin-like growth factor 1 in rats were reported (for example, increased tibia length) but, questionably, were discounted as unrelated to treatment.² Perhaps, more importantly, there are virtually no published reports of the presence or potency of truncated insulin-like growth factor 1 in milk of cows treated with bovine somatotrophin or of the effects

of insulin-like growth factor 1 on tissues of the digestive tract.

In addition to insulin-like growth factor 1, milk contains several other growth factors, which may have a role in neonatal gut development.⁷ Their importance for other consumers is problematical. As a recent National Institutes of Health panel report concluded: "Whether the additional amount of insulin-like growth factor 1 in milk from [recombinant bovine somatotrophin] treated cows has a local effect on the esophagus, stomach or intestines is unknown."⁸ Treatment with bovine somatotrophin also changes the nutrient content of milk—for example, the concentration of casein decreases and the fatty acid content is altered.⁹ Whether bovine somatotrophin would cause changes of biological importance to consumers would depend on the relative contribution of treated cows' milk to total milk and dairy products consumed and on the level of consumption. Milk is seldom free from contaminants, and mastitis in dairy herds results in low (and monitored) concentrations of certain antibiotics in milk. But if treatment with bovine somatotrophin increases the incidence of mastitis then consumers would be increasingly exposed to antibiotics in milk.³

These risks relate to consumption of milk and dairy products from cows treated with bovine somatotrophin. Paradoxically, greater risks might be associated with their non-consumption, because milk is an important source of calcium, protein, and vitamins. Public attitudes to bovine somatotrophin are influenced by the widespread suspicion that "commercial confidentiality" is being used as a cloak to obscure product deficiencies, with its manufacturers accused of selective publication of data. The implications for public health are clear: whether or not the suspicions are justified, consumer rejection of milk could lead to its substitution by far less nutritious alternatives.

Concern for consumer confidence led to calls for extending the criteria for assessing veterinary medical products to include an evaluation of likely socioeconomic impact. This proposal was recently incorporated into a draft regulation issued by the European Commission,¹⁰ for which bovine somatotrophin may well prove to be a test case. If bovine somatotrophin is not to cause yet another damaging food scare there is urgent need to adopt procedures for its assessment that are both open to public scrutiny and suitably comprehensive. Such assessments should include rigorously monitored blind trials.^{11 12}

T B MEPHAM

Reader in Lactational Physiology,
University of Nottingham,
Loughborough LE12 5RD

- 1 Ministry of Agriculture, Fisheries and Food. *News Release 394/90*. London: Ministry of Agriculture, Fisheries and Food, 1990.
- 2 Juskevich JC, Guyer CG. Bovine growth hormone: human food safety evaluation. *Science* 1990;249:875-83.
- 3 Kronfeld DS. Safety of bovine growth hormone. *Science* 1991;251:256.
- 4 Francis GL, Upton FM, Ballard FJ, McKneil KA, Wallace JC. Insulin-like growth factors 1 and 2 in bovine colostrum. *Biochem J* 1988;251:95-103.
- 5 Malven PV, Head HH, Collier RJ, Bounomo FC. Periparturient changes in secretion and mammary uptake of insulin and in the concentrations of insulin and insulin-like growth factors in milk of dairy cows. *J Dairy Sci* 1987;70:2254-65.
- 6 Prosser CG, Fleet IR, Corps AN. Increased secretion of insulin-like growth factor 1 in milk of cows treated with recombinantly-derived bovine growth hormone. *J Dairy Res* 1989;56:17-26.
- 7 Koldovsky O. Search for role of milk-borne biologically active peptides for the suckling. *J Nutr* 1989;119:1546-51.
- 8 National Institutes of Health. *Technology assessment conference statement: bovine somatotrophin*. Bethesda, Md: NIH, 1990.
- 9 Baer RJ, Tieszen KM, Schingoethe DJ, et al. Composition and flavor of milk produced by cows injected with recombinant bovine somatotrophin. *J Dairy Sci* 1989;72:1424-34.
- 10 Commission of the European Communities (Directorate General for Agriculture). Draft proposal for a council regulation concerning the use of certain substances and techniques intended for administration or application to animals to stimulate their productivity. Brussels: Commission of European Communities, 1991.
- 11 Mepham TB. Criteria for the public acceptability of biotechnological innovations in animal production. In: Heap RB, Prosser CG, Lamming GE, eds. *Biotechnology in growth regulation*. London: Butterworths, 1919:203-12.
- 12 Mepham TB. Questions of validity in mammary physiology: methodology and ethics. *Proc Nutr Soc* 1989;48:1-7.

Drug misuse and the Dutch

A matter of social wellbeing and not primarily a problem for the police and the courts

The Netherlands has a policy on drugs which is certainly individual and is sometimes described as soft because it condones their use. The Dutch believe, however, that a pragmatic approach aimed at seeking solutions is more effective than one that is emotional and dogmatic. They do not have a laissez faire attitude but they do not like a "war on drugs" either as that creates war language and a war atmosphere.

The Netherlands recognises that it is part of Europe, and it has made both possession of and trafficking in all drugs illegal. But the criminal law is enforced in a way that is intended primarily to decrease the drug supply and not to punish use. The government is aware that these two attitudes must conflict to some extent: legal restrictions are needed on the supply of drugs because of their harmful (primary) effects, but making drugs illegal creates additional (secondary) physical and social problems for both users and society. The effects of drug use are often confused with the effects of drug policy. The criminal nature of so many drug problems has blinded many people to the primary effects of drug use. This confusion has made the main focus of drug policy in most countries the fight against international drug trafficking.

Recognition of the side effects of drug use is the basis of the Dutch tolerance of users. This offers room for accepting pragmatic aims through which the side effects can be tackled.

Misuse of drugs is primarily seen, therefore, as a matter of social wellbeing and public health rather than as a problem for the police and the courts. The protection of health and social wellbeing is the primary aim, and the Ministry of Health has always played the leading part.

The criminal justice system maintains a clear cut distinction between drug users and traffickers. It avoids classifying the actions of users as offences since they would then no longer be accessible to any form of prevention or intervention. The use of discretion by the police and the prosecutions department allows tolerance of the possession of "hard drugs" (heroin, cocaine, amphetamines, and so on), which for cannabis extends not only to possession but also to the sale of small quantities. No special action is taken by the police to detect the possession of drugs for personal use or selling up to 30 g of cannabis products.

This policy has resulted in dealers selling small quantities of cannabis products in coffee shops—but it prevents users from going underground and sliding into the fringes of society where the risks may increase. If young people experimenting with drugs are obliged to buy the relatively less dangerous cannabis products on the illegal market where the harder drugs also circulate there is a great risk of their turning to the latter at some point.

Nevertheless, while drug users in possession of quantities