explain why nature requires the existence of two highly similar peptides. If chemical agonists or antagonists could be targeted—for example, by direct skin application—interesting therapeutic possibilities would be raised.

DOMHNALL J O'HALLORAN
Research Fellow
S R BLOOM
Professor of Endocrinology

Division of Endocrinology and Metabolism,
Department of Medicine,
Hammersmith Hospital,
London W12 0NN

- 1 Emeson RB, Hedjran F, Yeakley JM, Guise JW, Rosenfeld MG. Alternative production of calcitonin and CGRP mRNA is regulated at the calcitonin-specific splice acceptor. *Nature* 1989;**341**:76-80.
- 2 Kittur SD, Hoppener JWM, Antonarakis SE, et al. Linkage map of the short arm of human chromosome 11: location of genes for catalase, calcitonin and insulin-like growth factor II. *Proc Natl Acad Sci USA* 1985;**82**:5064-7.
- 3 Rosenfeld MG, Mermod JJ, Amara SG, et al. Production of a novel neuropeptide encoded by the calcitonin gene via tissue specific RNA processing. *Nature* 1983;**304**:129-35.
- 4 Morris HR, Panico M, Etienne T, Tippins J, Girgis SI, MacIntyre I. Isolation and characterisation of human calcitonin gene-related peptide. *Nature* 1984;**308**:746-8.
- 5 Alevizaki M, Shiraiishi A, Rassool FV, Ferrier CJM, MacIntyre I, Legon S. The calcitonin like sequence of the beta CGRP gene. *FEBS Lett* 1986;**206**:47-52.
- 6 Steenbergh PH, Hoppener JWM, Zandberg J, Lips CJM, Jansz HS. A second human calcitonin/ CGRP gene. *FEBS Lett* 1985;**183**:403-7.
- 7 Hoppener JWM, Steenbergh PH, Zandberg J, et al. The second human calcitonin/CGRP gene is located on chromosome 11. *Hum Genet* 1985;**70**:259-63.
- 8 Zaidi M, Breimer LH, MacIntyre I. Biological peptides from the calcitonin gene. *Q J Exp Physiol* 1987;**72**:371-408.
- 9 Goodman EC, Iverson LL. Calcitonin gene-related peptide: novel neuropeptide. *Life Sci* 1986;**38**:2169-78.
- 10 Amara SG, Arizaj JL, Leff SE, Swanson LW, Evans RM, Rosenfeld MG. Expression in brain of a messenger RNA encoding a novel neuropeptide homologous to calcitonin gene-related peptide. *Science* 1985;**229**:1094-7.
- 11 Tschopp FA, Henke H, Petermann JB, et al. Calcitonin gene-related peptide and its binding sites in the human central nervous system and pituitary. *Proc Natl Acad Sci USA* 1985;**82**:248-52.
- 12 Gibson SJ, Polak JM, Gaiad A, et al. Calcitonin gene-related peptide is expressed in sensory ganglia and also in spinal motoneurons in man and rat. *Neurosci Lett* 1988;**91**:283-8.
- 13 Wimalawansa SJ, Morris HR, Etienne A, Blench I, Pancio M, MacIntyre I. Isolation, purification and characterisation of β -h CGRP from human spinal cord. *Biochem Biophys Res Commun* 1990;**167**:993-1000.
- 14 Mulderry PK, Ghatei MA, Rodrigo K, et al. Calcitonin gene-related peptide in cardiovascular tissues of the rat. *Neuroscience* 1985;**14**:9474-54.
- 15 Wharton J, Gulbenkian S, Mulderry PK, et al. Capsaicin induces a depletion of calcitonin gene-related peptide immunoreactive nerves in the cardiovascular system of the guinea pig and rat. *J Auton Nerv Syst* 1986;**16**:298-309.
- 16 Wanaka A, Matsuyama T, Yoneda S, et al. Origins and distribution of calcitonin gene-related peptide-containing nerves in the wall of the cerebral arteries of the guinea pig with special reference to co-existence with substance P. *Cell Mol Biol* 1987;**33**:201-9.
- 17 Mulderry PK, Ghatei MA, Spokes RA, et al. Differential expression of alpha-CGRP and beta-CGRP by primary sensory neurons and enteric autonomic neurons of the rat. *Neuroscience* 1988;**25**:195-205.
- 18 Petermann JB, Born W, Chang J-Y, Fischer JA. Identification in the human central nervous system, pituitary, and thyroid of a novel calcitonin gene-related peptide, and partial amino acid sequence in the spinal cord. *J Biol Chem* 1987;**262**:542-5.
- 19 Tiller-Borich JK, Capili H, Gordan GS. Human brain calcitonin gene-related peptide (CGRP) is concentrated in the locus caeruleus. *Neuropeptides* 1988;**11**:55-61.
- 20 Henke H, Sigrist S, Lang W, Schneider J, Fisher JA. Comparison of binding sites for the calcitonin gene-related peptides I and II in man. *Brain Res* 1987;**410**:404-8.
- 21 Inagaki S, Kito S, Kubota Y, Girgis S, Hillyard CJ, MacIntyre I. Autoradiographic localisation of calcitonin gene-related peptide binding sites in human and rat brain. *Brain Res* 1986;**374**:287-98.
- 22 Jonas V, Lin CR, Kawashima E, et al. Alternative RNA processing events in human calcitonin/ calcitonin gene-related peptide gene expression. *Proc Natl Acad Sci USA* 1988;**82**:1994-8.
- 23 Hoppener JW, Steenberg PH, Slebos RJ, et al. Expression of the second calcitonin/calcitonin gene-related peptide gene in Ewing sarcoma cell lines. *J Clin Endocrinol Metab* 1987;**64**:809-17.
- 24 Edbrooke MR, Parker D, McVey JH, et al. Expression of the human calcitonin/CGRP gene in lung and thyroid carcinoma. *Embryo Journal* 1985;**4**:715-24.

Multiple births: too often a disaster

The buck stops at those treating infertility

Although much is known about twin pregnancy—its effects on maternal health and perinatal outcome and its costs to families and society—little is known about higher order births. To fill this gap the Office of Population Censuses and Surveys set up a national study of triplet and higher order births.¹ Its aim was to survey all such births in the United Kingdom over five years (1980 and 1982-5).

Interest in the topic was aroused by the increasing number of multiple deliveries after treatment of subfertility, and the report documents that 36% of mothers of triplets and 70% of

- 25 Kim S, Morimoto S, Kawai Y, Onishi T, Ogihara T. Circulating levels of calcitonin gene-related peptide in patients with medullary thyroid carcinoma. *J Clin Chem Clin Biochem* 1989;**27**:423-7.
- 26 Girgis SI, MacDonald DWR, Stevenson JC, et al. Calcitonin gene-related peptide: potent vasodilator and major product of calcitonin gene. *Lancet* 1985;ii:14-7.
- 27 Wimalawansa SJ, Morris HR, MacIntyre I. Both alpha- and beta-calcitonin gene-related peptide are present in plasma, cerebrospinal fluid and spinal cord in man. *J Mol Endocrinol* 1989;**3**:247-52.
- 28 Stevenson JC, MacDonald DWR, Warren RC, Booker MW, Whitehead MJ. Increased concentration of circulating calcitonin gene-related peptide during normal pregnancy. *BMJ* 1986;**293**:1329-30.
- 29 Foord SM, Craig RK. Isolation and characterisation of a human calcitonin gene-related receptor. *Eur J Biochem* 1987;**170**:373-9.
- 30 Denis-Donini S. Expression of dopaminergic phenotypes in the mouse olfactory bulb induced by the calcitonin gene-related peptide. *Nature* 1989;**339**:701-3.
- 31 Takami K, Kawai Y, Shio Saka S, et al. Immunohistochemical evidence for the coexistence of calcitonin gene-related peptide and choline acetyltransferase-like immunoreactivity in neurons of the rat hypoglossal, facial and ambiguus nuclei. *Brain Res* 1985;**328**:386-9.
- 32 New HV, Mudge AW. Calcitonin gene-related peptide regulates muscle acetylcholine receptor synthesis. *Nature* 1986;**323**:809-11.
- 33 Fontaine B, Klarsfeld A, Changeux J-P. Calcitonin gene-related peptide and muscle activity regulate acetylcholine receptor alpha-subunit mRNA levels by distinct intracellular pathways. *J Cell Biol* 1987;**105**:1337-42.
- 34 Takamori M, Yoshikawa HJ. Effect of calcitonin gene-related peptide on skeletal muscle via specific binding sites and G protein. *J Neurosci* 1989;**9**:99-109.
- 35 Gibson SJ, Polak JM, Katagiri T, et al. A comparison of the distribution of eight peptides in spinal cord from normals and cases of motor neurone disease with special reference to Onuf's nucleus. *Brain Res* 1988;**474**:255-78.
- 36 Girgis SI, Yates CM, Fink G, MacIntyre I. Calcitonin gene-related peptide and calcitonin immunoreactivity in brain and spinal cord in Alzheimer-type dementia. *J Neurol Sci* 1990;**99**:69-74.
- 37 Nelson MT, Huang Y, Brayden JE, Hescheler J, Standen NB. Arterial dilations in response to calcitonin gene-related peptide involves activation of K⁺ channels. *Nature* 1990;**344**:770-3.
- 38 Bunker CB, Terenghi G, Springall DR, Polak JM, Dowd PM. Deficiency of calcitonin gene-related peptide in Raynaud's phenomenon. *Lancet* 1990;**336**:1530-3.
- 39 Unger WG, Terenghi G, Ghatei MA, et al. Calcitonin gene-related polypeptide as a mediator of the neurogenic ocular injury response. *J Ocul Pharmacol* 1985;**2**:189-99.
- 40 Fuller RW, Conradson TB, Dixon CM, Crossman DC, Barnes PJ. Sensory neuropeptide effects in human skin. *Br J Pharmacol* 1987;**92**:781-8.
- 41 Williams G, Cardoso H, Ball JA, Mulderry PK, Cooke E, Bloom SR. Potent and comparable vasodilator actions of A- and B-calcitonin gene-related peptide on the superficial subcutaneous vasculature of man. *Clin Sci* 1988;**75**:309-13.
- 42 Rogers DF, Belvisi MG, Aursudkij B, Evans TW, Barnes PJ. Effects and interactions of sensory neuropeptides on airway microvascular leakage in guinea-pigs. *Br J Pharmacol* 1988;**95**:1109-16.
- 43 Brain SD, Williams TJ. Inflammatory oedema induced by synergism between calcitonin gene-related peptide (CGRP) and mediators of increased vascular permeability. *Br J Pharmacol* 1985;**86**:655-60.
- 44 Coupe MO, Mak JCW, Yacoub M, Oldershaw PJ, Barnes PJ. Autoradiographic mapping of calcitonin gene-related peptide receptors in human and guinea pig hearts. *Circulation* 1990;**81**:741-7.
- 45 McEwan J, Larkin S, Davies G, et al. Calcitonin gene-related peptide: a potent dilator of human epicardial coronary arteries. *Circulation* 1986;**74**:1243-7.
- 46 Uren NG, Davies GJ. Human alpha calcitonin gene-related peptide improves exercise workload to ischaemia and dilates stenosis of the epicardial coronary artery [Abstract]. *Eur Heart J* 1990;**11**(suppl):1070.
- 47 Johnson FG, Bell BA, Robertson IAJ, et al. Effect of calcitonin gene-related peptide on postoperative neurological deficits after subarachnoid haemorrhage. *Lancet* 1990;**335**:869-72.
- 48 McCormack DG, Mak JCW, Coupe MO, Barnes PJ. Calcitonin gene-related peptide vasodilation of human pulmonary vessels. *J Appl Physiol* 1989;**67**:1265-70.
- 49 Palmer JBD, Cuss FMC, Mulderry PK, et al. Calcitonin gene-related peptide is localised to human airway nerves and potentially constricts human airway smooth muscle. *Br J Pharmacol* 1987;**91**:95-101.
- 50 Barnes PJ. Neuropeptides in human airways: function and clinical implications. *Am Rev Respir Dis* 1987;**136**:S77-83.
- 51 Springall DR, Collina G, Barer G, Suggett AJ, Bee D, Polak JM. Increased intracellular levels of calcitonin gene-related peptide-like immunoreactivity in pulmonary endocrine cells of hypoxic rats. *J Pathol* 1988;**155**:259-67.
- 52 Leighton B, Cooper GJS. Pancreatic amylin and calcitonin gene-related peptide cause resistance to insulin in skeletal muscle in vitro. *Nature* 1988;**335**:632-5.
- 53 Leighton B, Foot FA, Cooper GJS, King JM. Calcitonin gene-related peptide-1 (CGRP-1) is a potent regulator of glycogen metabolism in rat skeletal muscle. *FEBS Lett* 1989;**249**:357-61.
- 54 Brown MJ, Morice AH. Clinical pharmacology of vasodilator peptides. *J Cardiovasc Pharmacol* 1987;**10**(suppl 12):82-7.
- 55 Struthers AD, Brown MJ, MacDonald DWR, et al. Human calcitonin gene-related peptide: a potent endogenous vasodilator in man. *Clin Sci* 1986;**70**:389-93.
- 56 Beglinger C, Born W, Hilderbrand P, et al. Calcitonin gene-related peptides I and II and calcitonin: distinct effects on gastric acid secretion in humans. *Gastroenterology* 1988;**95**:958-65.

survey found that about half the mothers of triplets and three quarters of those expecting quadruplets were initially admitted for rest alone. Women varied in their views about such a policy, some feeling reassured, others feeling anxious and stressed—particularly about the effects on their families. The report highlighted the costs to the NHS and their families of admission, an intervention without evidence of benefit.

Most reviews of managing triplet pregnancy concentrate on relating outcome to the mode of delivery.^{3,5} No consensus, however, has been reached about the optimum method as controlling for confounding variables, particularly gestational age, is difficult. In several studies only those women whose labours began very early were allowed to deliver vaginally. Although not helping to resolve this controversy, the survey by the Office of Population Censuses and Surveys highlighted the need for systematic evaluation of caesarean section. It also suggested open discussion with parents about the mode of delivery, likely neonatal problems, and possible outcomes.

Multiple births are likely to be more complicated and to occur at even earlier gestational ages than twins. Birth weights in this survey were similar to those described 40 years ago—that is, the more fetuses present the more likely they were to weigh less than 1500 g at birth.⁶ These very low birthweight infants require long periods of neonatal intensive care, which puts parents under severe strain. The report considers that liaison between hospitals, primary care, and social services could be improved. Better support—both advice and practical aid—could be provided, and this needs coordinating by someone other than the parents, which usually occurs at present.

About 5% of triplets in this review had a congenital malformation. (Although high, this figure is not directly comparable with national data.) Also of concern is the

increased prevalence of cerebral palsy and general ill health and disability in childhood. Because of the design of the follow up questionnaire and responses the authors believe that they have underestimated the size of developmental problems. They call for further long term follow up studies to assess physical and mental handicap and educational problems.

Finally, the cost to the NHS is calculated: each set of twins costs £5000, each set of triplets £12 000, and each set of quadruplets £25 000. Some 60% of the estimated cost is for neonatal care, and 20% is for antenatal care in hospital. The cost to parents of having three or more children at once is greater than having them in succession.

Selective reduction in the first trimester has been suggested as an alternative management in higher order multiple births⁷ but raises difficult legal and ethical issues.⁸ Preventing high order multiple pregnancies is clearly the best option given the high costs to mothers, babies, and services. The onus is therefore on those treating infertility to keep multiple births to a minimum.

DORIS M CAMPBELL

Senior Lecturer in Obstetrics and Gynaecology,
University of Aberdeen,
Aberdeen AB9 2ZD

- 1 Botting BJ, MacFarlane AJ, Price FV, eds. *Three, four, and more. A study of triplet and higher order births*. London: HMSO, 1990.
- 2 Hall MH. Rest in hospital and twin pregnancy. *Br J Obstet Gynaecol* 1990;97:869-71.
- 3 Campbell DM, MacGillivray I. Management of labour and delivery: higher multiples. In: MacGillivray I, Campbell DM, Thompson B, eds. *Twinning and twins*. Chichester: John Wiley and Sons, 1988:158-60.
- 4 Crowther CA, Hamilton RA. Triplet pregnancy: a 10 year review of 105 cases at Hapore Maternity Hospital, Nigeria. *Acta Genet Med Gemellol (Roma)* 1989;38:271-8.
- 5 Lipitz S, Reichman B, Paret C, et al. The improving outcome of triplet pregnancies. *Am J Obstet Gynecol* 1989;161:1279-84.
- 6 McKeown T, Record RG. Observations of fetal growth in multiple pregnancy in man. *J Endocrinol* 1952;8:386-401.
- 7 Berkowitz RL, Lynd L, Chitkara U, et al. Selective reduction of multifetal pregnancies in the first trimester. *N Engl J Med* 1988;318:1043-7.
- 8 Howie PW. Selective reduction in multiple pregnancy. Legal confusion and ethical dilemmas. *BMJ* 1988;297:433-4.

Large computer databases in general practice

Of little use unless high quality data are collected

Much research into health and disease depends on data collected from many subjects over long periods of time. Primary care is the obvious place to collect these data: general practitioners deal with most episodes of illness and have records of those episodes dealt with elsewhere. Until recently, however, the structure of general practice and the detail of recorded information have militated against such research,¹⁻³ except in trained and motivated "spotter" practices.⁴ The arrival of computers into general practice brought hopes of improvement.⁵⁻⁷

In 1987 the two main suppliers of computers to general practice, VAMP Health and AAH Meditel, offered computers to practices at low cost in return for access to aggregated patient data. (These could be sold to interested parties, pharmaceutical companies being the most obvious customers.⁸) Databases covering 1500 general practices and 3 million patients were envisaged, which could be used to examine prescribing habits. Other possibilities existed—for example, monitoring workload, preventive care,⁹ management, and post-prescription events—but recording details of these lay outside the normal contractual obligations.

The early hopes for large databases have not been fulfilled. VAMP's financial difficulties, ending its agreement to

purchase aggregated data from practices, brings the viability of these schemes into question.¹⁰ The main problem, however, has been incomplete recording. On p 766 Jick and colleagues record that in August 1988, when the VAMP scheme had enrolled 1000 doctors, only 75 practices had reached an acceptable standard of recording. Currently only one in three practices is up to standard.¹¹ In one winter month in 1989, 89 of 548 Meditel practices recorded no respiratory infections, and a further 26 could not supply data on list size. In those meeting these minimum criteria, influenza was recorded at a rate one quarter of that obtained from the weekly returns to the Birmingham research unit of the Royal College of General Practitioners (p 763).¹²

The poor quality of data is explained by practices consistently recording only what their contracts with the computer suppliers specify. All that are consistently retrievable are patients' ages and addresses, diagnoses resulting in a prescription (only about 70% of all diagnoses¹³), and the prescriptions themselves. All other data recorded by practices depend on individual preference, which varies within and among practices. The absence of medical, social, and occupational histories obviously limits the value of the databases for epidemiological research and hypothesis testing.

Perhaps their best use at present (as suggested by the two