Emergency contraception

A few days rather than a few hours to act

"Morning after"—when used to describe a form of contraception—confuses both doctors and patients by implying the need to start treatment within a few hours.¹ "Postcoital" is similarly misleading. The Family Planning Association prefers the term "emergency contraception," by which it means contraception used after intercourse but before implantation. This gives the patient and her doctor a few days rather than a few hours to act. Two methods exist: the Yuzpe method and the intrauterine contraceptive device (IUCD).

The Yuzpe method must begin within 72 hours of unprotected sexual intercourse. It entails taking four tablets of ethinylestradiol 50 µg and norgestrel 500 µg: two tablets immediately and two 12 hours later.² Overall failure rates are about 2-3%,³ with perhaps twice this failure rate when used at mid-cycle.⁴ This compares with a risk of pregnancy of about 20% when unprotected intercourse occurs at mid-cycle, 30% at ovulation, and 0-10% at other times.⁵ The Yuzpe method should not be used when absolute contraindications to oestrogens are present. (These include pregnancy, crescendo angina, transient ischaemic attacks, liver disease, undiagnosed genital tract bleeding, and a history of thromboembolism. Some absolute contraindications to long term use—such as breast cancer and arterial disease—are not contraindications for short term use.)

The Yuzpe method's main disadvantage is nausea and vomiting; another disadvantage is the risk of ectopic pregnancy when the method fails.

The IUCD may be inserted up to five days after unprotected intercourse and fails in less than 1% of cases.⁶ Although contraindicated if the patient has a history or high risk of ectopic pregnancy, an IUCD may be fitted in selected cases, providing that it is removed after the next period. Other contraindications to its use—pelvic infection or recent exposure to sexually transmitted infection—may be dealt with when the IUCD is inserted. An advantage of the IUCD is that it can remain in place as the long term method of contraception.

Women most commonly seek emergency contraception either because no contraception was used at the time of intercourse or because their chosen method—usually the sheath or cap—has failed.⁷ The need for emergency contraception is not the prerogative of the young and feckless:

women present from all age groups and walks of life.⁸ More awareness of emergency contraception could substantially reduce the numbers of abortions—perhaps by a third if the findings of a study in Tower Hamlets were repeated nationally. In this study of women attending an abortion service nearly half would have been suitable for emergency contraception, and 80% said that they would have used it had they known of it.¹

What could increase the use of emergency contraception? Firstly, society needs to accept that unprotected sexual intercourse occurs and will continue to occur regardless of its attitude towards such activity. Secondly, there should be open access to emergency contraception, with district health authorities ensuring a choice of clinics in general practice, hospitals, and the community. Access should be available, somewhere, seven days a week.⁸ Whoever coordinates district family planning and abortion services should be responsible for establishing emergency contraception services, which should receive extensive local publicity. Thirdly, all couples using contraception, especially those relying on barrier methods, should know about emergency contraception. And lastly, if the full potential of emergency contraception is to be realised then current reductions in community family planning clinics and patchy training of general practitioners and hospital doctors in contraceptive practice need urgent review.

F C READER

Honorary Senior Lecturer in Human Sexuality,
St George's Hospital Medical School,
London SW17 0RE

1 Burton R, Savage W, Reader F. The "morning after pill": is this the wrong name for it? *British Journal of Family Planning* 1990;15:119-21.

2 Yuzpe AA. Postcoital hormonal contraception: users, abusers, and abuses. *Int J Gynaecol Obstet* 1977;15:133-6.

3 Yuzpe AA, Smith RP, Rademaker AW. A multicenter clinical investigation employing ethinyl estradiol combined with dl-norgestrel as a postcoital contraceptive agent. *Fertil Steril* 1982;37:508-13.

4 Rowlands S, Guillebaud J, Bounds W, et al. Side effects of danazol compared with an ethinylestradiol/norgestrel combination when used for postcoital contraception. *Contraception* 1983;27:39-49.

5 Tietze C. Probability of pregnancy resulting from a single unprotected coitus. *Fertil Steril* 1960;11:485-8.

6 Friedman EHJ, Rowley DEM. Postcoital contraception—a two year evaluation service. *British Journal of Family Planning* 1987;13:139-44.

7 Yuzpe AA, Lances WJ. Ethinylestradiol and dl-norgestrel as a postcoital contraceptive. *Fertil Steril* 1977;28:932-6.

8 Bromwich PD, Parsons AD. The establishment of a postcoital contraception service. *British Journal of Family Planning* 1982;8:16-9.

A little something for the HIV epidemic?

General practitioners should be able to prescribe condoms on FP10s

General practitioners are ideally placed to counsel their patients on the risks of acquiring HIV. Although many are keen to care for HIV positive patients and help prevent the spread of the virus,¹ few discuss the subject with their patients. One study estimated that HIV is mentioned in only one in 220 consultations,² and where discussion of risk behaviour and HIV could logically be expected—accompanying general practitioners' contraceptive advice—it seems largely absent. More therefore needs to be done to help general practitioners to help those who are at risk from HIV. Enabling them to prescribe condoms on FP10s would be a start.³

Latex condoms prevent the transmission of many sexually transmitted diseases.⁴ Mechanical models have confirmed their effectiveness as physical and chemical barriers against HIV,^{5,6} and nonoxinol 9, incorporated in the lubricant of some condoms, may be useful if this barrier breaks, by inactivating HIV.^{7,8}

But condoms have many negative associations in Western culture. They are associated with furtive sexual activity in the adolescent and sexual incontinence in the adult. They may symbolise a relationship in which something vital is held back, and suggesting their use to prevent infection may be