DETECTION AND MANAGEMENT OF CONGENITAL ABNORMALITIES—I

Geoffrey Chamberlain

IN CONFIDENCE Certificate A

Not to be destroyed within three years of the date of operation

ABORTION ACT 1967

Certificate to be completed before an abortion is performed under Section 1(1) of the Act

I, _____ (Name and qualifications of practitioner in block capitals)

of _____ (Full address of practitioner)

Have/have not* seen/and examined* the pregnant woman to whom this certificate relates at _____ (Full address of place at which patient was seen or examined)

on _____ and I _____ (Name and qualifications of practitioner in block capitals)

of _____ (Full address of practitioner)

Have/have not* seen/and examined* the pregnant woman to whom this certificate relates at _____ (Full address of place at which patient was seen or examined)

on _____

We hereby certify that we are of the opinion, formed in good faith, that in the case of _____ (Full name of pregnant woman in block capitals)

of _____ (Usual place of residence of pregnant woman in block capitals)

(Ring appropriate number(s))

1. the continuance of the pregnancy would involve risk to the life of the pregnant woman greater than if the pregnancy were terminated;
2. the continuance of the pregnancy would involve risk of injury to the physical or mental health of the pregnant woman greater than if the pregnancy were terminated;
3. the continuance of the pregnancy would involve risk of injury to the physical or mental health of the existing child(ren) of the family of the pregnant woman greater than if the pregnancy were terminated;
4. there is substantial risk that if the child were born it would suffer from such physical or mental abnormalities as to be seriously handicapped.

This certificate of opinion is given before the commencement of the treatment for the termination of pregnancy to which it refers and relates to the circumstances of the pregnant woman's individual case.

Signed _____ Date _____

Signed _____ Date _____

*Where appropriate
Form A 2.1.1
H.M.S.O. 1967. 5000. 5/62. 4 P. 170

Certificate A of the Abortion Act 1967.

A congenital abnormality in their forthcoming baby is greatly feared by couples; we are not many generations away from the superstitious who looked on malformation as a retribution for misbehaviour. Congenital abnormalities are major causes of perinatal mortality and morbidity.

The known causes of abnormality are genetic or environmental. Genetic abnormality depends on the chromosomes we get from our parents together with the breaks and realignments occurring at fertilisation. Maternal aging increases the risk of abnormalities in genes. A good account of genetic abnormalities is found in the recent *ABC of Clinical Genetics*.

Environmental factors interfere with embryonic development at a precise stage of organogenesis. They are difficult to pinpoint and often are misassociated. Obvious insults such as exposure to thalidomide, x rays, and rubella can be identified; more difficult is the precise place of factors such as organic solvents in the cleaning industry and viral infections such as chickenpox.

An antenatal service should aim at diagnosing congenital abnormalities as early as possible. Though the ideal treatment is prevention, this is usually too late by the time the woman joins the antenatal clinic. If an abnormality can be detected early the couple may be offered the choice of a termination of pregnancy. This is allowed under statutory grounds IV of the Abortion Act 1967, and about 1500 abortions a year occur in England and Wales for this reason.

Not all couples want to abort their unborn child even if it has an abnormality. Antenatal diagnostic facilities should be made available not only to those who agree to a termination of pregnancy if an abnormality is found but to all affected couples for the following reasons:

- It gives the couple more time to accustom themselves and other children in the family to the idea that an abnormal child is to be born
- If the abnormality is not lethal, early diagnosis allows plans to be made for the woman to be delivered in a centre where full treatment may be given early
- Certain abnormalities that entail back pressure on essential organs such as the kidneys may be amenable to intrauterine decompression operations. This is less useful than first thought for, although the immediate physical pressure can be relieved, there are often underlying abnormalities of the organs themselves which lead to poor results in the long term.

In the early 1990s antenatal screening for congenital abnormalities is concerned with the detection of malformations of the central nervous system and abnormalities of chromosomal origin.

Testing in early pregnancy

Chorionic villus biopsy. Under ultrasound guidance there can be (a) a transcervical approach of the cannula to the edge of the developing placenta or (b) a transabdominal aspiration by needle from the middle of the trophoblast mass.

Risks of Down's syndrome. The risk increases after 35 and sharply after 40. The percentage of women in the United Kingdom who deliver by each age group is also shown. Although the risk is high after 40, the numbers of women delivering are small.

The risk of Down's syndrome in a woman of 40 according to serum α fetoprotein concentration at 14-20 weeks. The actuarial risk by age in a woman of 40 is 1:100.

Chromosomal problems

To examine chromosomes a small amount of fetal tissue is required. In chorionic villus sampling a minute piece of trophoblast tissue is removed for examination of the chromosomes in the cell nucleus and, with increasing confidence, DNA assessment. Such sampling is commonly performed at 9-11 weeks of gestation, and a result is available a couple of days after the test (compared with the delay of at least 21 days for amniocentesis). Sampling is under ultrasound control and is by the transcervical or the transabdominal route, depending on the site of placental implantation. Recent reports of fetal damage being associated with the transabdominal approach have led to a reduction in its use. Most obstetricians await the results of a large randomised controlled trial, which are due later this year.

The apparent attraction of early chorionic villus sampling and its quicker result are offset by the higher risks of stimulating a miscarriage. Abortion rates associated with chorionic villus sampling are 2-4% compared with 0.3-1.0% with amniocentesis. At 10 weeks of gestation, however, the rate of spontaneous miscarriage is biologically much higher than at 16 weeks, so the comparison is not only of techniques. Many obstetrics units are now using chorionic villus sampling for women at high risk, and with experience the miscarriage rates would be expected to fall.

A nationally organised randomised controlled trial is in process to sort out the comparative advantages of chorionic villus sampling and amniocentesis. Many hospitals offering such sampling are in this trial and so general practitioners may find that some of their patients referred for the test will have been randomised to the amniocentesis option. They should check with their local department whether it is part of this Medical Research Council trial, whose results are expected later this year.

Trisomy 21 (Down's syndrome) is much commoner in women over 35, but half of the babies with this condition are born to women under that age. Although the risks to mothers under 35 are less, the number of babies is much greater. To negate this, simple screening is required as both chorionic villus sampling and amniocentesis are unsuitable and invasive procedures that are very labour intensive. Hence the recent work assessing a combination of biochemical tests of maternal blood in early pregnancy to screen for Down's syndrome is of wide interest. Maternal α fetoprotein, human chorionic gonadotrophin, and oestriol concentrations are measured and compared with background data derived from large populations. The predictive value of the combination of tests is high so that a population can be pinpointed which is at a high risk of Down's syndrome. If that risk is greater than one in 100, it is then recommended that amniocentesis be performed to assess the fetal chromosomes. This test is new but is being tried in some health districts and has great promise of screening for Down's syndrome in younger mothers.

Certificate A of the Abortion Act 1967 has Crown copyright and is reproduced by permission of the Controller of Her Majesty's Stationery Office. The data for the graph showing the risk of Down's syndrome by α fetoprotein concentration are reproduced by permission of Churchill Livingstone from *Obstetrics* edited by A Turnbull and G Chamberlain.

Professor Geoffrey Chamberlain, FRCOG, is chairman of the department of obstetrics and gynaecology at St George's Hospital Medical School, London.