services combining medical and psychiatric expertise is needed.

1 Eastwood R, Trevelyan MH. Relationship between physical and psychological disorder. *Psychol Med* 1972;2:363-7.
2 Wells KB, Golding JM, Burnam MA. Affective, substance use, and anxiety disorders in persons with arthritis, diabetes, heart disease, high blood pressure, or chronic lung conditions. *Gen Hosp Psychiatry* 1989;11:320-7.
3 Wells KB, Golding JM, Burnam MA. Chronic medical conditions in a sample of the general population with anxiety, affective, and substance use disorders. *Am J Psychiatry* 1989;146:1440-6.
4 Goldberg D, Huxley P. *Mental illness in the community*. London: Tavistock, 1980.
5 Stewart AL, Greenfield S, Wells K, Rogers WH, Berry SD, Hays RD, et al. Functional status and well-being of patients with chronic conditions. *JAMA* 1989;262:907-13.
6 Mayou RA, Hawton KE. Psychiatric disorder in the general hospital. *Br J Psychiatry* 1986;149:172-96.
7 Regier DA, Goldberg ID, Taube CA. The de facto US mental health services system. *Arch Gen Psychiatry* 1978;35:685-93.
8 Kessler LG, Burns BJ, Shapiro S, Tischler GL, George LK, Hough RL. Psychiatric diagnoses of medical service users: evidence from the epidemiologic catchment area program. *Am J Public Health* 1987;77:18-24.
9 Hankin JR, Steinwachs DM, Regier DA, Burns BJ, Colburn ID, Hooper EW. Use of general medical care services by persons with mental disorders. *Arch Gen Psychiatry* 39:225-31.
10 Berkson J. Limitations of the application of fourfold table analysis to hospital data. *Biometrics Bulletin* 1946;2:47-53.
11 Barsky AJ, Wyshak G, Klerman GL. Medical and psychiatric determinants of out-patient medical utilization. *Med Care* 1986;24:548-60.

12 Fulop G, Strain JJ, Fahs MC, Hammer JS, Lyons JS. Medical disorders associated with psychiatric comorbidity and prolonged hospital stay. *Hosp Community Psychiatry* 1989;40:80-2.
13 Mayou RA, Hawton K, Feldman E. What happens to medical patients with psychiatric disorder? *J Psychosom Res* 1988;32:541-9.
14 Wallen J, Pincus HA, Goldman HH, Marcus SE. Psychiatric consultations in short-term general hospitals. *Arch Gen Psychiatry* 1987;44:163-8.
15 Hengeveld MW, Rooyman HGM, Vecht-van den Bergh R. Psychiatric consultations in a Dutch university hospital: a report on 1814 referrals, compared with a literature review. *Gen Hosp Psychiatry* 1984;6:271-9.
16 Acheson ED. *Medical record linkage*. Oxford: Oxford University Press, 1967.
17 Gill LE, Baldwin JA. Methods and technology of record linkage: some practical considerations. In: Baldwin JA, Acheson ED, Graham WJ, eds. *Textbook of medical record linkage*. Oxford: Oxford University Press, 1987;3:39-54.
18 Colman M, Douglas A, Herman C, Peto J. Cohort study analysis with a FORTRAN computer programme. *Int J Epidemiol* 1986;15:134-7.
19 Devis T. Population movements measured by the NHS central register. *Population Trends* 1984;36:18-24.
20 Breslow NE, Day NE. *Statistical methods in cancer research II—the design and analysis of cohort studies*. Lyons: International Agency for Research on Cancer, 1987;68-9, 94-6.
21 Rang EH, Fairbairn AS, Acheson ED. An enquiry into the evidence and prognosis of undiagnosed abdominal pain treated in hospital. *British Journal of Preventive and Social Medicine* 1970;24:47-51.
22 Fitzpatrick RM, Hopkins A. Referrals to neurologists for headaches not due to structural disease. *J Neurol Neurosurg Psychiatry* 1987;44:1061-7.
23 Mayou RA. Atypical non-cardiac chest pain. *J Psychosom Res* 1989;33:393-406.

(Accepted 23 July 1991)

Passive smoking and otitis media with effusion

G S Barr, A P Coatesworth

Department of
Otolaryngology, Queen
Elizabeth Hospital,
Birmingham B15 2TH
G S Barr, FRCS, senior
registrar
A P Coatesworth, MB, senior
house officer

Correspondence to:
Mr G S Barr, Department of
Otolaryngology, Walsgrave
Hospital, Coventry
CV2 2DX.

BMJ 1991;303:1032-3

Otitis media with effusion is reported to occur in over 80% of children at some stage, but surgery is indicated only when the condition is unresolved after three months. The main underlying causes of otitis media with effusion are eustachian tube malfunction, alteration of the mucociliary system, and nasopharyngeal disproportion. A family history of otitis media in parents or siblings and parental occupation and smoking may also be risk factors.¹ Hinton² found a higher proportion of parents who smoked among 115 children undergoing surgery for otitis media with effusion than among 35 children attending an orthoptic clinic. A study of 892 7 year old schoolchildren in Edinburgh found an association between salivary cotinine concentrations from passive smoking and the presence of tympanometric abnormalities.³ Such abnormalities, however, are not necessarily associated with otitis media with effusion.

Otitis media with effusion is diagnosed from the history and by otoscopy, audiometry, and tympanometry. No single entity is completely specific or sensitive and the condition is confirmed by myringotomy. We evaluated the relation between parental smoking habit and the presence of the established condition in a case-control study.

Subjects, methods, and results

In all, 115 children (70 boys, 45 girls; age range 17 months to 11 years 6 months, median 5 years 5 months) from the Cheltenham and Gloucester areas who had otitis media with effusion confirmed by myringotomy were matched according to age (within six months), sex, race, and social class to a control group of healthy children attending the ophthalmology and orthopaedic clinics. The children with otitis media with effusion had had hearing loss for at least three months and had been assessed by otoscopy, tympanometry, and audiometry. The control group had no history of ear problems and normal results on otoscopy and tympanometry with a portable Welch Allyn tympanometer.

The smoking habits of the parents of the children in

the two groups were recorded after we had explained the aims of our study to them. Data were compared by McNemar's test for the presence of at least one adult in the household who smoked and whether the mother smoked. The differences between the number of cigarettes smoked by mothers and by all of the adults in the household were calculated by comparing the median values of paired data by the binomial method.

There were 230 adults in the study group and 228 adults in the control group. Seven patients were from socioeconomic class I, 14 from class II, 38 from class III, 43 from class IV, and 13 from class V. Parental smoking habits in the two groups were the same. There were no differences between the median number of cigarettes smoked in the two groups by mothers alone and by all adults in the household (95% confidence interval 0 to 0 cigarettes for both sets of data). The 95% confidence interval of the difference in proportion of mothers who smoked was -0.08 to 0.16 and that for all adults in the household -0.19 to 0.05 (McNemar's test; table).

Parental smoking habits among 115 matched pairs of healthy control children and children with otitis media with effusion

Group	Smokers present	No of pairs	
		Mother only	All adults in household
Otitis media	No	46	29
Control	No		
Otitis media	No	22	29
Control	Yes		
Otitis media	Yes	27	21
Control	No		
Otitis media	Yes	20	36
Control	Yes		
Total		115	115

Comment

Nasal symptoms, particularly those related to adenoid hypertrophy, are associated with the development of otitis media with effusion. Histamine concentrations in adenoid tissue are proportional to size, but ultrastructural evidence shows that the morphology of adenoid mast cells is the same in children with and without otitis media with effusion (A B Drake-Lee, unpublished data). Exposure to cigarette smoke might induce instability of the mast cell walls and the onset of otitis media with effusion² but our data do not support this hypothesis.

The prevalence of otitis media with effusion is highest where social conditions are poor,⁴ and children of non-manual workers have significantly better hearing than do those of manual workers.⁵ Cigarette smoking is commoner in those from the poorer socioeconomic classes but it is unlikely to be a risk factor for otitis media effusion, although it may have an association.

We thank Messrs G C Fox, M Hardingham, and J M Robinson for allowing us to study their patients.

- 1 Maw AR. Otitis media with effusion (glue ear). In: Kerr AG, Groves J, Evans NG, eds. *Scott-Brown's otolaryngology*. 5th ed. Vol 6. London: Butterworth, 1987:159-76.
- 2 Hinton AE. Surgery for otitis media with effusion in children and its relationship to parental smoking. *J Laryngol Otol* 1989;103:559-61.
- 3 Strachan DP, Jarvis MJ, Feyerabend C. Passive smoking, salivary cotinine concentrations, and middle ear effusion in 7 year old children. *BMJ* 1989;298:1549-52.
- 4 Robinson GC, Anderson DO, Moghadam HK, Cambon KG, Murray AB. A survey of hearing loss in Vancouver school children. Part 1. Methodology and prevalence. *Can Med Assoc J* 1967;97:1199-207.
- 5 Richardson K, Peckam CS, Goldstein H. Hearing levels of children tested at 7 and 11 years. A national study. *Br J Audiol* 1976;10:117-23.

(Accepted 24 June 1991)

Menarche and reproduction after treatment for African Burkitt's lymphoma

Janet E Neequaye, Julianne Byrne, Paul H Levine

Burkitt's Tumor Project, Department of Child Health, University of Ghana Medical School, Accra, Ghana
Janet E Neequaye, MRCP, associate professor of child health

Epidemiology and Biostatistics Program, National Cancer Institute, National Institutes of Health, Bethesda, Maryland, USA
Julianne Byrne, PHD, senior staff fellow
Paul H Levine, MD, senior investigator

Correspondence to:
Dr J E Byrne, EPN 400, National Cancer Institute, NIH, Bethesda, MD 20892, USA.

BMJ 1991;303:1033

African Burkitt's lymphoma is a rapidly progressing cancer that responds quickly to cyclophosphamide.¹ Since 1966 the Burkitt's Tumor Project in Accra has treated children with Burkitt's lymphoma chiefly by cyclophosphamide. A troublesome side effect after cyclophosphamide, however, is sterility.² We sought to determine how chemotherapy might influence fertility in adults who had been treated for Burkitt's lymphoma as children.

Patients, methods, and results

Between 1984 and 1988 we attempted to interview and draw blood from every person treated at the University of Ghana Medical School for Burkitt's lymphoma between 1966 and 1988 who was at least 16 in 1988 and had survived at least five years from date of diagnosis. Because of difficulty in tracing boys results are given for girls only. Survivors were diagnosed at an average age of 9 years (range 3-17). All except three were prepubertal at diagnosis. All children were treated with cyclophosphamide and some also with other drugs. Cyclophosphamide was given fortnightly in boluses of 1400 mg/m² intravenously until remission, which was usually after three doses. The total dose ranged from 2.8 g/m² in two patients to roughly 9.0 g/m² in five (90-300 mg/kg).

Blood was sampled at the time of interview and date of the last menstrual period noted. Serum follicle stimulating hormone and luteinising hormone concentrations were measured by double antibody radioimmunoassay using the second international reference preparation as standard.

Most former patients lived in small rural villages with no access to postal or telephone systems. After excluding male patients and those who could not be located we interviewed 24 female survivors (two by proxy) and 13 sisters or neighbours as controls.

At follow up no survivor was still having treatment. All but two of the women were in good general health. One was blind and another had a chronic foot ulcer due to sensory deficit accompanying paraplegia. Both problems resulted from the tumour.

Survivors were similar to controls in age at interview, years of schooling, and sexual and marital experiences (table). Survivors, however, reached menarche significantly later than controls (age 13.5 v 12.2 years; $p<0.05$). This could not be attributed to treatment during puberty as the average age at menarche among the 16 survivors treated before age 11 was 13.3 years.

Of the 22 women trying for children, 21 had had at

least one pregnancy. One woman who was treated for three weeks at age 11, reached the menarche at 16, and was 24 at follow up reported irregular periods and infertility for seven years. Sixteen of 17 married survivors had at least one child. Blood from 13 women was tested for follicle stimulating hormone and luteinising hormone concentrations. In none of these women were values abnormal.

Reproductive experiences of female survivors of African Burkitt's lymphoma and controls

	Survivors	Controls
Total No studied	24	13
No (%) with menstrual problems	1 (4)	0
No (%) ever had sexual intercourse	23 (96)	12 (92)
No (%) married	17 (71)	12 (92)
No (%) married with children	16/17 (94)	12/12 (100)
Average age at first live birth (years)	18.6	19.5
Average age at menarche (years)	13.5	12.2*
Average age at marriage (years)	18.4	18.6

* $p<0.05$ (t test).

Comment

In this small study most female survivors of Burkitt's lymphoma seemed to be fertile. The single potential problem noted was the delay in menarche by more than one year. So far as we know this is the first such report after treatment with cyclophosphamide, which previously has been associated with dose dependent gonadal dysfunction.³ The doses received in our study were just below that range.³ To date the longest follow up of girls treated with cyclophosphamide (for the nephrotic syndrome) indicated normal fertility in 16 of 18 women in their mid-20s, but age at menarche was not studied.⁴

Follow up studies of survivors of cancer treated with various agents found changes in timing of menarche, some recording an earlier age at menarche,⁵ others a later age. Reasons for this effect are unclear. We also do not know what implication this would have for the timing of menopause.

This study was supported by the National Cancer Institute, National Institutes of Health, Bethesda (contract No NO1-CP-51009). We acknowledge the staff of the Burkitt's Tumor Project, Accra, for their help and in particular Mrs Emelia Buadoo, who interviewed the subjects, often under difficult circumstances.

- 1 Burkitt DP. Long-term remissions following one and two-dose chemotherapy for African lymphoma. *Cancer* 1967;20:756-9.
- 2 Koyama H, Wada T, Nishizawa Y, Iwanaga T, Aoki Y, Terasawa T, et al. Cyclophosphamide-induced ovarian failure and its therapeutic significance in patients with breast cancer. *Cancer* 1977;36:1401-9.
- 3 Rivkees SA, Crawford JD. The relationship of gonadal activity and chemotherapy-induced gonadal damage. *JAMA* 1988;259:2123-5.
- 4 Watson AR, Taylor J, Rance CP, Bain J. Gonadal function in women treated with cyclophosphamide for childhood nephrotic syndrome: a long-term follow-up study. *Fertil Steril* 1986;46:331-3.
- 5 Quigley C, Cowell C, Jimenez M, et al. Normal or early development of puberty despite gonadal damage in children treated for acute lymphoblastic leukemia. *N Engl J Med* 1989;321:143-51.

(Accepted 29 May 1991)