- 5 McLellan DL. Rehabilitation. *BMJ* 1991;303:355-7.
- 6 Department of Health. *Priorities and planning guidance for the NHS for 1992/93*. London: DoH, 1991. (EL(91)103.)
- 7 House of Commons Select Committee on Health. *Public expenditure on health and social services. Third report from the health committee: session 1990/91*. Vol 1. London: HMSO, 1991. (Paper 614.)
- 8 Sullivan LW. Foreward, In: *Healthy people 2000*. Washington, DC: Department of Health and Human Services, 1990.
- 9 Health Education Authority. *Health update 1: coronary heart disease*. London: HEA, 1990.
- 10 Social Survey Division, Office of Population Censuses and Surveys. *Smoking among secondary school children in 1990*. London: HMSO, 1991.
- 11 Department of Health. *Research for health: a research and development strategy for the NHS*. London: HMSO, 1991.
- 12 Smith A. *A national health service and the public health*. Harrogate: Yorkshire Regional Health Authority, 1991.

Junior Doctors. The New Deal

Using the potential of all staff

Angela Thomas

This is the last in a series of articles which explore the new deal on junior doctors' hours of work and explain how it will be implemented.

There are a host of other factors apart from long hours which cause low morale and stress among junior doctors. Some affect patient care directly and some affect the doctor alone. *The New Deal* contains important agreements which address these problems.¹

A recurrent complaint among juniors is the varying role that nurses play in different disciplines and in different hospitals. In some hospitals clinical work is shared between nursing or midwifery staff and medical staff—for example, in accident and emergency, oncology, gastroenterology, and maternity units. In

increasing headache as ward closures and the increased turnover of patients make placing of newly admitted patients difficult and time consuming for the junior doctors.

In the past these problems have not been solved because of the lack of priority given to them in the allocation of resources. Hard pressed haematology laboratories cut staff to save costs and lose their in-patient phlebotomists and cardiac departments lose their team of technicians, but the junior doctor is always there to do the jobs at no extra cost. A central organisation for locating beds run by a designated and appropriately qualified team of clerical staff would be of enormous benefit and would give junior doctors several more hours a day to devote to their training. Again, all members of the ministerial group were convinced that sorting these problems out, although not directly decreasing hours, would decrease the workload, improve the efficiency of junior doctors, and do much to alleviate the frustration and disillusionment that many now feel. The representatives of NHS management accepted that there would need to be a reallocation of resources to allow for the employment of the necessary support staff but they were enthusiastic that this should be done.

The Secretary of State for Health said when the deal was launched that the resources to achieve the hours deal would be made available.

others this does not happen and nursing staff are unable to use their skills to the full. This is particularly frustrating for them and for junior doctors if they have just moved from a hospital or department where an extended role is practised. A doctor can be disturbed at night simply to give an intravenous injection which could more easily and with less delay be given by the nurse on the ward. There are many other tasks which could be more appropriately undertaken by nursing or midwifery staff to the benefit of patients, nurses, and doctors but which are not because of a lack of agreed local policy. This leads to frustration and often resentment among doctors and nurses. There is scope for an extension of shared care and the ministerial group on junior doctors' hours has recognised this.²

"Making the best use of the skills of nurses and midwives" gives guidance on the need to review local policies on tasks which appropriately qualified nurses and midwives could reasonably undertake to improve the quality of care. The guidance is supported by all those participating in the discussions on hours, including the chief nursing officer and should enable all staff to be used to their full potential.

Another problem which *The New Deal* addresses³ and which the University of London has recently highlighted⁴ is the question of junior doctors spending more time than necessary on clerical and administrative tasks and too much routine technical work such as phlebotomy. London University has clearly stated that posts where junior doctors do too much of this sort of work may well have training approval withdrawn and some other universities have followed suit. A certain amount of technical work does have a place in the early training of a junior doctor but this does not justify requiring them to do two to three hours of blood taking before the normal day can start. Finding beds is an

Scandalously inadequate

Hospital is home for many junior doctors, particularly the most junior, but their living conditions are often scandalously inadequate, despite guidance on accommodation from the Department of Health, the latest in 1986 after vigorous representations by the Hospital Junior Staff Committee (now Junior Doctors Committee). Many residential areas, on call rooms, catering facilities, and communal areas still fall short of the minimum standards set by the government. Recently training recognition of juniors' posts was withdrawn at one hospital in London because of poor facilities. Surely it is not too much to expect clean linen in the on call room when you are on duty?

In an attempt to improve the position guidance on living and working conditions and job descriptions, which form part of the doctors' contract, has been issued under the new agreement.⁵ Is this guidance any more likely to be followed than the last? I believe so because, firstly, it has the full backing of the ministerial group, including the Minister for Health, and, secondly, there is now a path which leads back directly to the group if the guidance is not followed. If problems occur junior doctors will be able to raise them at national level.

All the above initiatives depend on local activity and commitment followed by local cooperation and agreement. If problems fail to be resolved there is a way that pressure can be exerted to bring about change. The regional task forces are charged with monitoring the progress of the implementation of all aspects of the

Sunley Research Centre,
London W6 8LW
Angela Thomas, MRCPATH,
senior registrar in haematology

BMJ 1991;303:1123-4

heads of agreement and not just hours of work. If satisfactory solutions are not found because of a shortage of money task forces can report this directly to the ministerial group, which will continue to meet every six months; the government will have to find the necessary resources.

Responsibility for providing information on all aspects of the agreement now lies partly with junior doctors. The BMA has sent its junior members a booklet explaining the new deal in detail, what information is required, and whom to contact for help.⁶ The Secretary of State for Health said when the deal was launched that the resources to achieve the hours deal would be made available. The aspects of the agreement I have referred to must be seen as part of a package of measures which, if implemented with the agreed

hours' reductions, should result in improved patient care, improved training for juniors, more efficient working, and perhaps even a few more hot meals at night.

- 1 NHS Management Executive. *Junior doctors. The new deal*. London: NHS Management Executive, 1991.
- 2 NHS Management Executive. *Junior doctors. The new deal. Making the best use of the skills of nurses and midwives*. London: NHS Management Executive, 1991.
- 3 NHS Management Executive. *Junior doctors. The new deal. Making better use of technical, administrative, clerical, and other support staff and systems*. London: NHS Management Executive, 1991.
- 4 University of London. *Inappropriate duties for preregistration house officers*. London: University of London, 1991.
- 5 NHS Management Executive. *Junior doctors. The new deal. Living and working conditions of doctors in training*. London: NHS Management Executive, 1991.
- 6 BMA Junior Doctors Committee. *Junior doctors. The new deal*. London: BMA, 1991.

Health and the Environment

Importance of a healthy environment

Fiona Godlee, Alison Walker

This is the first in a series of articles which examine the impact of environmental pollution on human health.

Doctors seem to spend their time these days exhorting their patients to adopt healthier habits: to exercise more, eat less, drink only a little and smoke not at all; to wear seat belts, crash helmets, and condoms. But perhaps the most important risks to health are beyond people's immediate control, caused by the unhealthy habits not of individuals but of an energy hungry and throwaway society.

The environmental price

Since the industrial revolution and the advent of the internal combustion engine we have consumed the earth's natural resources and altered our environment at a rate never seen before. The demand for energy is constantly growing, fuelled by the world's escalating population—now at 5.3 billion and estimated to reach 10 billion by 2050.¹ In 1989 the world's energy consumption reached 9.3 million tons of oil equivalent, causing the emission of nearly six billion tons of carbon into the environment (figure).² The World Energy Conference concluded that by 2020 the world would be using 75% more energy, and most of it would still be supplied by coal, oil, and nuclear power.³

The destructive consequences of human industry on the environment are in many cases already evident, in others controversial. Chlorofluorocarbons, still used as a propellant in aerosols in some countries and in the manufacture of refrigerators, are destroying the layer of ozone in the outer atmosphere which protects the earth from the sun's ultraviolet rays. If levels of chlorofluorocarbons continue to rise at the present rate—about 3% a year—a 6% reduction in stratospheric ozone has been predicted by 2040.⁴ The resultant increase in the amount of ultraviolet B reaching the earth's surface—1.5–2% for every 1% reduction in total column ozone—has the potential to cause ecological catastrophe. Phytoplankton, for example, the basis of the food chain for all aquatic creatures, are highly vulnerable to damage from ultraviolet B radiation.⁵

Greenhouse gases—carbon dioxide from the burning of fossil fuels, methane, nitrous oxide, and chlorofluorocarbons—are accumulating in the outer atmosphere, where concentrations of carbon dioxide are now higher than at any time in the past 16 000 years.⁶ Global warming may be the consequence. The

earth's surface temperature has risen by about 0.6°C in the past 100 years and further increases of 2–5°C have been predicted over the next 50 to 100 years, causing potentially catastrophic rises in sea level.⁵

Road vehicles, responsible for emitting a fifth of the carbon dioxide in air, also produce acidic gases—nitrogen oxides and sulphur dioxide—which cause acid rain. Though heavy industry is bringing its emissions under control, the use of road transport is increasing at a rate likely to swamp any reductions in pollution achieved by technological improvements in car engines. The Department of Transport estimates that in Britain the number of vehicle miles will double by 2025, causing an estimated doubling of emissions of carbon dioxide.⁷ For the same reason emissions of nitrogen oxides, which will fall over the next 20 years owing to the introduction of catalytic converters, will be on the increase again by 2020.

Acid rain lays waste large areas of agricultural land and forest, adding to deforestation caused by harvesting hardwood and the pressure for more agricultural land. Each year an area of tropical forest the size of Austria is destroyed.¹ Deforestation increases the build up of carbon dioxide because trees fix carbon from the air. It also spells disaster for biological diversity as more than half the species on earth live in the tropical rain forests.⁸

The human price

Man, the main cause of environmental degradation, is also one of its victims. Apart from the direct effects that increases in ambient temperature might have on health—higher mortality from heart and lung disease; the spread of tropical diseases to temperate climates; more deaths, disability, and psychological problems due to storms and natural disasters—global warming would cause the sea to expand and swamp large areas of habitable and agricultural land.⁹ A rise in sea level of one metre (the International Panel on Climate Change's 1990 estimate¹⁰) would cover five million square kilometres of the world's lowest lying land, causing the loss of one third of our crop growing areas and creating 50 million environmental refugees.⁶

Nuclear energy reduces reliance on fossil fuels but creates problems of its own. The fire at Chernobyl nuclear power station in 1986 is still having repercus-

British Medical Journal,
London WC1H 9JR
Fiona Godlee, MRCP,
assistant editor
Alison Walker, MRCP, former
editorial registrar

Correspondence to: Dr
Godlee.

BMJ 1991;303:1124-6