- 8 McNamee P, Celona J. *Decision analysis for the professional with Supertree*. Redwood City, California: Scientific Press, 1987.
- 9 Pauker SP, Pauker SG. Prenatal diagnosis: a directive approach to genetic counselling using decision analysis. *Yale J Biol Med* 1977;50:275-89.
- 10 Bingham P, Lilford RJ. Management of the selected term breech presentation: assessment of the risks of selected vaginal delivery versus Cesarean section for all cases. *Obstet Gynecol* 1987;69:965-78.
- 11 Hillner BE, Smith TJ. Efficacy and cost effectiveness of adjuvant chemotherapy in node negative breast cancer. *N Engl J Med* 1991;324:160-8.
- 12 Klein K, Pauker SG. Recurrent deep venous thrombosis in pregnancy: analysis of the risks and benefits of anticoagulation. *Med Decision Making* 1981;1:181-202.
- 13 Elstein AS, Holzman GB, Ravitch MM, Metheny WA, Holmes MM, Hoppe RB, et al. Comparison of physicians decisions regarding estrogen replacement for menopausal women and decisions derived from a decision analytic model. *Am J Med* 1986;80:246-58.
- 14 Feldman GB, Freiman JA. Prophylactic caesarean section at term? *N Engl J Med* 1985;312:1264-7.
- 15 Weinstein MC, Pliskin JS, Stason WB. Coronary artery bypass surgery: decision and policy analysis. In: Bunker JP, Barnes BA, Mosteller F, eds. *Costs, benefits and risks of surgery*. Oxford: Oxford University Press, 1977.
- 16 Tompkins RK, Burnes DC, Cable WE. An analysis of the cost effectiveness of pharyngitis management and acute rheumatic fever prevention. *Ann Intern Med* 1977;86:481-92.
- 17 Neutra R. Indications for the surgical treatment of suspected acute appendicitis. A cost effectiveness approach. In: Bunker JP, Barnes BA, Mosteller F, eds. *Costs, benefits and risks of surgery*. Oxford: Oxford University Press, 1977.
- 18 Neuhauser D, Lewicki AM. What do we gain from the sixth stool guaic? *N Engl J Med* 1975;293:226-8.
- 19 Van Crevel H, Habbema JDF, Braakman R. Decision analysis of the management of incidental intracranial saccular aneurysms. *Neurology* 1986;36:1335-9.
- 20 Lilford RJ. Trade off between gestational age and miscarriage risk of prenatal testing: does it vary according to genetic risk? *Lancet* 1990;336:1303-5.
- 21 Simes RJ. Treatment selection for cancer patients: application of statistical decision theory to the treatment of advanced ovarian cancer. *J Chron Dis* 1985;38:171-86.
- 22 Sacks HS, Berrier J, Reitman D, Ancona-Berk VA, Chalmers TC. Meta-analysis of randomised controlled trials. *N Engl J Med* 1987;316:450-5.
- 23 Lilford RJ. Clinical trial numbers. *Lancet* 1990;335:483-4.
- 24 Thornton JG, Bryce FC, Bhabra K. Withholding intensive care from premature babies. *Lancet* 1988;ii:332-3.
- 25 De Dombal FT. Computer aided diagnosis of acute abdominal pain. *BMJ* 1972;ii:9-13.
- 26 Adams ID, Chan M, Clifford PC, Cooke WM, Dullos V, de Dombal FT, et al. Computer aided diagnosis of abdominal pain: a multicentre study. *BMJ* 1986;293:800-4.
- 27 Elstein AS, Bordage G. Psychology of clinical reasoning. In: Dowie J, Elstein A, eds. *Professional judgement. A reader in clinical decision making*. Cambridge: Cambridge University Press 1988:109-29.
- 28 Balla JJ, Elstein AS, Christensen C. Obstacles to acceptance of clinical decision analysis. *BMJ* 1989;298:579-82.

(Accepted 29 December 1991)

Basic Molecular and Cell Biology

Gene regulation

David S Latchman

This is the first of two articles updating the series "Basic Molecular and Cell Biology" published in 1987. The articles will be published in a new edition of the book of the series later this year.

That the expression of human genes must be a highly regulated process should be clear to anyone who has ever dissected a human body. The vast range of different tissues and organs differ dramatically from each other and they all synthesise different proteins—haemoglobin in red blood cells, myosin in muscle, albumin in the liver, and so on. Moreover, with few exceptions all these different cell types contain the same sequence of DNA, which encodes all these different cell proteins, and this DNA is also identical to the DNA in the single celled zygote, from which all these different cells arise during embryonic development. Clearly, therefore, some process of gene regulation must operate to decide which genes within the DNA will be active in producing proteins in each cell type.

Levels of gene regulation

A number of stages exist between the DNA itself and the production of a particular protein (fig 1).¹ Thus the DNA must first be transcribed into a primary RNA transcript, which is subsequently modified at both ends by the addition of a 5' cap and a 3' tail of adenosine residues. Moreover, within this primary transcript, the RNA sequences which actually encode the protein are not present as one continuous block. Rather they are broken up into segments (exons) which are separated by intervening sequences (introns) that do not contain any protein coding information. As these introns interrupt the protein coding region and would prevent the production of an intact protein they must be removed by the process of RNA splicing² before the mature messenger RNA can be transported from the nucleus to the cytoplasm and translated into protein.

Clearly each of these stages is a potential point at which gene expression could be regulated, and there is evidence that several of them are actually used. Thus, for example, the production of many new proteins in the egg immediately after fertilisation and the start of embryonic development depends on the translation into protein of fully spliced, messenger RNAs that pre-existed in the cytoplasm of the unfertilised egg but

whose translation was blocked before fertilisation. This form of gene regulation is known as translational control. Similarly, by splicing the protein coding regions (exons) of a single primary transcript in different combinations two or more different mRNAs encoding different proteins in different tissues can be produced. This process of alternative splicing³ is well illustrated in the single gene that encodes both the calcium modulating hormone, calcitonin, and the

FIG 1—Stages in gene expression which could be regulated

Division of Molecular Pathology, University College and Middlesex School of Medicine, London W1P 6DB
David S Latchman, professor and head of division

BMJ 1992;304:1103-5

potent vasodilator, calcitonin gene related peptide (fig 2). Thus this gene is transcribed into a primary RNA transcript both in the thyroid gland and in the brain, but a different combination of exons are spliced together in each cell type to produce calcitonin mRNA in the thyroid and calcitonin gene related peptide mRNA in the brain.

FIG 2—Alternative splicing of primary transcript of calcitonin/calcitonin gene related peptide gene in brain and thyroid cells. Splicing followed by proteolytic cleavage of precursor protein produced in each tissue yields calcitonin in thyroid and calcitonin gene related peptide in brain

Although there are several cases of gene regulation after the first stage of transcription, a wide variety of evidence indicates that in most cases gene regulation is achieved at the initial stage of transcription, by deciding which genes should be transcribed into the primary RNA transcript.⁴ In these cases, once transcription has occurred, all the other stages in gene expression shown in figure 1 follow and the corresponding protein is produced. Thus the myosin gene is transcribed only in muscle cells, resulting in myosin being produced only in this cell type; the immunoglobulin gene is transcribed only in B lymphocytes, which produce immunoglobulin; and so on. Indeed, even in the case of calcitonin and calcitonin gene related peptide alternative splicing is acting as a supplement to transcriptional control since the calcitonin/calcitonin gene related peptide gene is transcribed only in the thyroid gland and the brain and not in other tissues. Thus the regulation of gene transcription has a critical role in the regulation of gene expression.

Regulation of transcription

For a gene to be transcribed it is necessary for specific protein factors known as transcription factors⁵ to bind to particular DNA binding sites in the regulatory regions of the gene and induce its transcription by the enzyme RNA polymerase. Some of these factors are present in all cell types while others are active only in specific cells or after exposure to a particular stimulus. The combination of particular binding sites in a particular gene determines the transcription factors which bind to it, and in turn the presence or absence of these factors determines in which cell type(s) the gene is transcribed. Thus, for example, the immunoglobulin genes contain a binding site for the octamer binding transcription factor Oct-2 in their regulatory region or promoter, upstream of the start site for transcription. The Oct-2 factor is synthesised only in B lymphocytes and hence binds only to the immunoglobulin gene promoter in B cells, resulting in the transcription of the immunoglobulin genes only in antibody producing B cells. Similarly, genes expressed only in muscle cells such as the creatine kinase gene

contain binding sites for the MyoD transcription factor, which is present only in muscle cells. This case is even more dramatic, however, because the artificial expression of MyoD in non-muscle cells such as fibroblasts is sufficient to convert them into muscle cells, indicating that MyoD activates transcription of all the genes whose protein products are necessary to produce a differentiated muscle cell.⁶

Unlike the case of MyoD, the expression of Oct-2 alone is not sufficient to produce differentiated B cells. This is because other transcription factors that are specifically active in B cells are also involved in producing the expression of genes specific to B cells such as those encoding the immunoglobulins. One such factor is NF κ B, which binds to a DNA sequence in the regulatory region of the immunoglobulin κ light chain gene. Interestingly, unlike Oct-2, the NF κ B protein is present in all cell types. In most cells, however, it is present in an inactive form in which it is complexed with an inhibitory protein, resulting in it being restricted to the cell cytoplasm. In mature B cells, however, NF κ B is released from the inhibitory protein and moves to the nucleus, where it can bind to its DNA target sequence and activate the transcription of the immunoglobulin κ light chain gene.⁷ Interestingly, this activation of NF κ B also occurs when resting T lymphocytes are activated by antigenic stimulation and is the main reason for the improved growth of the human immunodeficiency virus in activated compared with resting T cells since NF κ B can bind to two sites within the HIV promoter and activate viral transcription.

Hence the action of transcription factors on gene expression can be controlled not only by regulating their synthesis but also by regulating their activity (fig 3). The combination of these two processes allows transcription factors to regulate the expression of numerous different genes in different cell types.

Malregulation of gene expression in disease

FAILURE OF TRANSCRIPTION FACTOR FUNCTION

In view of the complex nature of gene regulation it is not surprising that it can go wrong and that a number of human diseases have now been shown to be due to defects in gene regulation. Thus one type of congenital severe combined immunodeficiency is caused by a failure of HLA class II gene transcription, resulting in the absence of these proteins. In turn this failure of transcription is dependent on the lack of a specific transcription factor necessary for the transcription of these genes.⁸ Conversely in haemophilia B the particular transcription factor necessary for transcription of the factor IX gene is present, but it fails to bind to the gene promoter owing to a mutation in the DNA sequence to which it would normally bind, hence resulting in a failure of gene transcription.⁹ Such defects in gene regulation can also affect steps of gene expression other than transcription. Thus the failure to produce one of the two alternatively spliced mRNAs derived from the porphobilinogen deaminase gene is the cause of one form of acute intermittent porphyria.¹⁰

PROTO-ONCOGENES

As well as cases such as these, where a disease is caused by the failure to express a particular gene, malregulation of gene expression can also result in disease if it causes genes to be expressed at the wrong time or in the wrong place. This form of malregulated gene expression is central to the development of certain cancers. Thus it is now clear that most, if not all, human cancers are caused by the mutation or overexpression of certain specific cellular genes known as proto-oncogenes, which results in their conversion into cellular oncogenes capable of causing cancer.¹¹

FIG 3—Transcription factors can activate gene expression in a particular tissue if (top) they are synthesised only in that tissue or (bottom) are present in an active form in that tissue

Although proto-oncogenes encode many different types of cellular proteins involved in growth regulation, such as growth factors or their receptors, several, such as *erbA*, *fos*, *jun*, *myb*, and *myc*, encode cellular transcription factors that are involved in regulating the expression of specific genes. After the conversion of these proto-oncogenes into oncogenes by mutation or overexpression, corresponding alterations occur in the expression of the genes which they regulate, resulting in cancer.

One example of this is provided by the related *Fos* and *Jun* proteins,¹² both of which encode cellular transcription factors. After treatment of cells with growth stimulating factors *Fos* and *Jun* are synthesised and activate the transcription of specific genes whose protein products are necessary for cellular growth. Normally, however, *Fos* and *Jun* are synthesised only transiently in response to exposure to the growth factor, resulting in only a transient activation of gene expression and thereby producing the controlled growth factor regulated proliferation characteristic of normal cells. If for any reason, however, *Fos* and *Jun* are continually synthesised, either owing to a mutation, resulting in their continual overexpression, or to infection with a virus expressing one or other of them, the cell is stimulated to grow continually even in the absence of growth factors. Such continuous uncontrolled growth is characteristic of the cancer cell.

Hence the *fos* and *jun* genes are proto-oncogenes whose products have a critical role in the growth of normal cells but which can be converted into oncogenes capable of transforming cells. Moreover, in contrast to the other diseases I have discussed, in this case malregulation of gene expression and disease is caused not by failure of transcription factor function but rather by failure to regulate correctly the activity of

the factor so that it is active in the wrong place at the wrong time. This indicates that, as with other cellular processes, gene expression is subject to complex regulatory mechanisms, the failure of which can be as disastrous as the failure of the basic process itself.

Conclusion

This review gives a brief overview of the major aspects of gene regulation mechanisms (for further details see elsewhere).⁴ But much remains to be understood. For example, it remains unclear how the expression of specific genes is regulated both spatially and temporally during development so that each cell type arises in the correct place and at the correct time. It is already clear, however, that the correct regulation of gene expression is central to health and correct development and that its malregulation is involved in a number of diseases.

- 1 Nevins JR. The pathway of eukaryotic mRNA transcription. *Ann Rev Biochem* 1983;52:441-6.
- 2 Sharp PA. Splicing of messenger RNA precursors. *Science* 1987;235:766-71.
- 3 Latchman DS. Cell-type specific splicing factors and the regulation of alternative RNA splicing. *The New Biologist* 1990;2:297-303.
- 4 Latchman DS. Gene regulation: a eukaryotic perspective. London: Unwin Hyman, 1990.
- 5 Latchman DS. Eukaryotic transcription factors. *Biochem J* 1990;270:281-9.
- 6 Olson EN. Myo D family: a paradigm for development? *Genes and Development* 1990;4:1454-61.
- 7 Lenardo MJ, Baltimore D. NF-kappa B: a pleiotropic mediator of inducible and tissue-specific gene control. *Cell* 1989;58:227-9.
- 8 Reith W, Satola S, Sanchez CH, Amaldi I, Lisowska-Grospiere B, Griscelli C, et al. Congenital immunodeficiency with a regulatory defect in MHC class II gene expression lacks a specific HLA-DR promoter binding protein RF-X. *Cell* 1988;53:897-906.
- 9 Crossley M, Brownlee GG. Disruption of a C/EBP binding site in the factor IX promoter is associated with haemophilia B. *Nature* 1990;345:444-6.
- 10 Grandchamp B, Picat C, Mignotte V, Wilson JHP, Te Velde K, Sandkuyil L, et al. Tissue-specific splicing mutation in acute intermittent porphyria. *Proc Natl Acad Sci USA* 1989;86:661-4.
- 11 Bishop JM. The molecular genetics of cancer. *Science* 1987;235:305-11.
- 12 Curran T, Franza BR. Fos and Jun: the AP-1 connection. *Cell* 1988;55:315-97.

ANY QUESTIONS

With the benefit of hindsight, was artificial pneumothorax a useful treatment for pulmonary tuberculosis?

As artificial pneumothorax was widely used it was useful, but no one knows how much good it did. There was a consensus among experienced doctors that selected patients with active, but not too active, disease located favourably in the lungs were likely to benefit from well maintained artificial pneumothorax, but the selection criteria varied widely. In a large proportion of patients thought suitable, induction proved impossible because of pleural adhesions. A common view was that if selective collapse of diseased parts could be attained and if this resulted in closure of cavities prognosis was greatly improved. Against this apparent benefit the ill effects of complications, notably pleural effusion, which might be transient but in some cases led to progressive pleural obliteration and thickening and in others became purulent, had to be considered.

The fact that many patients who were doctors, some of them experts in tuberculosis, elected to be treated by artificial pneumothorax indicates that on balance it was thought to be beneficial. But this view was based on uncontrolled observations and long term follow up studies in which the only possible comparative group was composed of patients with disease thought to be of similar extent and activity but who had not been treated by artificial pneumothorax. Since the most frequent reason why patients of this sort did not have artificial pneumothorax was that it had been attempted but found impracticable because of pleural adhesions, this comparison suffered from the defect of an obvious difference in the initial state of the groups compared as well as all the

problems of retrospective studies. But as no prospective controlled study of any of the policies on the use of artificial pneumothorax was conducted no one knew whether accepted views were correct.

There are good reasons for what may seem now to be a surprising omission. The most obvious is that artificial pneumothorax had been established as an accepted therapeutic procedure for many years before controlled trials were introduced into clinical studies. And the practical and ethical difficulties of the sort of trial that would have been required to test its efficacy would have been insuperable. Firstly, it would have been necessary to recruit patients with a defined sort of disease thought likely to benefit. Secondly, they would have had to be randomised into groups, in one of which artificial pneumothorax was to be attempted. Since nearly all doctors thought that appropriately selected patients were likely to benefit they would certainly have rejected any such trial as unethical. Indeed, it is difficult to imagine its being approved by an ethics committee today.

When artificial pneumothorax was uncomplicated and resulted in selective collapse and closure of cavities patients certainly benefited, but I am unsure how far the benefit for these patients was counterbalanced by the ill effects of pleural and other complications in others.

This shows the difficulty of assessing claims that procedures are beneficial but only to selected patients. By comparison, when specific antimycobacterial agents became available it was possible to design and carry out the series of controlled trials which led to generally applicable regimens of treatment, based on evidence open to general inspection. — J G SCADDING, *emeritus professor of medicine, University of London*