Many of these associations with microalbuminuria are even more pronounced in the presence of overt proteinuria,^{14 15 23} which predicts a much greater risk of death from coronary heart disease in diabetic and hypertensive patients and in the general population.²⁴⁻²⁶ In patients with insulin dependent diabetes mellitus overt proteinuria may be present, however, in fewer than half of those with established coronary heart disease.²⁷

Microalbuminuria may also be a non-specific marker for acute illness²⁸ (including myocardial infarction), when it may be operating as an acute phase reactant, and of malignancy, when it probably reflects a microvascular response to tumour related mediators.²⁹ In these circumstances microalbuminuria could have a prognostic value in reflecting the severity of the acute illness or response to treatment.

Microalbuminuria is uncommon without established cardiovascular disease or other cardiovascular risk factors. Screening for microalbuminuria currently seems justified in diabetic and hypertensive patients, and possibly in elderly patients, as it seems to identify those at greatest risk of premature death. Microalbuminuria and the subsequent excess risk of nephropathy and cardiovascular disease may be amenable to correction in diabetic patients. Whether similar gains are possible in elderly patients with microalbuminuria and patients with essential hypertension remains to be seen, but in such cases consideration might be given to more active management of hypertension.

P H WINOCOUR

Lecturer in Medicine
Department of Medicine,
Medical School,
University of Newcastle upon Tyne,
Newcastle upon Tyne NE2 4HH

- 1 Marshall SM, Shearing PA, Alberti KGMM. Evaluation of Micral—test strips for screening for microalbuminuria. *Clin Chem* (in press).
- 2 Marshall SM. Screening for microalbuminuria: which measurement? *Diabetic Med* 1991;8:706-11.
- 3 Winocour PH, Harland J, Millar JP, Laker MF, Alberti KGMM. Microalbuminuria and associated risk factors in the community. *Atherosclerosis* 1992;93:71-81.
- 4 Haffner SM, Stern MP, Gruber KK, Hazuda HP, Mitchell BD, Patterson JK. Microalbuminuria.

- Potential marker for increased cardiovascular risk factors in nondiabetic subjects? *Arteriosclerosis* 1990;10:727-31.
- 5 Allawi J, Rao PV, Gilbert R, Scott G, Jarrett RJ, Keen H, *et al*. Microalbuminuria in non-insulin dependent diabetes: its prevalence in Indian compared with Europid subjects. *BMJ* 1988;296:462-4.
 - 6 Damsgaard EM, Mogensen CE. Microalbuminuria in elderly hyperglycaemic patients and controls. *Diabetic Med* 1986;3:430-35.
 - 7 Gatling W, Knight C, Mullee MA, Hill RD. Microalbuminuria in diabetes: a population study of the prevalence and an assessment of three screening tests. *Diabetic Med* 1988;5:343-7.
 - 8 Mogensen CE. Microalbuminuria predicts clinical proteinuria and early mortality in maturity-onset diabetes. *N Engl J Med* 1984;310:60.
 - 9 Jarrett RJ, Viberti GC, Argyropoulos A, Hill RD, Mahmud U, Murrells TJ. Microalbuminuria predicts mortality in non-insulin-dependent diabetes. *Diabetic Med* 1984;1:17-20.
 - 10 Damsgaard EM, Froland A, Jorgensen OD, Morgensen CE. Microalbuminuria as predictor of increased mortality in elderly people. *BMJ* 1990;300:297-300.
 - 11 Yudkin JS, Forrester RD, Jackson CA. Microalbuminuria as a predictor of vascular disease in non-diabetic subjects. Islington diabetes Survey. *Lancet* 1988;ii:530-3.
 - 12 Jones SL, Close CF, Mattock MB, Jarrett RJ, Keen H, Viberti GC. Plasma lipid and coagulation factor concentrations in insulin dependent diabetics with microalbuminuria. *BMJ* 1989;298:487-90.
 - 13 Parving H-H, Jensen H, Mogensen CE, Evrin PE. Increased urinary albumin excretion rate in benign essential hypertension. *Lancet* 1974;ii:1190-92.
 - 14 Winocour PH, Durrington PN, Bhatnagar D, Ishola M, Mackness M, Arrol S. The influence of early diabetic nephropathy on a very low density lipoprotein (VLDL), intermediate density lipoprotein (IDL), and low density lipoprotein (LDL) composition. *Atherosclerosis* 1991;89:49-57.
 - 15 Winocour PH, Bhatnagar D, Ishola M, Arrol S, Durrington PN. Lipoprotein (a) and microvascular disease in type 1 (insulin dependent) diabetes mellitus. *Diabetic Med* 1991;8:922-7.
 - 16 Reaven GM. The role of insulin resistance in human disease. *Diabetes* 1988;37:1595-607.
 - 17 O'Donnell MJ, Le Guen CA, Lawson N, Gyde OHB, Barnett AH. Platelet behaviour and haemostatic behaviour in type 1 (insulin-dependent) diabetic patients with and without albuminuria. *Diabetic Med* 1991;8:624-8.
 - 18 Schmitz A, Ingerslev J. Haemostatic measure in type 2 diabetic patients with microalbuminuria. *Diabetic Med* 1990;7:521-5.
 - 19 Deckert T, Feldt-Rasmussen B, Borch-Johnsen K, Jensen T, Kofoed-Enevoldsen A. Albuminuria reflects widespread vascular damage. The Steno hypothesis. *Diabetologia* 1989;32:219-26.
 - 20 Parving HH, Jensen HAE, Westrup M. Increased transcapillary escape rate of albumin and IgG in essential hypertension. *Scand J Clin Lab Invest* 1977;37:223-7.
 - 21 Winocour PH, Dhar H, Anderson DC. The relationship between autonomic neuropathy and urinary sodium and albumin excretion in insulin-treated diabetics. *Diabetic Med* 1986;3:436-40.
 - 22 Ewing DJ, Campbell IW, Clarke BF. The natural history of diabetic autonomic neuropathy. *Q J Med* 1980;49:95-108.
 - 23 Winocour PH, Durrington PN, Ishola M, Anderson DC, Cohen H. Influence of proteinuria on vascular disease, blood pressure, and lipoproteins in insulin dependent diabetes mellitus. *BMJ* 1987;294:1648-51.
 - 24 Borch-Johnsen K, Andersen PK, Deckert T. The effect of proteinuria on relative mortality in type 1 (insulin dependent) diabetes mellitus. *Diabetologia* 1985;28:590-6.
 - 25 Nelson RG, Pettitt DJ, Carraher MJ, Baird MJ, Knowler WC. The effect of proteinuria on mortality in NIDDM. *Diabetes* 1988;37:1499-504.
 - 26 Bulpitt CJ, Beilohn LJ, Clifton P, Coles EC, Dollery CT, Gear JSS, *et al*. Risk factors for death in treated hypertensive patients. Reports from the DHSS hypertension care computing project. *Lancet* 1979;ii:134-7.
 - 27 Orchard TJ, Dorman JS, Maser RE, Becker DJ, Drash AL, Ellis D. Prevalence of complications in IDDM by sex and duration. Pittsburgh epidemiology of diabetes complications study. II. *Diabetes* 1990;39:1116-24.
 - 28 Gosling P, Shearman CP. Increased levels of urinary proteins: markers of vascular permeability? *Ann Clin Biochem* 1988;25 (suppl): 150-1.
 - 29 Sawyer N, Wadsworth J, Mijnen M, Gabriel R. Prevalence, concentration, and prognostic importance of proteinuria in patients with malignancies. *BMJ* 1988;296:1295-8.

Patterns of hospital medical staffing

Future implications

The muted response to the publication of the full text of Robin Dowie's report *Patterns of Hospital Medical Staffing*¹ is surprising given the publicity that surrounded the release of her interim report.² Several explanations are possible. The sheer scope and volume of information presented—eight specialty reports and the overview—may have discouraged some commentators; others may have found some of the conclusions unpalatable.

Despite the delay between fieldwork and publication the findings of the research are still timely, as Dowie says in her overview. As the most detailed examination yet of the distribution of medical workload across grades and specialties the report should be required reading for all members of regional task forces. Overall, the picture that emerges is one of poor management of scarce personnel: junior doctors having to perform inappropriate tasks; consultants in surgical specialties, especially trauma and orthopaedic surgery, denied adequate theatre time; lack of educational opportunities at all levels; and maldistribution of staff within and between specialties as well as geographically.

Each specialty report ends with a series of specific recommendations. The most novel of these is for the designation of

specific vocational trainee posts for general practitioners within the hospital service, to be funded centrally and separately from the posts for specialist trainees. (Dowie estimates the initial cost at about £50m.) This would release funds to employ more career grade staff. This seems attractive given that the outlay would be relatively small (for the NHS) and the returns could be substantial. From the perspective of junior doctors Dowie's dismissal of the possibility of appreciable consultant expansion as "unrealistic" given "the present economic climate" is therefore disappointing, especially as much of the overview contains cogent reasons why such expansion is needed. Virtually all the specialty reports include recommendations to increase the staff grade, which she feels should be more flexibly deployed.

Much of the argument for such a course of action is derived from an interpretation of the staffing projections contained in *Achieving a Balance—Plan for Action*.³ This anticipated a 15% decline in numbers of registrars between 1987 and 1998 coupled with a modest increase in the numbers of senior house officers, thus bringing about an absolute fall in the number of intermediate grade trainees. The projected increase in numbers has already been outstripped⁴ and the Joint Planning

Advisory Committee's latest report shows that more career registrar posts have already been approved⁵ than were envisaged in the "do nothing" projections in *Achieving a Balance*.

Preferentially increasing the staff grade while delaying or curtailing the expansion of consultant numbers will appeal to hard pressed managers seeking to provide adequate services with inadequate budgets. Indeed there may still be, as Dowie contends, "a substantial reservoir" of people qualified and willing to undertake such posts. Most junior doctors, however, undoubtedly still want to train as consultants.

Many of the report's recommendations should receive the wholehearted support of all doctors. They include the setting up of more preregistration posts in general practice, further research into the workloads of individual specialties, skill substitution, the costing of the use of hospital doctors' time, and better statistical support for planning medical manpower. We should not be content, however, to support a system in which many career grade hospital staff never achieve the status of fully trained specialists.

What is needed is a fundamental reappraisal of where

services are delivered and by whom. This is particularly pertinent at present, given that we face a growing elderly population, a fall in the number of applicants for medical school, and the possibility of a pan-European shortfall of medical staff within the next 15 years.⁶ Perhaps the time has come to blow the dust off the Short report⁷ at last.

ANDREW CARNEY

Chairman, North West Thames Junior Doctors Committee,
London WC1H 9SA

- 1 Dowie R. *Patterns of hospital medical staffing. Overview and specialty reports*. London: British Postgraduate Medical Federation, 1991.
- 2 Dowie R. *Patterns of hospital medical staffing. Interim report: junior doctors' hours*. London: British Postgraduate Medical Federation, 1990.
- 3 UK Health Departments, Joint Consultants Committee, and Chairmen of Regional Health Authorities. *Hospital medical staffing: achieving a balance—plan for action*. London, DHSS, 1987.
- 4 Department of Health. *Hospital medical staff—England and Wales*. London: DoH, Statistics and Management Information Division, 1991: national table 2 (30/09/90).
- 5 Joint Planning Advisory Committee. *Report for the years 1990/1*. London: DoH, 1991.
- 6 Saugmann P. Medical manpower in West Europe: towards a balance between supply and demand by the Year 2000. In: *Medical manpower in Europe. From surplus to deficit?* Copenhagen: Permanent Working Group of European Junior Hospital Doctors, 1991.
- 7 House of Commons Select Committee on Social Services. *Fourth report from the social services committee, session 1980/1*. London: HMSO, 1981. (Medical education, HC 31; Short report.)

Medical treatment for benign prostatic hyperplasia

Surgery still gives the best results

Transurethral prostatectomy remains the treatment of choice for symptomatic benign prostatic hyperplasia unless the prostate is large enough to warrant the retropubic approach. Recently attention has turned to alternative treatments for the condition: now we are in the middle of a surge of interest in possible medical treatments.

Interpreting the results of drug treatment requires an understanding of the very variable course of untreated prostatic obstruction: short term improvements may reflect the natural course of the disease rather than the effects of treatment. Patients should be carefully assessed before treatment is begun—not only to allow an accurate assessment of the drugs' subsequent effects but also to exclude prostatic malignancy, complications of prostatic obstruction, and other coexisting diseases.

The exclusion of prostatic cancer is currently provoking debate among urologists. Without a tissue diagnosis (automatically available with prostatectomy) focal carcinomas which would otherwise be diagnosed, will be missed in up to one in five patients. Most of these tumours will probably remain occult and not influence patients' survival. Rectal examination will miss most of them; measuring the serum concentration of prostate specific antigen before starting drug treatment¹ and performing transrectal ultrasonography of the prostate when the concentration of antigen is raised will provide the best diagnostic accuracy.

The first attempt at medical treatment was chemical castration with hormones, but this never achieved widespread acceptance because of its inevitable effects on sexual function. Interest in hormonal treatment has revived with the advent of a new class of drug that inhibits the conversion of testosterone to the more active dihydrotestosterone within the prostate by inhibiting the enzyme 5 α -reductase (this avoids any effect on circulating androgens and hence the unwanted systemic effects associated with castration). The results of clinical trials are awaited.

Currently the main medical treatment for benign prostatic hyperplasia is α adrenergic blockade. Phenoxybenzamine,

which blocks α_1 and α_2 adrenoceptors, has improved urinary flow and clinical symptoms in most clinical studies.^{2,3} Some 30% of patients experience side effects,² which have been attributed to the blockage of presynaptic α_2 adrenoceptors. This is believed to interfere with the normal negative feedback control of the release of noradrenaline at the presynaptic adrenergic nerve terminal, thus resulting in a high circulating concentration of noradrenaline. Phenoxybenzamine's side effects and evidence of its mutagenicity in bacterial and mouse cell cultures⁴ have limited its use in the treatment of benign prostatic obstruction.

The experimental work of Caine and associates⁵ and the incidental clinical observation reported in the *BMJ* in 1978 that urinary incontinence developed in women during anti-hypertensive treatment with prazosin⁶ contributed to the recognition of the potential role of selective α_1 adrenoceptor blockade as treatment for diseases of the lower urinary tract. Subsequent studies by several groups of workers have shown the functional and ultrastructural pre-eminence of α_1 receptors over α_2 receptors within the stromal compartment of the prostate,⁷ thereby providing a scientific basis for the use of specific α_1 blockade in non-surgical management of benign prostatic hyperplasia. Most studies of the specific α_1 antagonists have been conducted over relatively short periods of two to four weeks⁸⁻¹¹ and have shown objective improvements in mean and maximum urinary flow rates of 14-96% with subjective improvements in patients' symptom scores.

Longer term studies, which are of greater relevance to clinical practice, have investigated terazosin,¹² indoramin,¹³ prazosin,¹⁴ doxazosin,¹⁵ and alfuzosin.¹⁶ Only the study of alfuzosin reported by Ramsay *et al* failed to show any improvement in the outflow obstruction.¹⁷ All the other studies showed that drug treatment significantly improved urinary flow rates, although the improvement in flow rates after at least three months' treatment was less than that previously reported in shorter term studies. For example, in a study of the longer term use of terazosin, Lepor and Knapp-Maloney reported an increase in maximum flow rate of 4.7