4XN (0222 747747); Western General Hospital, Edinburgh EA4 2HU (031 3327917); Duncan Guthrie Institute, Glasgow G3 8SJ (041 339 8888); Kennedy Galton Centre, Northwick Park Hospital, Middlesex HA1 3UJ (081 422 8577); Clarendon Wing, General Hospital, Leeds LS2 9NS (0532 432799); Leicester Royal Infirmary, Leicester LE1 7LX (0533 541414); Royal Liverpool Hospital, Liverpool L69 3BX (051 709 0141); Institute of Child Health, London WC1N 1EA (071 242 7989); Seventh and Eighth Floors, Guy's Tower, Guy's Hospital, London SE1 9RT (071 955 4648); St George's Hospital Medical School, London SW17 0RE (071 672 9944); St Mary's Hospital, Manchester M13 0JH (061 276 6322); 19/20 Claremont Place, Newcastle upon Tyne NE2 4AA (0912 226000); City Hospital, Nottingham NG5 1PB (0602 691169); Churchill Hospital, Oxford OX3 7LE (0865 62834); 117 Manchester Road, Sheffield S10 5DN (0742 27511); Princess Anne Hospital, Southampton SO9 4HA (0703 796166).

- Gusella JF, Wexler NS, Conneally PM, Naylor SL, Anderson MA, Tanzi RE *et al.* A polymorphic DNA marker genetically linked to Huntington's disease. *Nature* 1983;306:234-8.
- Hayden MR, Robbins C, Allard D, Haines J, Fox S, Wasmuth J, *et al.* Improved predictive testing for Huntington's disease by using three linked DNA markers. *Am J Hum Genet* 1988;43:689-94.
- Harper PS, ed. Predictive tests in Huntington's disease. In: *Huntington's disease. Major problems in neurology*. Vol 22. London: W B Saunders, 1991:373-413.
- Craufurd D, Harris R. Ethics of predictive testing for Huntington's chorea: the need for more information. *BMJ* 1986;293:249-51.

- Morris M, Tyler A, Harper PS. Adoption and genetic prediction for Huntington's disease. *Lancet* 1988;ii:1069-70.
- World Federation of Neurology: Research Committee Research Group on Huntington's Chorea. Ethical issues policy statement on Huntington's disease molecular genetics predictive test. *J Neurol Sci* 1989;94:327-32.
- Harper PS, Newcombe RG. Age at onset and life table risks in genetic counselling for Huntington's disease. *J Med Genet* 1992;29:239-43.
- Sandkuyl L, Ott J. Determining informativity of marker typing for genetic counselling in a pedigree. *Hum Genet* 1989;82:159-62.
- Tyler A, Harper PS. Attitudes of subjects at risk and their relatives towards genetic counselling in Huntington's chorea. *J Med Genet* 1983;20:179-88.
- Mastromauro C, Myers RH, Berkman B. Attitude towards presymptomatic testing in Huntington's disease. *Am J Hum Genet* 1987;26:271-82.
- Harper PS, ed. The epidemiology of Huntington's disease. In: *Huntington's disease. Major problems in neurology*. Vol 22. London: W B Saunders, 1991:251-80.
- Craufurd D, Dodge A, Kerzin-Storror L, Harris R. Uptake of presymptomatic predictive testing for Huntington's disease. *Lancet* 1989;ii:603-5.
- Meissen GA, Myer RH, Mastromauro C, Koroshetz WJ, Klinger KW, Farrer LA, *et al.* Predictive testing for Huntington's disease with use of a linked DNA marker. *N Engl J Med* 1988;318:535-42.
- Brandt J, Quaid KA, Folstein SE, Garber P, Maestri NE, Abbott MH, *et al.* Presymptomatic diagnosis of delayed-onset disease with linked DNA markers. The experience in Huntington's disease. *JAMA* 1989;261:3108-14.
- Bloch M, Fahy M, Fox S, Hayden MR. Predictive testing for Huntington's disease; demographic characteristics, lifestyle patterns, attitudes and psychological assessments of the first fifty-one test candidates. *Am J Med Genet* 1989;32:217-24.
- Tyler A, Morris M, Lazarou L, Meredith L, Myring J, Harper PS, *et al.* Presymptomatic testing for Huntington's disease in Wales 1987. 1990. *Br J Psychiatry* (in press).
- Lam RW, Bloch M, Jones BD, Marcus AM, Fox S, Amman W, *et al.* Psychiatric morbidity associated with early clinical diagnosis of Huntington disease in a predictive testing programme. *J Clin Psychiatry* 1988;49:444-7.

(Accepted 23 March 1992)

Electrocardiographic abnormalities and associated factors in Chinese living in Beijing and in Mauritius

Nan Li, Jaakko Tuomilehto, Gary Dowse, K George Alberti, Paul Zimmet, Zhang Min, Pierrot Chitson, Hassam Gareebou, You Chonghua, Djamil Fareed for the Mauritius Non-Communicable Disease Study Group

Beijing Heart Lung and Blood Vessel Medical Centre, Beijing, China
Nan Li, research fellow
Zhang Min, systems analyst
You Chonghua, director

Department of Epidemiology, National Public Health Institute, 00510 Helsinki, Finland
Jaakko Tuomilehto, professor

International Diabetes Institute, Caulfield, Australia
Gary Dowse, epidemiologist
Paul Zimmet, professor

Medical School, Newcastle upon Tyne
K George Alberti, professor

Noncommunicable Diseases Office, Ministry of Health, Port Louis, Mauritius
Pierrot Chitson, medical officer
Hassam Gareebou, medical officer

World Health Organisation, Port Louis, Mauritius
Sir Djamil Fareed, WHO representative

Correspondence to: Professor J Tuomilehto, Department of Epidemiology, National Public Health Institute, Elimäenkatu 25A 6th Floor, 00510 Helsinki, Finland.

BMJ 1992;304:1596-601

Abstract

Objective—To compare the prevalence of electrocardiographic abnormalities and to evaluate the association between these abnormalities and the levels of coronary heart disease among Chinese living in different environments.

Design—Cross sectional surveys.

Setting—Beijing, China, and the island of Mauritius.

Subjects—Random samples of people aged 35-64 years in Beijing (621 men, 642 women) in 1984 and in Mauritius among Chinese (137 men, 130 women) and non-Chinese (1265 men, 1432 women) in 1987.

Main outcome measures—Prevalence of electrocardiographic abnormalities suggesting coronary heart disease and of associated risk factors.

Results—Prevalence of electrocardiographic abnormalities suggesting coronary heart disease was significantly lower in Beijing (4.0%) than in Mauritian Chinese (24.3%) and Mauritian non-Chinese (24.5%). Mean serum concentrations of total and non-high density lipoprotein cholesterol were lower in Beijing Chinese than in Mauritian Chinese, but smoking and hypertension were slightly more prevalent. Overall, men with electrocardiographic abnormalities had higher risk factor levels than those with a normal electrocardiogram regardless of ethnic origin.

Conclusions—The prevalence of coronary heart disease and associated risk factors was different among Chinese living in two different environments: in Beijing in the People's Republic of China and in Mauritius. Chinese, who traditionally have a very low frequency of coronary heart disease, are by no means protected against coronary heart disease and other non-communicable diseases. Therefore, primary prevention of coronary heart disease is a

major challenge for preventive medicine in China, as well as in many other developing countries.

Introduction

International comparisons have shown that the occurrence of coronary heart disease varies greatly from one country to another. In China the incidence of and mortality from coronary heart disease are low, as shown by, for example, the World Health Organisation monitoring trends and determinants in cardiovascular disease (MONICA) project.¹ The highest age standardised coronary heart disease mortality in men aged 35-64 years was found in the Siberian MONICA study (401/100 000) and the lowest in the Sino-MONICA-Beijing study (40/100 000). In women the highest coronary heart disease mortality was found in the Glasgow MONICA study (132/100 000) whereas in Beijing it was only 28/100 000.¹ These 10-fold differences in men and 4.7-fold differences in women may be explained by different levels of coronary heart disease risk factors,² but it has also been proposed that these risk factors may have different impact on the development of coronary heart disease in different geographic areas and in genetically different populations.

There have been many studies comparing coronary heart disease risk factor levels in different countries.²⁻⁶ Genetics, ethnic background, lifestyles, and study methods have often differed between the areas, hence the possibilities for valid conclusions from many earlier comparisons are limited. The Ni-Hon-San (Nippon-Honolulu-San Francisco) Japanese migrant study initiated in the mid-1960s overcame many methodological shortcomings and showed a clear gradient in coronary heart disease mortality among Japanese men

increasing from Japan to Hawaii and further to California, despite the subjects studied having the same genetic background.⁷⁻¹¹

The epidemiological evidence suggests that the occurrence of coronary heart disease is largely determined by the environment.^{3,8,11} The aim of this study was to compare the prevalence of electrocardiographic abnormalities and to assess the association between these abnormalities and the levels of major coronary heart disease risk factors among Chinese living in two different environments: in Beijing, China, and in Mauritius. Migrant studies provide one of the best means of generating hypotheses concerning the relation of environmental factors and disease. The island country of Mauritius, which has an entirely immigrant population, is an ideal place for migrant studies. The Chinese form one of three main ethnic groups in Mauritius.

Methods

STUDY POPULATION

The Sino-MONICA-Beijing study covers a population of approximately 700 000, about two thirds urban and one third rural, scattered in six urban and suburban districts and one rural county of Beijing. This distribution approximates that of the whole city. The first Sino-MONICA-Beijing risk factor survey was carried out during August and September of 1984 and August-September 1985 in Beijing.¹² Random sampling was performed in two steps. First, 40 out of 222 urban residence committees and 17 out of 125 rural villages were chosen. At the second step, all individuals in the residence committees or villages were randomised again and enumerated by name, sex, and age. A total of 1682 individuals aged 25-64 years were surveyed, with a response rate of 88%.

The subtropical island of Mauritius is situated in the south western Indian Ocean. The total population is about 1 million, an estimated 70% of whom originated from the Indian subcontinent; 2% are Chinese and 28% are Creole (mixed African, Indian, and European ancestry). The Mauritius non-communicable risk factor survey was completed during April and May 1987.¹³ The target population for the survey was Mauritian adults aged 25-74. A two stage cluster sampling scheme was used to select 10 areas from defined census divisions: six among them were rural and four urban, which approximates the actual rural to urban ratio of the total population. All adults aged 25-74 living in the selected areas were included in the sample. In addition, a specific area with high density of Chinese residents (Chinatown) was chosen to bolster the small number of Chinese subjects in the random sample. A total of 5074 individuals were surveyed, with an overall response rate of 86%: among the 409 Chinese the rate was 78%. Chinese ethnicity was based on looking Chinese, having a Chinese name, and the subjects identifying themselves as Chinese. Inter-marriages between Chinese and other ethnic groups in Mauritius are known to be rare.

Subjects 35-64 years of age for whom electrocardiographic data were available from these two populations were chosen for the present comparative analysis.

BLOOD PRESSURE MEASUREMENT

The same method was used to measure blood pressure in the Sino-MONICA-Beijing and Mauritius studies. Blood pressure was measured in the right arm with a standard mercury sphygmomanometer with a cuff width of 12 cm and length of 42 cm. Measurements were performed after five minutes' rest with the subjects in the sitting position. A minimum number of observers were trained and tested before starting the survey. Systolic and diastolic pressures were recorded

on the basis of the first the fifth phase of the Korotkoff sounds and were expressed to the nearest 2 mm Hg. Blood pressure was measured twice and calculated as the mean of the two measurements.

WEIGHT AND HEIGHT

Height and weight were recorded with the subjects wearing light clothing and without shoes. Accurate balance scales were used and height was recorded to the nearest cm and weight to the nearest 200 g (in Beijing) and to the nearest 100 g (in Mauritius). Body mass index was calculated as weight divided by height squared (kg/m^2).

BLOOD SAMPLING AND BLOOD LIPID DETERMINATION

Blood samples in the Sino-MONICA-Beijing survey were taken after an overnight fast of 12-16 hours. Serum was stored on ice and sent to the laboratory of Beijing Heart, Lung, and Blood Vessel Centre within four hours. Total cholesterol was measured using a cholesterol oxidase enzymatic method. The phosphotungstate-magnesium precipitation method was used to isolate high density lipoprotein cholesterol from low density lipoprotein cholesterol and very low density lipoprotein cholesterol. Regular quality control measures were done by internal and external standardisation with the World Health Organisation MONICA project quality control centre for lipid standardisation in Prague, Czechoslovakia.

Strict internal and external quality control measures also were performed to ensure validity in the Mauritius survey. Plasma total cholesterol and high density lipoprotein cholesterol concentrations were determined from fasting blood specimens by using manual colorimetric methods at the Victoria Hospital, Mauritius. Plasma was separated at the survey site within two hours and transported to the Victoria Hospital in the afternoon for analysis. An aliquot of every tenth specimen was stored frozen and these specimens were measured again by autoanalyser in Newcastle upon Tyne, England, using an enzymatic method. The Newcastle laboratory is part of two national quality assurance programmes in the United Kingdom. Results indicated that total cholesterol was slightly overestimated in the local laboratory in Mauritius. Hence all values have been adjusted according to the regression formula:

$$\text{total cholesterol (mmol/l)} = \frac{\text{original value} - 1.133}{0.81879}$$

ELECTROCARDIOGRAMS

Standard supine 12 lead electrocardiograms were recorded with a paper speed of 25 mm/s in both populations. All electrocardiographic equipment fulfilled the recommendations of the American Heart Association's technical specifications. Electrocardiographic recordings were carried out by technicians who had been specially trained in subject preparation, electrode placement, and other aspects. All electrocardiograms from Beijing were read and classified according to the Minnesota code in duplicate in Beijing, where three observers have been trained by the WHO MONICA project quality control centre for electrocardiogram coding. Discrepancies were adjusted by an experienced supervisor (NL), who also reviewed most of the electrocardiograms classified as normal. Items from normal electrocardiograms, such as heart rate, QRS axis, maximum R wave and S wave, and V5 lead T wave height, were also coded. Electrocardiograms from Mauritius were coded in duplicate in Helsinki and Prague. Electrocardiograms from either Beijing or Mauritius suggesting any ischaemic abnormality were checked by one investigator (NL) to assure that no obvious biases were included in the comparison of abnormalities. An additional recording of 152

TABLE 1—Prevalence of coronary heart disease in men and women by ethnic group according to Minnesota coding criteria of electrocardiograms. Values are numbers (percentages)

Code category	Men				Women			
	Beijing (n=621)	Mauritius			Beijing (n=642)	Mauritius		
		Chinese (n=137)	Non-Chinese (n=1265)			Chinese (n=130)	Non-Chinese (n=1432)	
Probable coronary heart disease*		1 (0.2)	3 (2.1)	26 (2.0)	1 (0.2)		9 (0.6)	
1.1 Q and QS	1 (0.2)	3 (2.2)	17 (1.3)			6 (0.4)		
1.2 Q and QS			9 (0.7)		1 (0.2)	3 (0.2)		
Possible coronary heart disease*		17 (2.7)	24 (17.5)	175 (13.8)	31 (4.8)	38 (29.2)	452 (31.6)	
1.3 Q and QS	4 (0.7)	5 (3.6)	22 (1.7)		2 (0.3)	37 (2.6)		
4.1-4.4 ST-T depression	8 (1.3)	7 (5.1)	63 (5.0)		17 (2.6)	211 (14.7)		
5.1-5.3 T wave inversion	12 (1.9)	21 (15.3)	155 (12.3)		29 (4.5)	422 (29.5)		
7.1 Left bundle branch block			4 (0.3)			4 (0.3)		
Total		18 (2.9)	27 (19.7)	201 (15.9)	32 (5.0)	38 (29.2)	461 (32.2)	
	Cases %	Cases %	Cases %	Cases %	Cases %	Cases %	Cases %	
Probable coronary heart disease	1 0.2	2 1.4	12 1.0	1 0.2		4 0.3		
Possible coronary heart disease	17 2.7	24 17.5	175 13.8	31 4.8	38 29.2	452 31.6		
Total probable and possible coronary heart disease	18 2.9	24 17.5	187 14.8	32 5.0	38 29.2	456 31.8		

The incidences are more than shown in the row "cases" because one person can have two or more different electrocardiographic abnormalities.

* Whitehall criteria (reference 14).

TABLE II—Prevalence of abnormal electrocardiograms suggesting coronary heart disease (probable and possible) by population group, age, and gender

	Men	Women	All
	No (%; 95% confidence interval)	No (%; 95% confidence interval)	No (%; 95% confidence interval)
Beijing Chinese:			
35-54 years	5 (1.2; 0.2 to 2.2)	16 (3.8; 2.0 to 5.6)	21 (2.5; 1.4 to 3.6)
55-64 years	13 (6.3; 3.0 to 9.6)	16 (7.3; 3.8 to 10.8)	29 (6.9; 4.5 to 9.3)
Mauritian Chinese:			
35-54 years	15 (16.3; 8.8 to 23.8)	23 (23.0; 14.8 to 31.2)	38 (19.8; 14.2 to 25.4)
55-64 years	9 (20.0; 8.3 to 31.7)	15 (50.0; 32.0 to 67.9)	24 (32.0; 21.4 to 42.6)
Mauritian non-Chinese:			
35-54 years	114 (11.8; 9.8 to 13.8)	309 (28.8; 26.1 to 31.5)	423 (20.8; 19.0 to 22.6)
55-64 years	73 (24.4; 19.5 to 29.2)	147 (40.8; 35.8 to 45.9)	220 (33.4; 29.8 to 37.0)

normal electrocardiograms from Beijing was done by an independent coder. The same investigator who had checked most of the electrocardiograms (both normal and abnormal) from Beijing also recorded a sample of 200 normal electrocardiograms from Mauritius. These final checkings confirmed that in both countries normal electrocardiograms were correctly coded as normal.

In the final classification of the observed electrocardiographic changes we used the commonly agreed Whitehall criteria¹⁴ where Minnesota codes 1.1, 1.2 (large Q wave) suggest probable coronary heart disease and Minnesota codes 1.3 (small Q wave), 4.1-4.4 (ST segment depression), 5.1-5.3 (T wave inversion), and 7.1 (left bundle branch block) suggest possible coronary heart disease.

STATISTICAL METHODS

Statistical analyses were performed on the National Public Health Institute's VAX computer using SAS (Statistical Analysis Systems) software. Mean values of continuous variables were adjusted for age by analysis of covariance. Significance across groups was assessed by the F test. Age standardised proportions were calculated by the direct method using the "world standard population" as the standard. Significance of differences in proportions across groups was assessed by conventional χ^2 test, and 95% confidence intervals for the prevalence of electrocardiographic abnormalities were computed assuming the Poisson distribution.

The independent effect and relative importance of several putative variables and their interactions on the occurrence of electrocardiographic abnormalities suggesting coronary heart disease among the Chinese were assessed using the multiple logistic regression model. In the final model, in addition to the population, we included age, non-high density lipoprotein cholesterol concentration, and mean arterial pressure, which in univariate analyses were associated with electrocardiographic abnormalities. To assess whether the observed

difference in the association of these three risk factors with electrocardiographic abnormalities was different between the Chinese living in China and in Mauritius, we included interaction terms between centre and other variables in the multiple logistic regression model.

Results

The prevalence of electrocardiographic abnormalities suggesting coronary heart disease was low in all study populations (table I). Since the prevalence of possible coronary heart disease was also low in Beijing, the overall prevalence of positive electrocardiographic findings (probable and possible coronary heart disease) was significantly lower in Beijing (4.0%) than in Mauritian Chinese (24.3%) and in Mauritian non-Chinese (24.5%). The prevalence of possible coronary heart disease was higher in women than in men in each population because ST-T changes (such as 4.2, 4.3, 5.2, 5.3) were more common in women than in men. The prevalence of abnormal electrocardiogram was significantly lower in the younger (35 to 54 years) than older (55 to 64 years) subjects among the three populations (table II).

In Beijing, men with a positive electrocardiogram had significantly higher serum total cholesterol and non-high density lipoprotein cholesterol concentrations, higher systolic and diastolic blood pressures, and a higher prevalence of hypertension than men with a normal electrocardiogram (table III). Only systolic blood pressure was significantly higher in women with a positive electrocardiogram compared with those with a normal electrocardiogram. In Mauritian Chinese, non-high density lipoprotein cholesterol concentration, body mass index, systolic blood pressure, and the prevalence of hypertension were significantly greater in men with electrocardiographic abnormalities than in those without. In women, only the prevalence of hypertension was higher in subjects with a positive electrocardiogram. In the non-Chinese population of Mauritius both men and women with a positive electrocardiogram had significantly increased systolic and diastolic blood pressures and prevalence of hypertension.

In Mauritius, about 40% of the Chinese subjects with an abnormal electrocardiogram had a high blood pressure (table IV). Because of relatively small numbers and similar trends in both sexes data for men and women were pooled. The overall association between electrocardiographic abnormalities and hypertension was similar in younger (35 to 54 year old) and older (55 to 64 year old) subjects. In both populations the increase in the prevalence of electrocardiographic

TABLE III—Selected variables by electrocardiogram (ECG) abnormalities suggesting coronary heart disease (Whitehall criteria¹⁴)

Characteristic	Chinese in Beijing			Chinese in Mauritius			Mauritian non-Chinese		
	ECG normal	ECG abnormal	p Value	ECG normal	ECG abnormal	p Value	ECG normal	ECG abnormal	p Value
<i>Men</i>									
Cholesterol (mmol/l)									
High density lipoprotein	1.2	1.2	0.705	1.1	1.1	0.518	1.3	1.3	0.735
Total	4.2	4.8	0.004	5.4	5.6	0.841	5.8	5.7	0.895
Non-high density lipoprotein	3.1	3.6	0.046	4.3	4.5	0.042	4.5	4.4	0.957
Body mass index (kg/m ²)	23.4	24.9	0.275	24.1	25.9	0.033	23.1	23.0	0.253
Blood pressure (mm Hg):									
Systolic	130	151	0.015	129	136	0.033	129	135	<0.001
Diastolic	85	93	0.002	82	84	0.579	81	85	<0.001
Smoking (%)	37.0	50.0	0.057	23.0	12.5	0.641	42.6	44.4	0.747
Hypertension (%)	24.2	55.6	0.004	18.6	45.8	0.004	16.1	40.1	<0.001
<i>Women</i>									
Cholesterol (mmol/l)									
High density lipoprotein	1.2	1.3	0.823	1.4	1.4	0.414	1.3	1.3	0.154
Total	4.3	4.3	0.165	5.4	5.0	0.400	5.6	5.7	0.091
Non-high density lipoprotein	3.1	3.0	0.177	4.1	3.5	0.549	4.3	4.5	0.062
Body mass index (kg/m ²)	24.5	24.5	0.875	24.1	23.6	0.654	24.9	25.4	0.042
Blood pressure (mm Hg):									
Systolic	128	137	0.003	131	133	0.128	126	134	<0.001
Diastolic	81	83	0.109	78	78	0.616	76	81	<0.001
Smoking (%)	5.9	0	0.805	0	0	1.0	2.4	2.6	0.437
Hypertension (%)	21.8	34.4	0.053	10.9	36.8	0.008	10.9	28.3	<0.001

TABLE IV—Prevalence (%) of electrocardiogram (ECG) abnormalities by hypertension status and age group

Age group	Chinese in Beijing		Chinese in Mauritius		All	
	Blood pressure normal	Blood pressure abnormal*	Blood pressure normal	Blood pressure abnormal*	Blood pressure normal	Blood pressure abnormal*
35-54 years:						
ECG normal	82.5	17.5	87.7	12.3	83.4	16.7
ECG abnormal	76.2	23.8	65.8	34.2	69.5	30.5
p Value (χ^2 test)		0.451		<0.001		0.003
Odds ratio† (95% confidence interval)						2.57 (1.36 to 4.88)
55-64 years:						
ECG normal	65.5	34.5	76.5	23.5	66.7	33.3
ECG abnormal	44.8	55.2	50.0	50.0	47.2	52.8
p Value (χ^2 test)		0.025		0.022		0.002
Odds ratio† (95% confidence interval)						2.62 (1.42 to 4.84)
35-64 years:						
ECG normal	77.0	23.0	84.9	15.1	78.1	21.9
ECG abnormal	58.0	42.0	59.7	40.3	58.9	41.1
p Value (χ^2 test)		0.002		<0.001		<0.001
Odds ratio† (95% confidence interval)						2.97 (1.94 to 4.55)

*Systolic ≥ 160 mm Hg or diastolic ≥ 95 mm Hg or patient treated for hypertension.

†Odds ratio computed controlling for the study population.

TABLE V—Multiple logistic regression of ischaemic electrocardiographic abnormalities suggesting coronary heart disease on various risk factors in Chinese subjects living in Beijing or Mauritius

Variable	Men (n=758)			Women (n=772)		
	Parameter estimate	Standardised error	p Value	Parameter estimate	Standardised error	p Value
Centre (1)*	4.81	1.64	0.003	0.59	1.20	0.623
Non-high density lipoprotein cholesterol	0.30	0.15	0.034	-0.22	0.14	0.097
Mean arterial pressure†	0.03	0.01	0.024	0.02	0.01	0.048
Age	0.06	0.03	0.023	0.07	0.02	<0.001
Mean arterial pressure* Centre (0)	-0.01	0.01	0.643	-0.002	0.01	0.861
Non-high density lipoprotein cholesterol* Centre (0)	-0.23	0.15	0.076	-0.12	0.14	0.195
Age* Centre (0)	-0.05	0.03	0.125	0.02	0.02	0.374

*0=Beijing; 1=Mauritius.

†Computed according to the formula; 1/3 systolic + 2/3 diastolic pressure.

abnormalities with age was parallel with the increase in the prevalence of hypertension.

The multiple logistic regression analysis confirmed that the higher prevalence of electrocardiographic abnormalities was not fully explained by the differences in serum non-high density lipoprotein cholesterol concentration and blood pressure (table V). The interaction between the study area (Beijing v Mauritius) and the three risk factors was non-significant confirming that their effect was similar in both Chinese populations and both in men and women.

Discussion

The present study provides a unique opportunity to assess the risk of coronary heart disease in the Chinese, who in their homeland are known for their low

incidence of and mortality from coronary heart disease.^{1,15} Although increasing evidence is accumulating on the importance of genetic susceptibility to the risk of coronary heart disease,¹⁶⁻¹⁹ genetic markers that might contribute to the lower coronary heart disease risk in the Chinese have not been identified. People of Mongoloid origin, including the Chinese, are particularly susceptible to hypertension, non-insulin dependent diabetes mellitus, and stroke if exposed to environmental risk factors.^{1,13,19-22} Therefore, the apparent lower susceptibility to coronary heart disease in China remains obscure and is probably related to lifestyles and culture.

The main finding of our study was that the prevalence of coronary heart disease in Chinese living in two different environments was different, being three to four times higher in migrant Chinese living in Mauritius than in those living in Beijing, and similar to the prevalence in Asian Indians and Creoles in Mauritius. Studies in migrants are important in testing hypotheses of the role of environmental factors in the aetiology of coronary heart disease.^{7-10,23,24} Until now there have been no comparative studies of the occurrence of coronary heart disease among the Chinese in mainland China and abroad. Our data are thus particularly valuable in that they show the increased susceptibility of the Chinese, given an appropriate environment.

BLOOD PRESSURE, GLUCOSE INTOLERANCE, AND PLASMA LIPIDS

Our earlier results from Mauritius showed that the Chinese are certainly not protected against impaired

glucose tolerance and non-insulin dependent diabetes mellitus,^{13,25} and the Chinese in Mauritius are also prone to develop high blood pressure.²⁶ Several recent reports have suggested that the well known association between blood pressure and glucose intolerance may be part of a metabolic syndrome characterised by insulin resistance and leading to coronary heart disease.²⁷⁻³⁰ Hyperinsulinaemia and insulin resistance are also associated with an atherogenic plasma lipoprotein profile.^{29,31} Both total and non-high density lipoprotein cholesterol were significantly increased in the Chinese in Mauritius when compared with the Chinese in Beijing. Unfortunately, glucose tolerance was not assessed in Beijing. Other studies have shown very low rates of non-insulin dependent diabetes mellitus in the Chinese in China.³²

Even though absolute levels of blood pressure and plasma lipids in Mauritian Chinese in our study were not exceptionally high, they were by no means low, and were similar to those of the Indian and Creole Mauritians. Earlier studies have shown that Chinese patients with acute myocardial infarction have significantly higher blood pressure and plasma lipid concentrations than do control subjects.^{33,34} This is consistent with the commonly agreed fact that the risk of coronary heart disease increases linearly with blood pressure and plasma cholesterol.^{3,35,36} Other factors than may account for the higher rate of coronary heart disease in Chinese in Mauritius than in Beijing could be differences in dietary habits and physical activity that are likely to be more favourable in Beijing,^{37,38} although comparative data are not available.

CORONARY HEART DISEASE

The World Health Organisation MONICA project baseline data showed that mortality from coronary heart disease among the 40 participating centers was the lowest in men from Beijing, and although the Chinese women did not rank the lowest, their absolute rate was low.¹ The reported incidence (attack rate) of acute myocardial infarction—26/100 000/year in men and 8/100 000/year in women aged 25-64—confirmed that the true occurrence of acute myocardial infarction is very low.¹⁵ We studied the same population in Beijing, and can thus be confident that the low prevalence of coronary heart disease shown by our survey reflects a low incidence of coronary heart disease in mainland China.

Repolarisation findings in electrocardiograms are not specific for coronary heart disease but can also be related to other conditions, such as carditis; they can also be due to automatic influences, chest configuration, obesity, and the fasting state. Electrocardiograms are, however, often used in cardiovascular surveys to provide estimates for the occurrence of coronary heart disease. Particularly in developing countries, where epidemiological studies of coronary heart disease are rare, a cross sectional analysis of electrocardiographic abnormalities will often be the only feasible way to obtain such information. Although these do not provide a complete picture of the occurrence of coronary heart disease in the population, they are valuable for determining the subsequent risk for coronary heart disease.³⁹⁻⁴³

CONCLUSIONS

Our data confirm the previous observation based on routine statistics that coronary heart disease is common in Mauritius⁴⁴ and they show that heart disease is certainly important in the Chinese community. Similarly, it is a major public health problem in the Chinese population of Singapore (where rates

are already higher than in Australia), the United States, and many European countries.⁴⁵⁻⁴⁶ Therefore primordial prevention of coronary heart disease in the People's Republic of China is the major challenge for preventive medicine there.^{12,47,48} We believe that the situation seen in Singapore and Mauritius is not inevitable and that it is possible to prevent such a major epidemic of coronary heart disease from occurring in China. Further assessment of trends in coronary heart disease mortality, morbidity, and risk factors will show whether this will be successful in reality.

The study in Mauritius was undertaken with the support and collaboration of the Ministry of Health of Mauritius, the World Health Organisation (Geneva and Mauritius), the Royal Southern Memorial Hospital (Melbourne, Australia), the University of Newcastle upon Tyne (United Kingdom), and the National Public Health Institute (Helsinki, Finland). Field and technical assistance was provided by staff from all of these parent organisations, and individual contributions are gratefully acknowledged. We are grateful for the support of the British Diabetic Association and Boehringer Mannheim. The study was partially supported by the National Institute of Health (grant DK-25446) and the Finnish Heart Foundation. The Beijing Heart, Lung, and Blood Vessel Centre became one of the collaborating centres of the WHO MONICA project in 1981 and was named Sino-MONICA-Beijing. We are grateful for the support of the principal investigators, Professor Wu Ying-Kai and Drs Wu Zhao-Su, Yao Chong-Hua, Hong Zhao-Kuang. This study was also supported by the Ministry of Public Health of the People's Republic of China.

- 1 Tuomilehto J, Kuulasmaa K, Torppa J, for WHO MONICA Project. Geographic variation in mortality from cardiovascular diseases. Baseline data on selected population characteristics and cardiovascular mortality. *World Health Stat Q* 1987;40:171-84.
- 2 Pajak A, Kuulasmaa K, Tuomilehto J, Ruokokoski E. The WHO MONICA Project: geographic variation in the major risk factors of coronary heart disease in men and women, aged 35-64 years. *World Health Stat Q* 1988;41:115-40.
- 3 Keys A. *Seven countries. A multivariate analysis of death and coronary heart disease*. Cambridge, Massachusetts: Harvard University Press, 1980.
- 4 Byington R, Dyer AR, Garside D, Liu K, Hoss D, Stamler J, et al. Recent trends of major coronary risk factors and CHD mortality in the United States and other industrialised countries. In: Havlik RJ, Feinleib M, eds. *Proceedings of the Conference on the Decline in Coronary Heart Disease Mortality*. Washington, DC: US Department of Health, Education and Welfare, Public Health Service, National Institutes of Health, 1979;340-80. (NIH publication No 79-1610.)
- 5 INTERSALT Cooperative Research Group. INTERSALT: An international study of electrolyte excretion and blood pressure: results for 24 hour urinary sodium and potassium excretion. *BMJ* 1988;297:319-28.
- 6 ERICA Research Group. The CHD risk-map of Europe. The first report of the WHO-ERICA project. *Eur Heart J* 1988;9(suppl):1-36.
- 7 Kato H, Tillotson J, Nichaman MZ, Rhoads GG, Hamilton HB. Epidemiologic studies of coronary heart disease and stroke in Japanese men living in Japan, Hawaii, and California. Serum lipids and diet. *Am J Epidemiol* 1973;97:372-85.
- 8 Kagan A, Harris BR, Winkelstein WJ, Johnson KG, Kato H, Syme SL, et al. Epidemiologic studies of coronary heart disease and stroke in Japanese men living in Japan, Hawaii, and California. Demographic, physical, dietary, and biochemical characteristics. *Journal of Chronic Diseases* 1974;27:345-64.
- 9 Worth RM, Kato H, Rhoads GG, Kagan A, Syme SL. Epidemiologic studies of coronary heart disease and stroke in Japanese men living in Japan, Hawaii, and California. Mortality. *Am J Epidemiol* 1975;102:481-90.
- 10 Robertson TL, Kato H, Rhoads GG, Kagan A, Marmot M, Syme SL, et al. Epidemiologic studies of coronary heart disease and stroke in Japanese men living in Japan, Hawaii, and California. Incidence of myocardial infarction and death from coronary heart disease. *Am J Cardiol* 1977;39:239-43.
- 11 Stamler J. Opportunities and pitfalls in international comparisons related to patterns, trends and determinants of CHD mortality. *Int J Epidemiol* 1989;18(suppl 1):s3-18.
- 12 Yao CH, Wu ZS, Hong ZG, Xu XM, Zhang M, Wu YK. Risk factors of cardiovascular diseases in Beijing. *Chinese Med J* 1988;101:901-5.
- 13 Dowse GK, Gareeboo H, Zimmet PZ, Alberti KGM, Tuomilehto J, Fareed D, et al. High prevalence of NIDDM and impaired glucose tolerance in Indian, Creole, and Chinese Mauritians. *Diabetes* 1990;39:390-6.
- 14 Rose G, Reid DD, Hamilton PJS, McCartney P, Keen H, Jarrett RJ. Myocardial ischaemia, risk factors and death from coronary heart disease. *Lancet* 1977;i:105-9.
- 15 Wu ZS, Yao CH, Chen DY, Li N, Zhang M, Wu YY. The Sino-MONICA-Beijing study: report on results between 1984 and 1986. *Acta Med Scand* 1987;(suppl 728):60-6.
- 16 Rhoads GG, Dahlen G, Berg K, Morton NE, Dannenberg AL. Lp(a) lipoprotein as a risk factor for myocardial infarction. *JAMA* 1986;256:2540-4.
- 17 Trembath RC, Thomas DJB, Hendra TJ, Yudkin JS, Galton DJ. Deoxyribonucleic acid polymorphism of the apoprotein AI-CIII-AIV gene cluster and coronary heart disease in non-insulin-dependent diabetes. *BMJ* 1987;294:1577-8.
- 18 Austin MA, King M-C, Bawol RD, Hulley SB, Friedman GD. Risk factors for coronary heart disease in adult female twins: genetic heritability and shared environmental effects. *Am J Epidemiol* 1987;125:308-18.

- 19 Xiang KS, Cox NJ, Sanz N, Huang P, Karam JH, Bell GI. Insulin-receptor and apolipoprotein genes contribute to development of NIDDM in Chinese Americans. *Diabetes* 1989;38:17-23.
- 20 Yu YH. Prevalence and unawareness of hypertension in the petrochemical industrial population in China. *Prev Med* 1986;15:643-51.
- 21 Zhang HX, Wang ZY, Gou ZY, Jin LI, Hao XZ. Ten-year report on the community control of hypertension and stroke in a rural district of Shijiazhuang, China. In: Yamori Y, Strasser T, eds. *New horizons in preventing cardiovascular diseases*. Amsterdam: Elsevier, 1989:259-62.
- 22 Zimmet P, Dowse G, LaPorte R, Finch C, Moy C. Epidemiology—its contribution to understanding of the etiology, pathogenesis, and prevention of diabetes mellitus. In: Creutzfeldt W, Lefebvre P, eds. *Diabetes mellitus: pathophysiology and therapy*. Berlin: Springer-Verlag, 1989:5-26.
- 23 Linda MG. Diabetes mortality among Chinese migrants to New York city. *Human Biol* 1984;56:449-58.
- 24 McKeigue PM, Miller GJ, Marmot MG. Coronary heart disease in South Asians overseas: a review. *J Clin Epidemiol* 1989;42:597-609.
- 25 Dowse GK, Gareebou H, Zimmet PZ, Alberti KGMM, Tuomilehto J, Finch CF, et al. Abdominal obesity and physical inactivity as risk factors for NIDDM and impaired glucose tolerance in Indian, Creole and Chinese Mauritians. *Diabetes Care* 1991;14:271-81.
- 26 Nan L, Tuomilehto J, Dowse G, Zimmet P, Gareebou H, Chitson P, et al. Prevalence and medical care of hypertension in four ethnic groups in the newly industrialised nation of Mauritius. *J Hypertension* 1991;9:859-66.
- 27 Reaven GM, Hoffman BB. A role for insulin in the aetiology and course of hypertension? *Lancet* 1987;ii:435-7.
- 28 DeFronzo RA, Cooke C, Andres R, Faloona GR, Davis J. The effect of insulin in renal handling of sodium, potassium, calcium, and phosphate in man. *J Clin Invest* 1975;55:845-55.
- 29 Tobey TA, Greenfield M, Kraemer F, Reaven GM. Relationship between insulin resistance, insulin secretion, very low density lipoprotein kinetics and plasma triglyceride levels in normotriglyceridemic man. *Metabolism* 1981;30:165-71.
- 30 Modan M, Halkin H, Almog S, Lusky A, Esnol A, Shefi M, et al. Hyperinsulinemia: a link between hypertension, obesity and glucose intolerance. *J Clin Invest* 1985;75:809-17.
- 31 Orchard TJ, Becker DJ, Bates M, Kuller LH. Plasma insulin and lipoprotein concentrations: an atherogenic association? *Am J Epidemiol* 1983;118:326-37.
- 32 Zhong XL. Diabetes mellitus survey in China. *Chinese Med J* 1982;95:423-30.
- 33 Schwartzkopff W, Schleicher J, Pottins I, Yu SB, Han CZ, Du DY. Lipids, lipoproteins, apolipoproteins, and other risk factors in Chinese men and women with and without myocardial infarction. *Atherosclerosis* 1990;82:253-9.
- 34 Luo CI, Lui M, Ping Y, Gou Z. Blood lipid analysis in coronary heart disease and significance. *Tianjin Medical Journal* 1986;1:10.
- 35 Levy D, Kannel WB. Cardiovascular risk: new insights from Framingham. *Am Heart J* 1988;116:266-72.
- 36 Stamler J, Wentworth D, Neaton JD. Is the relationship between serum cholesterol and risk of premature death from coronary heart disease continuous and graded? *JAMA* 1986;256:2823-8.
- 37 Yao CH, Hao EH, Wu YY, Dong L, Wu YK. Dietary survey in Beijing in 1983. *Chinese Med J* 1985;98:439-41.
- 38 Gong LS, Wang CX, Qian JA, Xu DH, Qian YS. Cardiovascular effects of Taijiquan and Qigong exercises. In: Yamori Y, Strasser T, eds. *New horizons in preventing cardiovascular diseases*. Amsterdam: Elsevier, 1989:163-7.
- 39 Rose G, Baxter PJ, Reid DD, McCartney P. Prevalence and prognosis of electrocardiographic finding in middle-aged men. *Br Heart J* 1978;40:636-43.
- 40 Uusitupa M, Pyörälä K, Raunio H, Rissanen V, Lampainen E. Sensitivity and specificity of Minnesota code Q-QS abnormalities in the diagnosis of myocardial infarction verified at autopsy. *Am Heart J* 1983;106:753-7.
- 41 Liao Y, Liu K, Dyer A, Schoenberger JA, Shekelle RB, Collette P, et al. Sex differential in the relationship of electrocardiographic ST-T abnormalities to risk of coronary death: 11.5 year follow-up findings of the Chicago Heart Association detection project in industry. *Circulation* 1987;75:347-52.
- 42 Knutsen R, Knutsen S, Curb JD, Reed D, Kautz J, Yano K. The predictive value of resting electrocardiograms for 12-year incidence of coronary heart disease in the Honolulu heart program. *J Clin Epidemiol* 1988;41:293-7.
- 43 Kannel WB, Abbott RD. Incidence of prognosis of unrecognised myocardial infarction: an update from the Framingham study. *N Engl J Med* 1984;311:1144-7.
- 44 Brissonnette LGG, Fareed DS. Cardiovascular diseases as a cause of death in the island of Mauritius, 1972-1980. *World Health Stat Q* 1985;38:163-75.
- 45 Hughes K, Yeo PPB, Lun KC, Sothy SP, Thai AC, Wang KW, et al. Ischaemic heart disease and its risk factors in Singapore in comparison with other countries. *Ann Acad Med Singapore* 1989;18:245-9.
- 46 Hughes K, Yeo PPB, Lun KC, Thai AC, Sothy SP, Wang KW, et al. Cardiovascular diseases in Chinese, Malays, and Indians in Singapore. II. Differences in risk factor levels. *J Epidemiol Community Health* 1990;44:29-35.
- 47 Tao SC, Huang ZD, Wu XG, Zhou BF, Xiao ZK, Hao JS, et al. CHD and its risk factors in the People's Republic of China. *Int J Epidemiol* 1989;18 (suppl 1):S159-63.
- 48 Vartiainen E, Dianjun D, Marks JS, Korhonen H, Guanyi G, Ze-Yu G, et al. Mortality, cardiovascular risk factors, and diet in China, Finland, and the United States. *Public Health Rep* 1991;106:41-6.

(Accepted 2 April 1992)

Growth hormone and tumour recurrence

A L Ogilvy-Stuart, W D J Ryder, H R Gattamaneni, P E Clayton, S M Shalet

Abstract

Objective—To determine whether using growth hormone to treat radiation induced growth hormone deficiency causes tumour recurrence.

Design—Comparison of tumour recurrence rates in children treated with growth hormone for radiation induced deficiency and an untreated population. Computed tomograms from children with brain tumours were reviewed when starting growth hormone and subsequently.

Setting—North West region.

Patients—207 children treated for brain tumour, 47 of whom received growth hormone and 161 children with acute lymphoblastic leukaemia 15 of whom received growth hormone.

Main outcome measures—Tumour recurrence and changes in appearances on computed tomography.

Results—Among children with brain tumour, five (11%) who received growth hormone had recurrences compared with 42 (26%) who did not receive growth hormone. Also adjusting for other variables that might affect tumour recurrence the estimated relative risk of recurrence was 0.82 (95% confidence interval 0.28 to 2.37). The only child with acute lymphoblastic leukaemia who relapsed while taking growth hormone had relapsed previously before starting treatment. Two of the five children with brain tumours who relapsed had abnormal appearances on computed tomography when growth hormone was started. 14 other children who remained relapse free and had follow up computed tomography showed no deterioration in radiological appearance during treatment.

Conclusions—In this population growth hormone

did not increase the risk of tumour recurrence but continued surveillance is essential. Abnormal results on computed tomography are not a contraindication to treatment with growth hormone.

Introduction

Acute lymphoblastic leukaemia and brain tumours are the two commonest childhood malignancies, the treatment of which has consisted of cranial irradiation with or without adjuvant cytotoxic chemotherapy. Long term management of the endocrine sequelae of treatment, including growth failure, is fundamental to the improved quality of life of these children. The use of growth hormone in children with radiation induced growth hormone deficiency is now widely accepted, but questions still exist about the safety of this mitogenic hormone and whether it might cause a recurrence of a brain tumour or leukaemia.

Studies with small numbers of patients at our centre¹ and others^{2,3} suggested that growth hormone is not responsible for recurrence of brain tumours, but none of these studies applied statistical analysis. An analysis of deaths in recipients of pituitary growth hormone showed brain tumour recurrence to be one of the most common causes, but a comparative group who had not received growth hormone was not available for analysis.⁴ Therefore, it could not be established whether growth hormone contributed to tumour recurrence.

In children treated with growth hormone after treatment for a brain tumour radiographs of the central nervous system often appear abnormal at the start of treatment. No information is available to determine if

Departments of Endocrinology, Radiotherapy, and Statistics, Christie Hospital and Holt Radium Institute, Manchester M20 9BX
A L Ogilvy-Stuart, research fellow in paediatric endocrinology
W D J Ryder, statistician
H R Gattamaneni, consultant radiotherapist
P E Clayton, research fellow in paediatric endocrinology
S M Shalet, consultant endocrinologist

Correspondence to: Dr Shalet.

BMJ 1992;304:1601-5