- et al. Improved prognosis of diffuse histiocytic and undifferentiated lymphoma by use of high dose methotrexate alternating with standard agents (M-BACOD). *J Clin Oncol* 1983;1:91.
- 28 Skarin AT, Canellos GP, Rosenthal DS, Case D, McIntyre J, Pinkus C, et al. Moderate dose methotrexate (m) combined with bleomycin (B), Adriamycin (A), cyclophosphamide (C), Oncovin (O), and dexamethasone (D), m-BACOD is advanced diffuse histiocytic lymphoma (DHL) [abstract]. *Proceedings of the American Society of Clinical Oncology* 1983;2:220.
- 29 Longo DL, DeVita VT, Duffey PL, Wesley MN, Ihde DC, Hubbard SM, et al. Superiority of ProMACE-CytaBOM over ProMACE-MOPP in the treatment of advanced diffuse aggressive lymphoma: results of a prospective randomized trial. *J Clin Oncol* 1991;9:25.
- 30 Klimo P, Connors JM. MACOP-B chemotherapy for the treatment of diffuse large cell lymphoma. *Ann Intern Med* 1985;102:596.
- 31 Coiffier B, LePage E. Prognosis of aggressive lymphomas: a study of five prognostic models with patients included in the LNH-84 regimen. *Blood* 1989;74:558.
- 32 Shipp MA, Harrington DP, Klatt MK, Jochelson MS, Pinkus GS, Marshall JL, et al. Identification of major prognostic subgroups of patients with large cell lymphoma treated with m-BACOD or M-BACOD. *Ann Intern Med* 1986;104:757.
- 33 Jagannath S, Velasquez WS, Tucker SL, Fuller LM, McLaughlin PW, Manning JT, et al. Tumor burden assessment and its implication for a prognostic model in advanced diffuse large cell lymphoma. *J Clin Oncol* 1986;4:859.
- 34 Hoskins PJ, Ng V, Spinelli J, Klimo P, Connors JM. Prognostic variables in patients with diffuse large-cell lymphoma treated with MACOP-B. *J Clin Oncol* 1991;9:220.

Lesson of the Week

Potential of aluminium absorption by effervescent analgesic tablets in a haemodialysis patient

J Main, M K Ward

Gastrointestinal aluminium absorption is greatly increased by citrate, which may unknowingly be prescribed in effervescent or dispersible tablets

It has recently been shown that in renal failure patients taking aluminium containing preparations as phosphate binders, concurrent administration of citrate enhances intestinal aluminium absorption.¹ This is an important interaction because of the acute and chronic effects of aluminium on brain and bone and the fairly common practice of prescribing citrate to renal failure patients either as a calcium salt or Shohl's solution.^{2,4} We report enhanced aluminium absorption due to concurrent administration of effervescent analgesics.

Case history

A 45 year old woman on home haemodialysis was on a regular treatment with aluminium hydroxide capsules 950 mg thrice daily as a phosphate binding agent. Monthly checks were made of serum aluminium concentrations, which never exceeded 2.5 µmol/l. She developed ophthalmic herpes zoster and was treated with acyclovir tablets and a compound analgesic, co-codamol (paracetamol 500 mg, codeine 8 mg), up to six tablets daily. Ten days later routine blood testing revealed a sudden 10-fold increase in serum aluminium concentration to 18.5 µmol/l (figure). Inquiry revealed that the co-codamol had been dispensed in effervescent form, each tablet containing 1.5 g sodium citrate. After stopping this the aluminium concentrations fell to

acceptable levels over the next three weeks. The patient developed no symptoms or signs of acute aluminium toxicity.

Discussion

Increased serum aluminium concentrations and acute aluminium toxicity produced by the concurrent administration of aluminium salts and citrate containing compounds in patients with renal failure have recently been described.^{2,4} In these cases the citrate was given either to combat metabolic acidosis or as calcium citrate to treat renal bone disease. The increase in aluminium concentrations is due to enhanced gastrointestinal absorption mediated by citrate's properties of increasing aluminium solubility and opening gastrointestinal mucosal tight junctions.¹

These reports recommend that patients with renal failure taking aluminium compounds should not be prescribed citrate containing preparations. Our experience draws attention to the presence of citrate in a commonly used and widely available group of drugs—effervescent or dispersible analgesics. Such compounds can be bought without prescription. Clearly, it is important that renal patients taking aluminium containing compounds are not prescribed and do not self administer citrate based effervescent analgesics. The presence of citrate in these compounds is frequently omitted from descriptions of the drugs in both the *British National Formulary* and the *Data Sheet Compendium*. We suggest that when citrate is used in drug formulations it is clearly labelled to enable renal failure patients and those who prescribe for them to avoid such drugs if indicated.

The Committee on Safety of Medicines has been informed of this interaction.

- 1 Partridge NA, Regnier FE, White JL, Hem SL. Influence of dietary constituents on intestinal absorption of aluminium. *Kidney Int* 1989;35:1413-7.
- 2 Kirschbaum BB, Schoolwerth AC. Acute aluminium toxicity associated with oral citrate and aluminium-containing antacids. *Am J Med Sci* 1989;297:9-11.
- 3 Molitoris BA, Froment DH, Mackenzie TA, Huffer WH, Alfrey AC. Citrate: a major factor in the toxicity of orally administered aluminium compounds. *Kidney Int* 1989;36:949-53.
- 4 Nolin CR, Califano JR, Butzin CA. Influence of calcium acetate or calcium citrate on intestinal aluminium absorption. *Kidney Int* 1990;38:937-41.

(Accepted 27 September 1991)

Renal Offices, Royal Victoria Infirmary, Newcastle upon Tyne NE1 4LP
J Main, senior registrar in nephrology
M K Ward, consultant nephrologist

Correspondence to: Dr Main.

BMJ 1992;304:1686

Change in serum aluminium concentration with time. Arrow indicates date of starting effervescent analgesic treatment