Fat and female fecundity: prospective study of effect of body fat distribution on conception rates

Boukje M Zaadstra, Jacob C Seidell, Paul A H Van Noord, Egbert R te Velde, J Dik F Habbema, Baukje Vrieswijk, Jan Karbaat

Abstract

Objectives—To study the effect of body fat distribution in women of reproductive age on fecundity.

Design—Prospective cohort study of all women who had entered a donor insemination programme.

Setting—One fertility clinic serving a large part of the midwest of the Netherlands.

Subjects—Of 542 women attending the clinic for artificial insemination for the first time, 500 women were eligible for study.

Main outcome measures—Probability of conception per cycle and number of insemination cycles before pregnancy or stopping treatment.

Results—A 0.1 unit increase in waist-hip ratio led to a 30% decrease in probability of conception per cycle (hazard ratio 0.706; 95% confidence interval 0.562 to 0.887) after adjustment for age, fatness, reasons for artificial insemination, cycle length and regularity, smoking, and parity. Increasing age was significantly related to lower fecundity ($p < 0.05$); very lean and obese women were less likely to conceive ($p < 0.10$) as were women with subfertile partners ($p < 0.10$). All other exposure variables were not significantly related to fecundity.

Conclusions—Increasing waist-hip ratio is negatively associated with the probability of conception per cycle, before and after adjustment for confounding factors. Body fat distribution in women of reproductive age seems to have more impact on fertility than age or obesity.

Introduction

Little is known about the effects of body fat distribution on reproductive function in women. Obesity is common among multiparous women¹ but this may indicate that weight gain is the result of multiple pregnancies rather than a prerequisite for conception.² Menstrual disorders have been found to be related to obesity as well as to a predominance of fat in the abdominal region.^{3,4} An abdominal type of fat distribution may be related to increased androgenicity of these women.^{5,6} Frisch suggested that underweight women would also have impaired reproductive function owing to a lack of oestrogen produced in adipose tissue.⁷ It is not known, however, whether body fat distribution has any effect on the ability to conceive (fecundity) independent of weight and menstrual disorders.

A population of women presenting for artificial donor insemination provides an opportunity to study prospectively determinants of fecundity while controlling for possible confounders.⁸ We studied 500 women aged 20 to 42 years presenting for artificial insemination to determine effects of body fat distribution while controlling for weight and other variables on the chance of conception.

Subjects and methods

From January 1986 to January 1988 all women attending a single fertility clinic for the first time

($n = 542$) were asked to participate in the study. Three women refused to participate. The 39 women who did not start treatment before January 1989 were excluded from the analyses. Thus 500 women filled in a questionnaire and had anthropometric measurements taken at intake.

The self administered questionnaire included questions on age, smoking habits, and other lifestyle characteristics. Regularity and length of the menstrual cycles, parity, degree of infertility of partner were ascertained by a doctor based on structured patient history. Women were wearing only light indoor clothing without shoes when anthropometric measurements were taken. These included weight (kg), height (cm), waist circumference at the umbilical level and hip circumference at the level of the widest symphysis (cm). All measurements were taken by one doctor. Quetelet's index (kg/m^2) was calculated as a measure of total fatness (body mass index) and waist to hip ratio as a measure of body fat distribution.

Women were referred to the clinic by a gynaecologist or other specialist (302, 61%) or by a general practitioner (101, 20%) or came to the clinic on their own initiative (85, 17%). (No information was available on 12 (2%).)

The main reasons for presenting for artificial insemination were infertility (azoospermia) or subfertility (oligospermia) of the partner (152 (30%) and 236 (47%) respectively). Other reasons included sterilisation of partner (14, 3%), genetic abnormalities of partner (11, 2%), no partner (63, 13%), and other reasons (24, 5%). For the analysis women with azoospermic or sterilised partners and women with no partner were categorised in the subgroup with infertile partners. Most women had tried to become pregnant for two or three years, and most (412, 82%) had never been pregnant when they entered the study.

Before the first insemination all women were examined by one doctor. The mean menstrual cycle length was 28 days. Cycle length was classified as short (≤ 24 days) in 30 women, normal (25–35 days) in 431, and long (≥ 36 days) in 39. At the start of the treatment 368 women reported having regular cycles (less than four days difference on average). Forty seven women with very irregular and long cycles had ovulation induced by clomiphene. None of the other women received drugs to improve fecundity at the start of insemination treatment.

The youngest woman entering the study was 20 years old, and 10 were 40 or older. Smoking was reported by 260 women.

INSEMINATIONS AND FOLLOW UP

Intracervical inseminations were applied in subsequent menstrual cycles. Timing was based on previous cycle length, examination of the cervical mucus, and basal temperature charts. Frozen semen was used from donors aged between 25 and 45 years with proved fertility (having fathered at least one child) and with sperm properties satisfying the World Health Organisation criteria.

Only first conceptions as a result of artificial in-

TNO Institute of Preventive Health Care, Child Health Division, Postbox 124, 2300 AC Leiden, Netherlands
Boukje M Zaadstra, epidemiologist

Department of Human Nutrition, Agricultural University, Wageningen, Netherlands
Jacob C Seidell, epidemiologist
Baukje Vrieswijk, nutritionist

Department of Epidemiology, University of Utrecht Medical School, Utrecht, Netherlands
Paul A H Van Noord, epidemiologist

Department of Obstetrics and Gynaecology, University Hospital, Utrecht, Netherlands
Egbert R te Velde, professor of reproductive medicine

Department of Public Health and Social Medicine, Erasmus University Medical School, Rotterdam, Netherlands
J Dik F Habbema, professor of medical decision sciences

Cryo Biological Laboratories Bijdorp, Barendrecht, Netherlands
Jan Karbaat, medical director

Correspondence and requests for reprints to: Dr Zaadstra.

BMJ 1993;306:484-7

semination were used for the analysis. Insemination was defined as successful if the woman did not menstruate when expected and subsequently had a positive pregnancy test result. The follow up period lasted three years, and the largest number of cycles observed was 33.

STATISTICAL METHODS

The cumulative probability of conception by insemination cycle was calculated with Kaplan-Meier estimates for two waist-hip ratio groups.⁹ The 41 women who did not report the result of their last insemination cycle (over the entire follow up period) were recorded as not successful up to their last but one cycle. The anthropometric measures were missing for 11 women.

Univariate proportional hazard regression analysis^{10 11} was used to analyse the relation between probability of conception per cycle and each of the following variables separately: waist-hip ratio, body

Cumulative pregnancy rate in 489 women having artificial insemination according to waist-hip ratio (277 women had ratio <0.8 and 212 a ratio ≥0.8). Kaplan-Meier logrank $p=0.0085$

TABLE 1—Distribution of study population over categories of studied variables and percentages of pregnant women within 12 insemination cycles

Variable	No (%) of women (n=489)*	No (%) pregnant after 12 cycles
Waist-hip ratio:		
<0.70†	32 (7)	20 (63)
0.70-0.75	99 (20)	50 (51)
0.75-0.80	146 (30)	69 (47)
0.80-0.85	138 (28)	57 (41)
≥0.85	74 (15)	24 (32)
Body mass index (kg/m ²):		
<20.0	104 (21)	42 (40)
20.0-25.0†	303 (62)	145 (48)
25.0-30.0	60 (12)	29 (48)
≥30.0	22 (5)	4 (18)
Age (years):		
20-24†	67 (14)	34 (51)
25-29	229 (47)	113 (49)
30-34	131 (27)	56 (43)
≥35	62 (13)	17 (27)

* For 11 women no anthropometric measurements available.

† Reference category. For waist-hip ratio and body mass index upper limits are exclusive.

TABLE 2—Univariate and multivariate proportional hazard analysis of waist-hip ratio, age, body mass index, reasons for artificial insemination, cycle length, cycle regularity, smoking, and parity on probability of conception per cycle over total follow up period*

Variable	Univariate analysis			Multivariate analysis		
	Hazard ratio	95% Confidence interval	p Value	Hazard ratio	95% Confidence interval	p Value
Waist-hip ratio (0.1)	0.704	0.568 to 0.871	<0.001	0.706	0.562 to 0.887	<0.0025
Age:			<0.0025			<0.05
23	1.000			1.000		
27	0.977	0.763 to 1.251		0.970	0.750 to 1.253	
31	0.866	0.605 to 1.240		0.869	0.597 to 1.266	
37	0.603	0.382 to 0.952		0.638	0.396 to 1.027	
Body mass index:			<0.05			<0.10
19.0	0.854	0.675 to 1.073		0.837	0.662 to 1.058	
22.3	1.000			1.000		
26.4	0.901	0.749 to 1.084		0.939	0.775 to 1.139	
33.1	0.367	0.152 to 0.886		0.431	0.171 to 1.087	
Reason for artificial insemination:			NS			<0.10
Infertile partner	1.000			1.000		
Subfertile partner	0.778	0.597 to 1.013		0.726	0.555 to 0.950	
Other reasons	1.238	0.734 to 2.088		1.254	0.738 to 2.130	
Cycle length (1)	0.999	0.974 to 1.026	NS	1.003	0.974 to 1.033	NS
Cycle regularity	0.899	0.66 to 1.23	NS	0.886	0.642 to 1.221	NS
Smoking	0.870	0.675 to 1.123	NS	0.914	0.704 to 1.187	NS
Parity	1.015	0.717 to 1.435	NS	0.811	0.566 to 1.162	NS

The relations were analysed by classes if the variable was categorical (cycle regularity, reason for insemination, smoking, parity), linear equation if the variable was continuous and significantly not quadratical (cycle length, waist-hip ratio), and quadratic equation (variable+variable²) if addition of quadratic was significant (body mass index, age). Significance of variables was based on the effect of adding the parameters to the null model (univariate analysis) or the model with all other parameters (multivariate analysis).

For cycle regularity, smoking, and parity the hazard ratios express the difference of the second class with the first and for cycle length and waist-hip ratio the hazard ratios express an increase of 1 day or 0.1 unit respectively. For body mass index and age four values were chosen (medians of the classes in table 1). The hazard ratio and confidence intervals were calculated for these values, where the highest value was taken as "baseline." The result is analogous to the presentation of other variables when it is taken into account that many other values could have been chosen. Consequentially the significance of the parameters cannot be deduced from confidence intervals.

* N=448 women, due to missing values in one of the exposure or confounding variables.

mass index, age, regularity and length of menstrual cycle, smoking, parity, degree of infertility of partner. Multivariate proportional hazard regression^{10 11} was used to analyse the relation between waist-hip ratio and probability of conception per cycle while controlling for the above variables. Unadjusted and adjusted hazard ratios and 95% confidence intervals were calculated. Both regression analyses were calculated from data on 448 women because of missing values in one of the exposure or confounding variables.

Results

Insemination was successful within 12 cycles in 226 women, not successful in 274 women, including 38 women with an unknown result. Table 1 shows the distribution of the women according to waist-hip ratio, body mass index, and age, and the percentage that became pregnant in each category. The distribution of age, waist-hip ratio, and body mass index of the 38 women lost to follow up was not significantly different from that of women with known outcome ($p>0.05$). Abdominal fat preponderance (waist-hip ratio ≥0.80) was quite common (212 women). The percentage of women who became pregnant fell with increasing waist-hip ratio from 63% to 32%. Women with a body mass index under 20 kg/m² were less likely to become pregnant than those with an index of 20-24.9 kg/m². Only 22 women were obese (body mass index ≥30 kg/m²). Just four of these women became pregnant, suggesting a curvilinear relation between weight and fecundity. The percentage of pregnant women declined with older age and among the 62 who were 35 years or older only 17 (27%) became pregnant.

The figure shows that the cumulative pregnancy rate by insemination cycle for women with waist-hip ratios <0.80 (pear shape) was significantly higher than for women with ratios ≥0.80 (apple shape) ($p=0.008$). Because of the small number of women completing 24 cycles or more the suggestion in our data that both groups will eventually have the same pregnancy rate, must be interpreted with caution.

Table 2 shows the unadjusted hazard ratios and 95% confidence intervals for the effect of waist-hip ratio, body mass index, age, regularity and length of cycle, smoking, parity, and reason for artificial insemination on probability of conception per cycle. Waist-hip ratio, age and body mass index were separately related to probability of conception per cycle. The multivariate proportional hazards regression in table 2 shows that the relation between waist-hip ratio and probability of conception per cycle, after controlling for all the other

mentioned variables, remained essentially the same: waist-hip ratio yielded the highest independent significant contribution to probability of conception per cycle (hazard ratio 0.706, 95% confidence interval 0.562 to 0.887).

Discussion

This study shows that increasing waist-hip ratio is associated with fecundity in healthy women, independently of age, fatness, length and irregularity of menstrual cycle, parity, degree of infertility of partner, and smoking habits. Both underweight and overweight women had a lower chance of becoming pregnant than women with normal weight, although this effect was greater for the obese than for the lean women. In addition, increasing age was associated with a reduced conception rate, which agrees with the results of our previous studies.^{12 13}

WAIST-HIP RATIO

Hartz *et al* showed that in about 12 000 American women aged 20-39 years participating in a slimming programme the degree of obesity and waist-hip ratio were independently and positively related to the prevalence of reported irregular menstruation, oligomenorrhoea (cycles longer than 36 days), and hirsutism.³ Such a relation has been explained as the result of increased androgenicity.^{5 6 14}

Based on information of cycle length and cycle irregularity in the period before insemination only 9% of the women in our study required clomiphene to regulate their cycle. Moreover, no obvious hirsutism was present in any of the women. Our study population therefore contained mainly healthy women of normal weight with apparently normal cycles, which makes comparison with the American study impossible.

Obese women or women with abnormal fat distribution with normal length of cycles (25-35 days) may have less regular cycles than women of normal weight and fat distribution. If this were true the decreased chance of conception could be explained by difficulties in timing of insemination, because timing may be less precise and therefore less successful in women with less regular cycles. However, we found no relation between abnormal fat distribution or obesity and cycle regularity or whether ovulation had to be induced. Therefore, after correcting for all these potentially confounding factors, the highly significant relation between waist-hip ratio and the chance of conceiving, essentially remained the same (table II). The reasons for our finding remain unclear. Insulin resistance (which is clearly associated with abnormal fat distribution or obesity)¹⁵ at an early stage might increase the androgenic microenvironment of the follicle thus decreasing oocyte quality and the chance of conceiving before androgen levels are raised in serum and before cycles become irregular. Moreover, recent findings suggest that increased levels of luteinising hormone, which may also be associated with hyperinsulinaemia, might decrease the spontaneous chance of conceiving and increase spontaneous abortion rates.¹⁶ Unfortunately, no pre-treatment endocrine profiles were available in our study.

Weight loss often restores normal cycle regularity¹⁷ and sex hormone concentrations in obese women,¹⁸ which indicates that obesity itself and not primary endocrine dysfunction is the cause of the reproductive problems. Further research is necessary to investigate mechanisms responsible for the low conception rate in normal weight women with an increased waist-hip ratio.

OTHER FACTORS AFFECTING FECUNDITY

We also observed a tendency of a reduced chance of

conception in women with a body mass index under 20 kg/m². The association between leanness and fertility has been extensively explored by Frisch, who has suggested that menstruation and ovulatory cycles can only start above a certain threshold ratio of lean body mass to fat mass.⁷ At body weights close to the required minimum women may still have anovulatory cycles even if they are menstruating because of a lack of peripherally produced oestrogen in adipose tissue.⁷ The problem can be reversed by gaining weight.⁷ However, this explanation does not hold for our results because most of the lean women in our population had regular cycles with biphasic basal body temperatures and did not require induction of ovulation. Moderate weight loss seems to lead to subtle ovulation disorders and therefore to decreasing fecundity.

Changes in weight and waist-hip ratio during the three year follow up period could have led to misclassification of women. However, it is known that body mass index on average changes only 0.10 kg/m² a year.¹⁹ Likewise no appreciable changes in waist-hip ratio occur with small fluctuations in weight.²⁰ Proportional hazard regression analysis over a 12 cycle follow up yielded essentially the same results as that over the total follow up period.

Length and regularity of menstrual cycle were based on structured patient history at the start of the treatment. Prospective information on cycle variability during treatment would have led to potential selection bias. A minimum number of three cycles is needed to estimate variability, and women with low waist-hip ratios—that is more fertile women—would be more likely to be excluded from such an analysis, leading to an under-estimation of the relation between waist-hip ratio and fecundity.

Smoking has been associated with delayed conception.²¹ In our study women who smoked had lower conception rates, though this did not reach significance. In the adjusted analysis subfertility of the partner became indicative for the fertility of the woman because of the fact that within this subgroup women with lower fertility are over-represented among those entering a donor insemination programme.²²

CONCLUSIONS

In summary, in a prospective study of healthy women presenting for artificial insemination treatment we observed that increasing waist-hip ratio was negatively associated with the probability of conception per insemination cycle, before and after adjustment for age, fatness, cycle characteristics, smoking, parity, degree of infertility, and partner.

Our results seem to agree with recent findings of a breast cancer study among postmenopausal women by Sellars *et al*.²³ They observed that a high waist-hip ratio, low parity, and greater age at first pregnancy were more important risk factors for breast cancer among women with a family history of the disease than among those without such a history. However, older age at first pregnancy and fewer children were not independent factors. They were unable to determine whether this risk profile reflected cultural inheritance or difficulty in becoming pregnant. Our results suggest that the second is true: women with high waist-hip ratio have difficulty in becoming pregnant.

As the prevalence of a high waist-hip ratio among healthy women is larger than the prevalence of obesity (15% and 5% respectively in our study) body fat distribution apparently has more impact on fertility than obesity.

Our findings are based on a cohort of women seeking insemination treatment because their partners had a fertility problem. Thus the population of women is primarily selected by the fertility of the partner and not the women and probably is a fair reflection on

the normal population of women.⁸ Though our results need replication, this suggests that our results may be applicable to women attempting to conceive naturally.

The authors thank the women who participated in the study and the staff of Bijldorp Laboratories for their invaluable help; S P Verloove-Vanhorick and J J Radder for critically reading the manuscript; C W N Looman for constructive help; M M van der Klaauw, L M Rhodenborgh for support in data handling; and M Huls-van Vliet for secretarial help.

- 1 Hartz AJ, Barbieri PN, Wong A, Katayama KP, Rimm AA. The association of obesity with infertility and related menstrual abnormalities in women. *Int J Obesity* 1979;3:57-73.
- 2 Newcombe RG. Development of obesity in parous women. *J Epidemiol Community Health* 1982;36:306-9.
- 3 Hartz AJ, Rupley DC, Rimm AA. The association of girth measurements with disease in 32 856 women. *Am J Epidemiol* 1984;119:71-80.
- 4 Rogers J, Mitchell GW. The relation of obesity to menstrual disturbances. *N Engl J Med* 1952;247:53-5.
- 5 Evans DJ, Hoffman RG, Kalkhoff RK, Kissebah AH. Relationship of androgenic activity to body fat topography and metabolic profiles in premenopausal women. *J Clin Endocrinol Metab* 1983;57:304-10.
- 6 Seidell JC, Cigolini M, Chazewska J, Ellsinger BM, Biase G di, Björntorp P, et al. Androgenicity in relation to body fat distribution and metabolism in 38 year old women—the European fat distribution study. *J Clin Epidemiol* 1990;43:21-34.
- 7 Frisch RE. Fatness and fertility. *Sci Am* 1988;258:70-7.
- 8 Noord-Zaadstra BM van, Karbaat J, te Velde ER, Habbema JDF, Van der Maas PJ. The study of risk habits in reproductive and perinatal epidemiologic research: the use of a donor inseminated population of women. *Paediatr Perinat Epidemiol* 1989;3:11-8.
- 9 Kaplan EK, Meier P. Non-parametric estimation from incomplete observations. *Journal of the American Statistical Association* 1958;53:457-81.

- 10 Cox DR. Regression models and life tables. *Journal of the Royal Statistical Society [series B]* 1972;34:187-202.
- 11 Aitkin MD, Anderson AO. *Statistical modelling in GLIM*. Oxford: Clarendon Press, 1989.
- 12 Noord-Zaadstra BM van, Habbema JDF, Karbaat J, Van der Maas PJ. Conceptiepercentages bij donorinseminatie: een onderzoek by 659 vrouwen die met vers donorsemen werden geïnsemineerd. *Ned Tijdschr Geneesk* 1980;124:581-7.
- 13 Noord-Zaadstra BM van, Looman CWN, Alsbach HA, Habbema JDF, te Velde ER, Karbaat J. Delaying childbearing: effect of age on fecundity and outcome of pregnancy. *BMJ* 1991;302:1361-5.
- 14 Friedl KE, Plymate SR. Effect of obesity on reproduction in the female. *Journal of Obesity and Weight Regulation* 1985;4:129-45.
- 15 Pasquali R, Antenucci D, Casimiri F, Venturoli S, Paradisi R, Fabbri R, et al. Insulin as a factor of increased androgen production in women with obesity and polycystic ovaries. *J Endocrinol Invest* 1987;10:575-9.
- 16 Regan L, Owen EJ, Jacobs HS. Hypersecretion of luteinizing hormone, infertility, and miscarriage. *Lancet* 1990;336:1141-4.
- 17 Mitchell GW, Rogers J. The influence of weight reduction on amenorrhea in obese women. *N Engl J Med* 1953;249:835-7.
- 18 Grenman S, Ronnema T, Iriala K, Kaihola HL, Gross M. Sex steroid, gonadotrophin, cortisol and prolactin levels in healthy, massively obese women: correlation with abdominal fat cell size and effect of weight reduction. *J Clin Endocrinol Metab* 1986;63:1257.
- 19 Rookus MA, Burema J, Hof van't MA, Durenberg R, Hautvast JGAJ. The development of body mass index in young adults. II. Interrelationships of level, change and fluctuation, a four-year longitudinal study. *Hum Biol* 1987;59:617-30.
- 20 Shimokata H, Andres R, Coon PJ, Elahi D, Muller D, Tobin JD. Studies in the distribution of body fat. II. Longitudinal effects of change in weight. *Int J Obesity* 1989;13:455-64.
- 21 Baird DD, Wilcox AJ. Cigarette smoking associated with delayed conception. *JAMA* 1985;253:2979-83.
- 22 Emperaire JC, Gauzere-Soumireu E, Audebert AJM. Female fertility and donor insemination. *Fertil Steril* 1982;37:90-3.
- 23 Sellers TA, Kushi LH, Potter JD, Kaye SA, Nelson CL, McGovern PG, et al. Effect of family history, body-fat distribution, and reproductive factors on the risk of postmenopausal breast cancer. *N Engl J Med* 1992;326:1323-9.

(Accepted 9 December 1992)

Diary keeping in asthma: comparison of written and electronic methods

M E Hyland, C A P Kenyon, R Allen, P Howarth

Abstract

Objective—To determine the extent to which symptom diaries of asthmatic patients are inaccurate or based on retrospective recall.

Design—Comparison of electronic and pencil and paper diaries. Both forms were completed twice daily at home for 14 days.

Setting—Outpatient clinic.

Subjects—24 asthmatic outpatients also tested for severity of asthma and for anxiety.

Results—More sessions were missed in the evening than in the morning for both types of diaries. Significantly more retrospective entries were made in the evening (26 entries, 14 patients) than in the morning (6 entries, 3 patients). Discrepant entries of peak expiratory flow accounted for 15% of those made on the appropriate day, and three quarters of patients made at least one discrepant entry. Variation in peak expiratory flow was significantly related to number of discrepancies and number of missing days, and anxiety score was significantly related to number of missing days. About a fifth of written entries may have errors.

Conclusion—Poor diary completion may result from having unreasonable expectations of patients and giving incomplete instructions. Electronic, time coded diaries could ensure better quality of records.

Introduction

Daily recording of diary cards of subjective and objective assessments of the severity of disease is commonly used to monitor chronic conditions, such as asthma, which exhibit an intrinsic variability. Self completed diaries are preferable to single questionnaires or clinic based consultation because accurate

recall of symptoms may be affected by memory bias¹ and single measures may be misleading. The validity of the diary method requires, however, that patients complete their diaries on the day stated rather than retrospectively, and that the patient's record—for example, of peak expiratory flow—is accurate. Few researchers who use this method regularly will not have doubted, at one time or other, that these assumptions are not always met. For example, symptom diaries that have been completed in blocks of different coloured ink arouse a suspicion that the patient may have been completing the diary in blocks of days rather than daily. Some researchers try to counteract this problem by frequently collecting diary forms,² but this is a burdensome procedure and the actual extent to which diaries are inaccurate or based on retrospective recall is unknown.

Developments in electronic diaries allow the problem of retrospective recall to be tested experimentally. The Asthma Question Box is a modified Psion Organiser that is programmed to ask patients questions about their symptoms and to allow them to record peak expiratory flow. The computer is also programmed automatically to store the actual date of data entry, without displaying this information.

The purpose of this study is to compare results from the Asthma Question Box with a conventional paper and pencil symptom diary to obtain information about missing records, retrospective completion (from an analysis of electronic date records), and accuracy (from comparison of data entry by the two methods).

Methods

As part of a larger study in which asthmatic outpatients were told that assessments were being

Department of Psychology,
University of Plymouth,
Plymouth PL4 8AA
M E Hyland, reader
C A P Kenyon, senior lecturer

University Department of
Medicine, Southampton
General Hospital,
Southampton SO9 4XY
R Allen, medical student
P Howarth, consultant
physician

Correspondence to:
Dr Hyland.

BMJ 1993;306:487-9