Controversies in Treatment

Should smokers be offered coronary bypass surgery?

When resources are restricted and some form of rationing of health care is effectively taking place is it ethical to restrict the access of patients to particular treatments if there are alterable factors in their lifestyle which make the treatment less likely to work?

Mr John Bailey, cardiothoracic surgeon, and Dr Matthew Shiu, general practitioner, have disagreed in a television debate on whether smokers should be given the same opportunity for coronary artery bypass graft surgery as non-smokers—or any opportunity at all. We invited them to put their differing views, and then invited a surgeon from another discipline and another general practitioner, both interested in medical ethics, to comment.

Coronary bypass surgery should not be offered to smokers

M J Underwood, J S Bailey

Cigarette smoking accelerates the progression of coronary artery disease^{1,2} and is a determinant of early death in patients awaiting coronary bypass surgery.³ Optimal surgical treatment results in substantial reduction of symptoms with increased exercise tolerance, freedom from angina, and in certain subgroups prolonged survival.⁴⁻⁶

Autogenous saphenous vein continues to be widely used as a conduit for coronary revascularisation.⁷ Unfortunately, coronary vein graft disease is a major cause of mortality and morbidity and is responsible for recurrent angina, myocardial infarction, and compromised ventricular function.⁸ One in 10 vein grafts undergo thrombotic occlusion within two weeks of operation,⁹ and diffuse intimal hyperplasia is a common finding after one year.⁹ By five years, frank atherosclerotic lesions are identifiable in many grafts.¹⁰

Early patency rates may be improved with antiplatelet agents such as aspirin and dipyridamole¹¹ but this protective effect is greatly reduced in patients who continue to smoke,¹² presumably as a result of increased thrombogenicity¹³⁻¹⁶ and enhanced platelet reactivity,¹⁷ which is well recognised in smokers. Late thrombotic occlusion is also considerably increased in patients who continue to smoke after surgery,¹⁸ and this correlates well with reports of increased graft occlusion in smokers undergoing peripheral vascular reconstruction.^{19,20} The atherosclerotic process in vein grafts is identical to that of native coronary arteries⁸ and the risk factors, including smoking, are the same. These atheromatous changes correlate positively with serum lipid concentrations^{21,22} and a reduction of the protective high density lipoprotein cholesterol^{23,24} is well recognised in smokers.²⁵ Not surprisingly, a higher incidence of vein graft atherosclerosis has been reported after surgery in smokers.^{21,26}

Vein graft failure is the major indication for reoperation,²⁷ and reoperation is nearly twice as expensive as "first time" coronary surgery,²⁸ carries twice the mortality,^{29,30} and is associated with a 50% net decrease in the overall rehabilitation status of the patient afterwards.²⁸ It is in the interest of patients, surgeons, and health departments that, before surgery, patients

are carefully assessed to ensure that they will obtain maximum benefit from the operation and that the risk of subsequent reoperation is minimised.

The problems associated with performing coronary surgery on patients who continue to smoke are not confined to risks of vein graft occlusion and reoperation in the future. Immediate postoperative complications are commoner, leading to increased mortality and morbidity; these include pulmonary complications (increased almost fourfold),^{31,32} sternal infection, mediastinal infection, and consequently a prolonged hospital stay.³³

"Subjecting patients to the increased risks of surgery in the face of a remediable cause is not justified."

We recognise that smoking is an addiction and that many patients need expert help to stop. With this in mind, all patients are counselled before operation in our department by staff trained to give expert advice regarding diet, exercise, and cessation of smoking and to identify other factors, such as hyperlipidaemia, that may require specialist treatment. Such schemes are successful in increasing high density lipoprotein cholesterol and reducing the proportion of smokers in groups of patients undergoing coronary artery surgery.³⁴

The current national target is for 300-400 coronary operations per million population to be performed per year,³⁵ although a figure of 500-600 is probably a more realistic reflection of modern cardiological and surgical practice worldwide. Without substantial increases in resources, additional pressure will be put on the already growing waiting lists for coronary surgery. Patients who smoke spend longer in hospital and have poorer results from surgery than non-smokers. Treating them deprives patients who have never smoked or who have stopped smoking of more efficient and effective surgery.

Department of
Cardiothoracic Surgery,
Groby Road Hospital,
Leicester LE3 9QE

M J Underwood, research
fellow

J S Bailey, consultant surgeon

David Rydderch Health
Centre, Birmingham
B32 3SD

Matthew Shiu, general
practitioner

King's College School of
Medicine and Dentistry,
London SE5 8RX

Roger Higgs, professor of
general practice and primary
care

Wessex Neurological
Centre, Southampton
General Hospital,
Southampton SO9 4XY
John Garfield, consultant
neurosurgeon

BMJ 1993;306:1047-50

Considerable evidence exists regarding the adverse effect of continued smoking on vein graft patency rates and early operative mortality and morbidity. One study has reported a reduction in the relative survival of younger patients who smoke,³⁶ and, although little work has been done, it would seem likely that continued smoking will negate the beneficial effect of surgery in patients whose life expectancy would be expected to improve after operation. Subjecting patients who continue to smoke, and for whom the only indication for operation is the relief of angina, to the increased risks of surgery in the face of a remediable cause is not justified.

- 1 Nikutta P, Lichtlen PR, Wiese B, Jost S, Dekkers J, Rafflenbeul W, *et al*. Influence of cigarette smoking on the progression of coronary artery disease within three years: results of the INTACT study. *J Am Coll Cardiol* 1990;15:181A.
- 2 Fielding JE. Smoking. Health effects and control (part one). *N Engl J Med* 1985;313:491-8.
- 3 Suttrop MJ, Kingma JH, Vos J, Koomen EM, Tijssen JGP, Vermeulen FEC, *et al*. Determinants for early mortality in patients awaiting coronary artery bypass graft surgery: a case-control study. *Eur Heart J* 1992;13:238-42.
- 4 CASS Principal Investigators and their Associates. Coronary artery surgery study (CASS): a randomised trial of coronary artery surgery: quality of life in patients randomly assigned treatment groups. *Circulation* 1983;68:951-60.
- 5 Peduzzi P, Hultgren H, Thomsen J, Detre K. Ten year effect of medical and surgical therapy on quality of life: Veterans Administration cooperative study of coronary artery surgery. *Am J Cardiol* 1987;59:1017-23.
- 6 Silverman KJ, Grossman W. Angina pectoris: natural history and strategies for evaluation and management. *N Engl J Med* 1984;310:1712-7.
- 7 Cameron EWJ, Walker WS. Coronary artery bypass surgery. *BMJ* 1990;300:1219-20.
- 8 Badimon JJ, Ip J, Badimon L, Fuster V. Thrombosis and accelerated atherosclerosis in coronary bypass surgery and restenosis after percutaneous transluminal coronary angioplasty: implications for therapy. *Coronary Artery Disease* 1990;2:170-9.
- 9 ACC/AHA Task Force. Guidelines and indications for coronary artery bypass graft surgery. A report of the American College of Cardiology/American Heart Association Task Force on assessment of diagnostic and therapeutic cardiovascular procedures. *J Am Coll Cardiol* 1991;17:543-89.
- 10 Smith SH, Geer JC. Morphology of saphenous vein-coronary artery bypass grafts: 7-116 months postoperatively. *Arch Pathol Lab Med* 1983;107:13-8.
- 11 Goldman S, Copeland J, Moritz T, Henderson W, Zadina K, Ovitt T, *et al*. Improvement in early saphenous vein patency after coronary artery bypass surgery with antiplatelet therapy: results of a Veterans Administration cooperative study. *Circulation* 1988;77:1342-52.
- 12 Bosch X, Perez-Villa F, Sanz G. Effect of smoking habits on the preventive effect of aspirin and dipyridamole on early aortocoronary bypass occlusion. *Eur Heart J* 1991;12:170.
- 13 Levine PH. An acute effect of cigarette smoking on platelet function. *Circulation* 1973;48:619-23.
- 14 Galea G, Davidson RJL. Haematological and haemorheological changes associated with cigarette smoking. *J Clin Pathol* 1985;38:978-84.
- 15 Fitzgerald GA, Oates JA, Nowak J. Cigarette smoking and haemostatic function. *Am Heart J* 1988;115:267-71.
- 16 Wilhelmsson L, Svardsudd K, Korsan-Bengtsson K. Fibrinogen as a risk factor for stroke and myocardial infarction. *N Engl J Med* 1984;311:501-5.
- 17 Fuster V, Chesebro JH, Frye RL, Elveback LR. Platelet survival and the development of coronary artery disease in the young adult: effects of cigarette smoking, strong family history and medical therapy. *Circulation* 1981;63:546-51.
- 18 Solymoss BC, Nadeau P, Millette D, Inge B, Campeau L. Late thrombosis of saphenous vein coronary bypass grafts related to risk factors. *Circulation* 1988;78(suppl I):140-3.
- 19 Wiseman S, Kenchington G, Dain R, Marshall CE, McCollum CN, Greenhalgh RM, *et al*. Influence of smoking and plasma factors on the patency of femoropopliteal grafts. *BMJ* 1989;298:643-6.
- 20 Greenhalgh RM, Laing SP, Cole PV, Taylor GW. Smoking and arterial reconstruction. *Br J Surg* 1981;68:605-7.
- 21 Nietzel GF, Baboriak JJ, Pintar K, Quershi I. Atherosclerosis in aorto-coronary bypass grafts. Morphological study and risk factor analysis 6-12 years after surgery. *Arteriosclerosis* 1986;6:594-600.
- 22 Campeau L, Enjalbert M, Lesperance J, Bourassa MG, Kwiterovich P, Wacholder S, *et al*. The relation of risk factors to the development of atherosclerosis in saphenous vein bypass grafts and the progression of the disease in the native circulation. A study 10 years after aortocoronary bypass surgery. *N Engl J Med* 1984;311:1329-32.
- 23 Gordon DJ, Probstfield JL, Garrison RJ, Neaton JD, Castelli P. High density lipoprotein cholesterol and cardiovascular disease: four year prospective American studies. *Circulation* 1989;79:8-15.
- 24 Stampfer MJ, Sacks FM, Salvini S, Willett WC, Hennekens CH. A prospective study of cholesterol, apolipoproteins and the risk of myocardial infarction. *N Engl J Med* 1991;325:373-81.
- 25 Criqui MH, Wallace RB, Heiss G, Mislakel M, Schonfeld G, Jones GTL. Cigarette smoking and plasma high-density cholesterol. *Circulation* 1980;62(suppl IV):70-6.
- 26 FitzGibbon GM, Leach AJ, Kafka HP. Atherosclerosis of coronary artery bypass grafts and smoking. *Can Med Assoc J* 1987;136:45-7.
- 27 Vekkala K, Jarvinen A, Virtanen K, Keto P, Pellinen T, Salminen US, *et al*. Indications for and risks in reoperation for coronary artery disease. *Scan J Thorac Cardiovasc Surg* 1990;24:1-6.
- 28 Dougenis D, Naik S, Hedley Brown A. Is repeated coronary surgery for recurrent angina cost effective? *Eur Heart J* 1992;13:9-14.
- 29 Nair UR, Campell CC, Dark JF, Deiranya AK, Lawson RA, Mousalli H, *et al*. Reoperation for recurrent coronary artery and graft disease. A review of 73 patients in a group of 2573 consecutive first operations. *J Cardiovasc Surg* 1989;30:656-60.
- 30 Ivert TS, Ekstrom S, Peterffy A, Welti R. Coronary artery reoperations. Early and late results in 101 patients. *Scand J Thorac Cardiovasc Surg* 1988;22:111-8.
- 31 Warner M, Offord KP, Warner ME, Lennon RL, Conover MA, Jansson-Schumacher U, *et al*. Role of preoperative cessation of smoking and other factors in postoperative pulmonary complications: a blinded prospective study of coronary artery bypass patients. *Mayo Clin Proc* 1989;64:609-16.
- 32 Carrel T, Schmid ER, von Segesser L, Vogt M, Turina M. Preoperative assessment of the likelihood of infection of the lower respiratory tract after cardiac surgery. *Thorac Cardiovasc Surg* 1991;39:85-8.
- 33 Nagachinta T, Stephens M, Reitz B, Polk BF. Risk factors for surgical wound infection following cardiac surgery. *J Infect Dis* 1987;156:967-73.
- 34 Engblom E, Ronnemaas T, Hamalainen H, Kallio V, Vantinen E, Knutz LR. Coronary heart disease risk factors before and after bypass surgery: results of a controlled trial on multifactorial rehabilitation. *Eur Heart J* 1992;13:232-7.
- 35 Brook RH, Park RE, Winslow CM, Chassin MR, Winslow CM, Hampton JR. Diagnosis and treatment of coronary disease: a comparison of doctors attitudes in the USA and the UK. *Lancet* 1988;ii:750-3.
- 36 Ramanathan KB, Vander Zwaag R, Maddock V, Kroetz FW, Sullivan JM, Mirvis DM. Interactive effects of age and other risk factors on long term survival after coronary artery surgery. *J Am Coll Cardiol* 1990;15:1493-9.

Refusing to treat smokers is unethical and a dangerous precedent

Matthew Shiu

Should cigarette smokers be offered coronary artery bypass graft operations? Some specialists are adamant that they should not unless they promise to give up smoking. With increasing resource limitations, such views may soon become accepted policy. Some of the argument would seem well founded on several fronts: the known adverse effect of smoking on the use of anaesthetics; the known adverse effects of continued smoking on graft survival and progression of coronary artery disease; and limited resources with ever increasing demand.

There is no doubt that cigarette smoking is harmful to coronary arteries. The medical profession is united in condemning smoking and has been fairly successful. Working as a general practitioner and with a special interest in cardiology, I spend much of my time getting patients off cigarettes. There is, however, a small group of patients who truly cannot give up their habit. All anti-smoking clinics have such examples. The medical profession does not have a satisfactory treatment for true cigarette addiction. So is it right to completely deny this group of high risk patients a

recognised form of treatment because they cannot give up smoking? According to a spokesperson at the Medical Defence Union, should a case ever be brought to the court it would be difficult to defend such a stand if treatment was refused solely on those grounds. Some patients are denied investigations on the grounds that surgery will not be offered.

The outcome of coronary artery surgery can be compromised by other factors such as hypertension, diabetes, age, female gender, obesity, and hypercholesterolaemia. These are all accepted as deleterious but not absolute overriding contraindications. Is it right to isolate one risk factor and make a firm policy without taking other issues into consideration? Diabetic patients are operated on—even the ones who cannot stay on a strict diet. Will patients be refused surgery because they cannot stick to a cholesterol-lowering diet?

Resources are limited and will continue to be limited. If we examine the resource issue more closely, however, the argument does not seem so strong. The financial cost of not operating is often much higher