risk of suicide, relevant to the next few days or even less. They are at least an invaluable double check on individual clinical assessment. Their presence when risk is judged otherwise to be minimal should make doctors query their original assessment. Surely, for example, we would expect a house physician to think twice, and yet again, before acceding to a request for rapid discharge by an elderly man who has been admitted after a deliberate overdose, especially if he has been misusing alcohol, is depressed, has important physical illness, and has recently suffered a major loss such as bereavement.

Given the limitations of predicting risk factors, how can we help the doctor to assess short term risk? This is central to day to day clinical work and by no means esoteric. Risk factors derived from well conducted follow up studies such as those already described are, of course, important. But clinical audit may help to identify a new range of risk factors by focusing on day to day issues such as problems with behaviour and relationships, which relate more immediately to short term risk.⁵ Clinical audit should also help to define more clearly the

clinical techniques and skills relevant to assessing and managing the risk of suicide. Audit should thus ensure that the right thing is done⁶ and has the potential to reveal anomalies and gaps in clinical practice which, if remedied, should help us meet the targets we have been set. We might then approach these with confidence and enthusiasm.

GETHIN MORGAN
Professor of mental health

Department of Mental Health,
University of Bristol,
Bristol BS2 8DZ

- 1 Department of Health. *The health of the nation: strategy for health in England*. London: HMSO, 1992. (Cm 1986.)
- 2 Nordentoft M, Breum L, Munck LK, Nordestgaard AG, Hunding A, Bjaeldager PAL. High mortality by natural and unnatural causes: a 10 year follow up study of patients admitted to a poisoning treatment centre after suicide attempts. *BMJ* 1993;306:1637-41.
- 3 Hawton K, Fagg J, Platt G, Hawkins M. Factors associated with suicide after parasuicide in young people. *BMJ* 1993;306:1641-4.
- 4 Vassilas CA, Morgan HG. General practitioner contact with suicides. *BMJ* (in press).
- 5 Morgan HG, Priest P. Suicide and other unexpected deaths amongst psychiatric inpatients. The Bristol confidential inquiry. *Br J Psychiatry* 1991;158:368-74.
- 6 Smith R. Audit and research. *BMJ* 1992;305:905-6.

Management at the MRC

Old fashioned and in need of reform

The main thrust of the government's reforms of the NHS was to push management responsibility downwards so that hospitals, units within them, general practices, and community units could be rewarded if they performed well—or punished if they did badly. The aim was to get away from the top down management style of the monolithic NHS. William Waldegrave, who was secretary of state for health on the day the changes were introduced, said at the time: "We've battered away over the past 40 years trying to improve outcomes in essentially a top down way. And when things weren't going well we set up another tier of administration or took one away or changed its boundaries. Ultimately, you can't run an organisation employing nearly a million people like that. You have to give the people down the line the power to improve it themselves."¹

Readers will have their own ideas on whether these reforms have been successful, but in management and economic terms they make perfect sense. Planned economies—like those that operated in the Eastern bloc—are doomed to failure because those doing the planning can never have enough information. Huge companies—like General Motors, IBM, and ICI—have to give real power to those close to customers and make sure that their central organisations add something to the company or they collapse. All of these companies have gone too slowly with the process and are in trouble as a result. Even in small organisations the management trend is to flatten the structure, break down departmental divisions, and reward those who show real initiative and ability to learn and change: otherwise, the organisation fails to adapt quickly enough to the rapidly changing world and is destroyed by its competitors.

The most rapidly changing world of all is that of science and technology, and managing a community of scientists and researchers is one of the most difficult challenges in management. The challenge must be to give power to those close to the research while making sure that they operate within a strategy that will bring benefits to those paying for the research. And the failure of the government's strategy for science, which was introduced last month, is that it stayed with the traditional top down management structure and did

nothing to shift power downwards.^{2,3} Ironically, the strategy was produced by the same minister, William Waldegrave, who was so clear on the necessity for the NHS reforms to give power to those close to the patients.

The measure of that failure is illustrated by two papers in this week's journal, which look at the management structure of the Medical Research Council (MRC) from the point of view of the directors of its smaller units.^{4,5} One of the authors, Roy Gillett, is a researcher who works in one of the units and had to take on most of the management load within the unit. To help him develop his skills he took a master of business administration degree, and his project was to survey the MRC's management of small units. All of the unit directors cooperated with the study, but the central MRC bureaucracy was unhelpful. This in itself is a bad sign: organisations that want to move forward welcome opportunities for critical examination.

The picture that emerges is that the unit directors feel impeded rather than aided by the central MRC bureaucracy. One in eight of the MRC staff works in the head office, which is positioned in an expensive part of central London and does no research, yet the directors feel that they must do most of the managing of the units themselves. Head office staff are rarely seen in the units, apart from during the peer review process, which is so prolonged that it manages to blight almost half of the units' time. The centre is seen by the unit directors as bureaucratic and dictatorial, while they are operating a management style that is oriented towards particular tasks. A survey of the middle managers at the MRC's head office described the organisation as "introspective, secretive, paternalistic, bureaucratic, compartmentalised, lacking a team spirit, perfectionist, slow and amateur in approach to managing."⁴

The authors of the two papers published this week and most of the unit directors would like to see profound changes in the management style of the MRC. The centre should be shrunk and the culture of what remains revitalised. Units should be aided in their management tasks by general managers along the lines of those seen in the best general

practices. Indeed, the best solution to the underlying problems might be to introduce a purchaser-provider model like that introduced into the NHS. The Advisory Council on Science and Technology suggested this to Mr Waldegrave, but he turned it down. With this proposal the MRC (and the other research councils) would become purchasers of research and have no responsibility for managing providers. The units would become providers of research and would compete for contracts from the research councils, the charities, government departments, and other bodies. The remote, heavy handed management style of the councils would become irrelevant, and the units would develop their own management styles and flourish or die. This system would allow what the management guru Tom Peters calls "fast failure,"⁶ which would be overt, swift, and predictable in contrast to the slow, covert, and questionable procedures currently used to close units. Resources could quickly be released for more rewarding research.

Mr Waldegrave seems to recognise the need to shift research management from being top down to bottom up, and

he hopes that this will be achieved through his proposed "foresight procedures"²: in these procedures researchers and others are systematically questioned about research prospects and the information derived is combined with information from marketing people to develop a strategy that is scientifically advanced and fits with what the market needs. But Mr Waldegrave seems to have neglected the effector arm of this process. If the management styles of the research councils are not radically reformed—perhaps to the point of a purchaser-provider split being introduced—then nothing will change.

RICHARD SMITH
Editor

BMJ

- 1 Smith R. William Waldegrave: thinking on the new NHS. *BMJ* 1991;302:636-40.
- 2 Chancellor of the Duchy of Lancaster. *Realising our potential: a strategy for science, engineering, and technology*. London: HMSO, 1993. (Cmnd 2250.)
- 3 Smith R. At last a strategy for British science. *BMJ* 1993;306:1492-3.
- 4 Gillett R, Harrow J. Prescriptions for medical research. I. Management within the Medical Research Council. *BMJ* 1993;306:1668-72.
- 5 Gillett R, Harrow J. Prescriptions for medical research. II. Is medical research well served by peer review? *BMJ* 1993;306:1672-5.
- 6 Peters T. *Thriving on chaos. Handbook for a management revolution*. London: Macmillan, 1986.

Human milk and the preterm baby / |

Mothers should breast feed

Five years ago an editorial on the feeding of preterm infants concluded "Breast not necessarily best."¹ It argued that specialised formulas for low birthweight infants met the high nutrient requirements of babies born prematurely better than human milk. How much does this matter, and what outcome measures are important during this critical period of growth?

Although the terms "feeding" and "nutrition" are often used interchangeably, they are different, particularly where newborn infants are concerned. For example, human milk has antimicrobial and trophic properties, which have been extensively investigated in vitro. Scepticism remains about their relevance to the clinical care of preterm infants despite clear evidence from controlled, prospective studies that even pasteurised human milk significantly reduces the incidence of infection in low birthweight infants at risk.²

It has been asserted that this is important only in developing countries, but a British study of 926 infants weighing < 1850 g at birth has shown that formula fed infants are over six times more likely to develop necrotising enterocolitis than are breast fed babies.³ Both the size and the design of this study (two thirds of the infants were randomised to receive low birthweight formula or banked human milk) suggest that this finding is a true one, and it has been corroborated by a case-control study from Australia.⁴ How human milk protects is unclear: its antimicrobial and anti-inflammatory properties could clearly be relevant, but homologous milk may promote intestinal adaptation more efficiently—as in animals.⁵ Low birthweight infants certainly tolerate human milk more rapidly than formula.⁶

Use of the generic label "human milk" unfortunately blurs the important distinction between a baby's own mother's milk and that donated by mothers of healthy term infants (banked human milk). Babies exclusively fed low birthweight formula undoubtedly grow more quickly than those fed banked human milk,^{7,9} but there is a widespread misconception that this translates into neurodevelopmental advantage. In fact, the mean developmental quotients at 9 months of preterm

babies randomly allocated to exclusive feeding with banked human milk or low birthweight formula did not differ significantly.¹⁰

Although failure to detect a difference was ascribed to lack of statistical power, differences in the developmental outcome at 18 months of a similar sample of babies fed low birthweight formula or standard formula were large and significant.¹¹ This suggests that qualitative differences between human milk and formula influence brain growth or development in infants with somatic undernutrition, a hypothesis supported by the independent advantage in IQ (on the Wechsler intelligence scale for children) of 8.3 points (95% confidence interval 4.9 to 11.7) among preterm infants fed their own mothers' milk.¹² Differences in the content of long chain polyunsaturated fatty acids in human milk and formula may explain these findings, but the possibility that other growth factors are absorbed and exert remote effects cannot be dismissed.¹³ For example, lactoferrin (molecular weight 78 000) is absorbed intact by preterm infants and excreted into the urine.¹⁴ Furthermore, even the most careful multivariate analysis cannot control for unrecognised demographic confounders or intangible differences in parenting associated with breast feeding.

Not surprisingly, evidence that human milk reduces infection and necrotising enterocolitis and enhances the developmental outcome of preterm infants has resulted in a revival of interest in its use. Variability in its composition and the high nutrient demands of preterm infants, particularly for protein, energy, calcium, and phosphorus, can nevertheless present management problems. Commercial human milk fortifiers, containing mainly cows' milk protein hydrolysate, sodium, calcium, and phosphorus, are in vogue, but there is some danger of therapeutic enthusiasm leading to abuse. Small explanatory studies have certainly shown that supplementation of maternal milk can improve the average rate of growth and retention of nitrogen without disturbing metabolic homeostasis,¹⁵ but it should be emphasised that the protein content of milk produced by mothers who deliver prematurely is extremely variable and that the nutritional