Interactions between a general practitioner and representatives of drug companies

Brian O'Mahony

The Surgery, Lismore,
County Waterford,
Republic of Ireland
Brian O'Mahony, general
practitioner

BMJ 1993;306:1649

Nearly all general practitioners meet representatives of drug companies.¹ The drug industry spent an estimated £5000 per general practitioner promoting its products in the United Kingdom in 1985.² Drug representatives are a crucial element in the promotion of products, and expenditure on representatives accounts for over 50% of promotional budgets.¹ Information is available on the frequency and duration of meetings between general practitioners and drug companies' representatives.¹ I describe my meetings with representatives during one year, including the samples, gifts, and literature given to me.

Subjects, methods, and results

The study was based at my singlehanded general practice in a rural town, population 1000, in the south east of Ireland. The practice was not dispensing. For 50 of the 52 weeks between 29 August 1990 and 28 August 1991 I and two receptionists recorded details of the visits by representatives of drug companies to my surgery. We recorded each representative's name and gender, the name of the company represented, the time spent with me, the name and number of products promoted, and any samples, gifts, or literature provided. The representatives were unaware that the study was taking place. For each visit I adopted a passive role, allowing the representative to dictate its content and duration. I did not request any samples but accepted all samples and gifts that were offered. The value of the samples was calculated from the prices quoted for September 1991 in the *Monthly Index of Medical Specialities*.

I received a total of 109 visits, a mean of 2.2 per week, from 49 representatives of 41 drug companies, a mean number of 2.7 visits per company. The visits lasted for a mean of 6.17 (range 2-16) minutes, and the sex of the representatives made no significant difference to the length of the visits ($t=1.63$; $df=107$; $p=0.106$). In each visit a mean of 3.3 (1-6) products were promoted. Data sheets were offered in 11 of the visits whereas advertising brochures were offered in 51. Two representatives offered copies of articles from peer reviewed journals. I received a gift in 64 visits: these usually displayed an advertisement for the product being promoted, and consisted of stationery, pens, diaries, calendars, or trinkets (mugs, tapes, towels, clothes brush, desk tidy, air freshener, ice scraper, and flower seeds). I was given samples of

drugs in 103 of the visits and received 174 samples altogether. The table lists these samples according to the classification used in the *British National Formulary*.

Free drug samples received by a general practitioner in one year. Drugs classified in accordance with British National Formulary

	No (%) of samples	Estimated cost (£)
Drugs used in treatment of:		
Cardiovascular system	36 (21)	376.35
Infections	29 (17)	250.94
Respiratory system	24 (14)	327.09
Gastrointestinal system	21 (12)	155.93
Musculoskeletal and joint diseases	20 (11)	147.46
Skin	12 (7)	66.19
Central nervous system	11 (6)	47.79
Ear, nose, and oropharynx	6 (3)	36.77
Eye	5 (3)	14.54
Others	10 (6)	62.16
Total	174 (100)	1485.22

Comment

The code of marketing practice for the pharmaceutical industry states: "The size and quantity of the sample supplied should be appropriate for either familiarisation with the presentation and appearance of the product, or the provision to patients for immediate use of an initial small supply of a product."³ The total value of the samples recorded in this study was £1485, which constitutes a substantial gift to a general practitioner.

Representatives are perceived by the industry as having a dual role as educators and promoters.¹ During the visits I received, however, the emphasis was on promotion. Advertising has been shown to influence physicians' prescribing habits so that they tend to prescribe analgesics according to advertised guidelines rather than published scientific literature.⁴

The gifts I received were not expensive items. Their main purpose seemed to be an attempt to keep a product's name before me, presumably in the hope that this would be consciously or unconsciously assimilated. Acceptance of gifts can have complex practical and ethical repercussions, including the establishment of a relationship in which a doctor may feel obliged to respond.⁵

I thank Maureen Arrigan for her secretarial help.

- 1 Lexchin J. Doctors and detailers: therapeutic education or pharmaceutical promotion? *Int J Health Serv* 1989;19:663-79.
- 2 Smith R. Doctors and the drug industry; too close for comfort. *BMJ* 1986;293:905-6.
- 3 Federation of Irish Chemical Industries. Code of marketing practice for the pharmaceutical industry (third edition, 1984). In: *Data sheet compendium 1991-92*. Dublin: FICI, 1991:v-ix.
- 4 Avorn J, Chen M, Hartley R. Scientific versus commercial sources of influence on the prescribing behaviour of physicians. *Am J Med* 1982;73:4-8.
- 5 Chren MM, Landefeld CS, Murray TH. Doctor, drug companies and gifts. *JAMA* 1989;262:3448-51.

(Accepted 18 April 1993)

A national survey of the provision of support services for people with cancer

J M Addington-Hall, M W Weir, C Zollman,
M B McIlmurray

As well as technical competence in diagnosis and management people with cancer need psychosocial support.¹ They may also require help to overcome the physical effects of the disease and of treatment.² Although both these needs are increasingly being

recognised, little is known about the distribution of services aimed at helping people cope with cancer. This paper describes the results of a national survey of support services provided by district health authorities for people with cancer.

Methods and results

In January 1991 the district general managers of all 190 English health districts were sent a short postal questionnaire asking about "any rehabilitation or support services provided for patients following treatment or diagnosis of cancer." Four weeks later non-responders were recontacted.

The response rate was 89% (170). Six districts said

Correspondence to:
Dr Addington-Hall.

BMJ 1993;306:1649-50

**Department of
Epidemiology and Public
Health, University College
London, London
WC1E 6EA**

J M Addington-Hall, *lecturer
in health services research*

Wirral Health Authority

M W Weir, *director of
community medicine and
quality assurance*

**Oncology Unit, Bristol
Hospital for Sick Children,
Bristol**

C Zollman, *senior house
officer*

**Royal Lancaster Infirmary,
Lancaster**

M B McIlmurray, *consultant
physician*

they provided no relevant services. Sixteen districts provided only practical support (physiotherapy, dietary advice, stoma and mastectomy care), one provided only emotional care (counselling, support groups), and 94 provided both types of support. Fifty three provided some complementary therapies (guided imagery, meditation/relaxation, healing, and art therapy) in addition to these services. The numbers of districts providing each service are listed in the table.

Although 111 districts had informal links with independently run cancer support services, usually with palliative care services (51) or local cancer support groups (45), there was limited evidence that these services were compensating for deficiencies in NHS services: no districts without support services had links with independent organisations; in four of those offering only practical support emotional care was provided outside the NHS; and in one both emotional care and complementary therapies were available from independent organisations.

Comment

Our results show that there are wide variations in provision of supportive care for patients with cancer. Most districts provided some practical support, usually stoma and mastectomy care. The full range of support, however, was provided in only 32 districts. This may be an underestimation as many hospital dietetics and physiotherapy departments may have provided services for patients with cancer but been overlooked in the completion of the questionnaire because they were not separate specialist cancer services.

Results suggest that the NHS is already responding to the emotional needs of patients with cancer. The availability of counsellors and support groups may, however, be lower than that suggested by our data as we did not ask about the services' nature and accessibility. Some, for example, may have been available only to patients with breast cancer. Some districts at least were referring patients to routinely available clinical psychology or psychiatric services which may not have been available to, or appropriate for, distressed patients with cancer. Macmillan nurses and hospices were often mentioned, suggesting that some services were orientated towards patients in the last stages of their illness.

More districts provided complementary therapies than we had expected. Clearly, the integration of these

Provision of support services for people with cancer by district health authorities

	Districts reporting provision of service	
	No	%
Practical support services		
Specialist physiotherapy	38	22
Dietary advice	116	68
Stoma care	156	92
Mastectomy care	139	82
None of these	7	4
One	10	6
Two	52	31
Three	69	41
Four	32	19
Emotional care		
Counselling	141	83
Support groups	97	57
None of these	24	14
One	54	32
Two	92	54
Complementary therapies		
Guided imagery	18	11
Relaxation-meditation	41	24
Healing	11	7
Art therapy	18	11
None of these	117	69
One	26	15
Two or more	27	16

non-traditional methods into clinical practice has begun, despite the lack of evidence of benefit and of any clear understanding of their role in cancer care.

Our findings suggest that cancer support services are developing in a fragmented and poorly coordinated way. There is no agreed strategy for providing supportive services for patients with cancer and little evidence on which to draw to decide what should be provided and how it should be organised. Further evaluations of models of supportive care¹ and of specific interventions⁴ are urgently needed. In the meantime we urge those responsible for providing and purchasing services for cancer patients to continue to address the need to provide effective and accessible services aimed at meeting physical and emotional needs.

1 Smith T. Cancer services. *BMJ* 1990;301:1406-7.

2 Watson PG. Cancer rehabilitation: the evolution of a concept. *Cancer Nursing* 1990;13:2-12.

3 McIlmurray MB, Gorst DW, Holdcroft PE. A comprehensive service for patients with cancer in a district general hospital. *BMJ* 1986;292:669-71.

4 Fallowfield L, Roberts R. Cancer counselling in the United Kingdom. *Psychology and Health* 1992;6:107-17.

(Accepted 15 April 1993)

Relation between sialic acid concentrations and the haemostatic system in the elderly

Kazuomi Kario, Takefumi Matsuo

**Department of Internal
Medicine, Awaji-Hokutan
Public Clinic, Hokutan,
Hyogo 656-16, Japan**
Kazuomi Kario, *physician*

**Department of Internal
Medicine, Hyogo
Prefectural Awaji Hospital,
Sumoto, Hyogo 656, Japan**
Takefumi Matsuo, *physician*

Correspondence to:
Dr K Kario.

BMJ 1993;306:1650-1

A recent prospective study showed that mortality due to cardiovascular disease was increased in people with high serum sialic acid concentrations.¹ Recent reports suggest that smoking and lipid concentrations may be confounders,^{2,3} but the precise relation between the serum sialic acid concentrations and death from cardiovascular disease remains uncertain. Fibrinogen is a well known risk factor for cardiovascular disease and, like sialic acid, is an acute phase reactant. Plasma fibrinogen concentrations increase with aging and smoking,⁴ characteristics which are similar to those of sialic acid.^{2,3} The possibility that fibrinogen might be a strong confounder that explains the relation between sialic acid concentrations and cardiovascular death has

not been investigated. We therefore investigated the relation between serum sialic acid concentrations and plasma concentrations of various haemostatic variables and the effect of smoking.

Subjects, methods, and results

We studied 138 healthy Japanese men aged over 60 years (mean age 78 years) who were selected from among men undergoing annual health examinations. Subjects with abnormal results on routine laboratory tests (C reactive protein, blood urea nitrogen, creatinine, transaminases, etc) were excluded. Fifty subjects had hypertension and 48 were smokers. Blood samples were collected after an overnight fast, and serum sialic acid concentrations were measured by an enzymatic assay.⁵ Serum total cholesterol and triglyceride concentrations were determined by enzymatic assays using cholesterol esterase, cholesterol oxidase, and glycerol-3-phosphate oxidase. The clotting activity of plasma fibrinogen and factor VII was determined by automated one-stage clotting assays. Plasma concentrations of antithrombin III, heparin cofactor II, plasminogen,